

Barcode No **89781007**
Patient Name **B/OAMANDEEP KOUR**
Age/Sex **3 DAYS(S)/Male**
Referred By **DR. NICU**

Lab No **16952412190001**
Reg Date **19/Dec/2024 04:27PM**
Sample Coll. Date **19/Dec/2024 03:30 PM**
Sample Rec.Date **20/Dec/2024 07:05 AM**

Report Date **20/Dec/2024 11:25AM**

HAEMATOTOLOGY

Test Name With Methodology	Result	Unit	Biological Ref.Interval
G6PD Deficiency (Quantitative)			
G6PD (Quantitative)	13.60	U/g Hb	>2.8

EDTA Whole Blood, Spectrophotometry

CUT-OFF VALUES

Sex/G6PD Statu	U/g Hb
Male/G6PD deficiency	<2.8
Male/G6PD normal	≥2.8
Female/G6PD deficiency	<2.8
Female/G6PD intermediate	2.8 – <7.4
Female/G6PD normal	≥7.4

Comment:

The goal of G6PD activity testing is to detect a G6PD deficiency and to determine its potential severity. It is primarily ordered on male children as they are the most frequently affected but may also be ordered on female children if the doctor suspects that a G6PD deficiency may be present. G6PD activity testing may be ordered on children who had persistent jaundice as a newborn that was not due to another identified cause. It may also be ordered on patients of any age who have had one or more episodes of hemolytic anemia.

If there is decreased G6PD activity, there is some degree of G6PD deficiency. In general, the lower the activity level, the more likely the patient is to experience symptoms when exposed to an oxidative stress. The results, however, cannot be used to predict how an affected patient will react in a given set of circumstances. The severity of symptoms will vary from patient to patient and from episode to episode.

If a male patient has normal G6PD activity levels, it is likely that they do not have a deficiency. However, if the test was performed during an episode of hemolytic anemia, it should be repeated a few weeks later when the RBC population has had time to replenish and mature.

Heterozygous females will have both G6PD-deficient and non-deficient cells. They will usually have normal or near normal G6PD activity levels, and few will experience any symptoms. A carrier may not be able to be detected through G6PD activity testing. In the rare homozygous female, G6PD activity will be significantly and detectably decreased.


Dr. Urvashi, DNB, Path
(Consultant Pathologist)
Reg. No. DMC-84904

Dr. Prashant Goyal (DCP)
(Director & Chief Pathologist)
Reg. No. DMC-53016

Page 1 of 1

A Unit of Hexamed Healthcare Services LLP

Corporate Office: Flat no. 203, Sagacity heights, Plot No. 27, Kavuri Hills Road, CBI Colony, Madhapur, Telangana 500033

Jammu Address: 25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 📞 support@hexamedhealthcare.com

☎ +91 95419 01145

☎ 1800 212 6547

🌐 www.hexamedhealthcare.com



Scan QR for Location



Scan QR for Enquiry Form