

Name : Mr. SHUBHAM
 Lab No. : 463577644
 Ref By : DR ADEEB SINGH
 Collected : 24/1/2025 1:40:00PM
 A/c Status : P
 Collected at : Hexamed Healthcare Services LLP
 25-270/6, Taluka Bahu, Main Bypass Road, NH1A,
 Channi Himmat,
 Jammu

Age : 29 Years
 Gender : Male
 Reported : 28/1/2025 12:26:15PM
 Report Status : Final

Test Report

STONE ANALYSIS BY FTIR



Sample: Kidney Stones

Aggregate Weight: 1.920 gm

Composition : Calcium Oxalate Monohydrate 80% Calcium Oxalate Dihydrate 20%

(Representative stone analysed)

Kidney stone (Nephrolithiasis) develops when urine becomes supersaturated with crystal forming substances such as Calcium, uric acid, oxalate etc.(resulting from dehydration or genetic predisposition) or due to decrease in urinary concentration of substances such as magnesium, citrate etc. that inhibit stone formation. Kidney stones tend to recur; approximately 50% of patients who form one stone form another in 10 years. Nephrolithiasis is considered to be a disease of affluence because it is more prevalent in conditions like Obesity, Hypertension & Diabetes. For patients with multiple kidney stones 24-hour urine chemistries should be measured to identify abnormalities that predispose to kidney stones, which guides dietary and

Page 1 of 4

A Unit of Hexamed Healthcare Services LLP

Corporate Office: Flat no. 203, Sagacity heights, Plot No. 27, Kavuri Hills Road, CBI Colony, Madhapur, Telangana 500033

Jammu Address: 25-270/6, Main Bypass Road, NH 1A, Channi Rama, Jammu, J & K 180015

feedback@hexamedhealthcare.com enquiry@hexamedhealthcare.com support@hexamedhealthcare.com



Scan QR for Location



Scan QR for Enquiry Form

Name : MR. SUDHAKAR
 Lab No. : 463577644
 Ref By : DR ADEEB SINGH
 Collected : 24/1/2025 1:40:00PM
 A/c Status : P
 Collected at : Hexamed Healthcare Services LLP
 26-270/6, Taluka Bahu, Main Bypass Road, NH1A,
 Channi Himmat,
 Jammu

Age : 29 Years
 Gender : Male
 Reported : 28/1/2025 12:26:15PM
 Report Status : Final

pharmacologic therapy to prevent future stone events.

STONE TYPE	FREQUENCY (%)	CAUSES
Calcium oxalate / mix	75	- Hypercalciuria with Hypercalcemia (Hyperparathyroidism, Vitamin D excess, malignancy) - Idiopathic hypercalciuria - Hypocitraturia - Alkaline urine, low urine volume - High dose of vitamin C - Metabolic syndrome - Inflammatory bowel disease
Calcium Phosphate	5	- Distal Renal Tubular acidosis - Primary Hyperparathyroidism - Chronic urinary tract infection - Hypercalciuria and/or Hypophosphaturia
Magnesium Ammonium Phosphate (Struvite)	10-20	Recurrent urinary tract infection
Uric Acid	5-10	- Hyperuricosuria and /hyperuricosemia - Metabolic syndrome (Diabetes, obesity, Hypertension) - Acidic urine (pH<5.5) - Gout
Cystine	1	Cystinuria (A genetic disorder)

DIETARY RECOMMENDATIONS FOR PREVENTION OF RECURRENCE

General Dietary measures:

- Fluid intake should be between 2.5-3L/day or it should be sufficient to produce urine output of 2L/day
- Diet rich in fibres, fruits (orange, lemon, apple etc) & vegetables
- Maintain adequate calcium intake (1000-1200 mg/day)
- Aim to achieve/maintain healthy weight
- Reduce intake of animal protein (meat & dairy)

Specific measures based on metabolic evaluation and stone composition:


STONE COMPONENT	METABOLIC PROFILE	SUGGESTED DIETARY RECOMMENDATION
Calcium oxalate or combined calcium oxalate/phosphate stone	Hypercalciuria	Salt restriction
	Hyperoxaluria	Calcium supplements with meal Restrict items rich in oxalate eg



Lab No. : 463577644
 Ref By : DR ADEEB SINGH
 Collected : 24/1/2025 1:40:00PM
 A/c Status : P
 Collected at : Hexamed Healthcare Services LLP
 25-270/6, Taluka Bahu, Main Bypass Road, NH1A,
 Channi Himmat,
 Jammu

Age : 29 Years
 Gender : Male
 Reported : 28/1/2025 12:26:15PM
 Report Status : Final

Uric Acid		Soya protein, Spinach, Nuts, Strawberries, Cocoa, Chocolates, beetroot, carrot, radish etc
	Hyperuricosuria	- Limit animal protein & purine intake (Fish, lamb, beef, chicken, pork etc), cauliflower, Mushroom - Reduce alcohol intake
	Hypocitraturia	Increase citrate rich food (Orange, lemon, apple etc)
	Hyperuricosuria & Hyperuricemia	- Limit animal protein & purine intake (Fish, lamb, beef, chicken, pork etc), cauliflower, Mushroom - Reduce alcohol intake
Cystine	Metabolic syndrome	Lifestyle changes, weight loss, Diabetes control
		- Increase fluid intake to 3-4L, Urine output should at least be 3L - Restrict sodium & protein intake


 DMC - 67327

 Dr Anjalika Goyal
 MD, Biochemistry
 Consultant Biochemist


 DMC - 89819

 Dr Himangshu Mazumdar
 MD, Biochemistry
 Sr. Consultant Biochemist


 DMC - 9550

 Dr Nimmi Kansal
 MD, Biochemistry
 Technical Director - Clinical Chemistry
 & Biochemical Genetics


 MCI 15-19066

 Dr. Richa Sirohi
 MD, Biochemistry
 Sr. Consultant Biochemist

End of report



Scan QR for Location

A Unit of Hexamed Healthcare Services LLP Page 3 of 4

Corporate Office: Flat no. 203, Sagacity heights, Plot No. 27, Kavuri Hills Road, CBI Colony, Madhapur, Telangana 500033

Jammu Address: 25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com support@hexamedhealthcare.com

+91 95419 01145

1800 212 6547

www.hexamedhealthcare.com



Scan QR for Enquiry Form