

Barcode No	87943858	Lab No	16932501270003
Patient Name	Mrs.PABLEEN KOUR	Reg Date	27/Jan/2025 10:25AM
Age/Sex	24 YRS/Female	Sample Coll. Date	27/Jan/2025 09:30 AM
Referred By	DR. NITIKA GANDOTRA	Sample Rec.Date	27/Jan/2025 12:21 PM
		Report Date	27/Jan/2025 01:50PM


.IMMUNO BIOCHEMISTRY-1

Test Name With Methodology	Result	Unit	Biological Ref.Interval
Insulin Fasting			
Insulin Fasting	5.84	uIU/mL	2.6-24.9

Serum, Electro Chemi Luminescent Immuno Assay

Comment:

This test is used to estimate the levels of Insulin in serum. This may be used to diagnose insulinomas and to help diagnose acute or chronic (fasting) hypoglycemia. Insulin levels may also be used to monitor endogenous insulin, check for insulin resistance, and to help determine when a type 2 diabetic might need to start taking insulin injections to supplement oral medications.


Dr. Prashant Goyal (DCP)
{Director & Chief Pathologist}
Reg. No. DMC-53016**A Unit of Hexamed Healthcare Services LLP****Corporate Office:** Flat no. 203, Sagacity heights, Plot No. 27, Kavuri Hills Road, CBI Colony, Madhapur, Telangana 500033**Jammu Address:** 25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 🌐 support@hexamedhealthcare.com






Barcode No	87943858	Lab No	16952501270003
Patient Name	Mrs.PABLEEN KOUR	Reg Date	27/Jan/2025 10:25 AM
Age/Sex	24 YRS/Female	Sample Coll. Date	27/Jan/2025 09:30 AM
Referred By	DR. NITIKA GANDOTRA	Sample Rec.Date	27/Jan/2025 12:20 PM
Ref. Lab/Hosp		Report Date	27/Jan/2025 01:34PM


Test Name With Methodology	Result	Unit	Biological Ref.Interval
Testosterone- Free			
Free Testosterone Serum, CLIA	2.16	pg/mL	0.5-4.2

SUMMARY AND EXPLANATION OF THE TEST

Testosterone is a male sex hormone, secreted by leydig or interstitial cells of the testes. The amount of testosterone synthesized is regulated and controlled through negative feedback on the hypothalamus and pituitary gland by the pituitary hormone and luteinizing hormone (LH). Testosterone circulating in the blood binds to three proteins: sex hormone binding globulin (SHBG, 60-80%), albumin and cortisol binding globulin'. About 1-2% of the total circulating testosterone remains unbound or free. Only measurement of free testosterone permits the estimating of the biologically active hormone. Free testosterone determinations are recommended to overcome the influences caused by variations in transport proteins on the total testosterone concentration SHBG-bound testosterone remains in the circulation with no binding function in this form. Higher testosterone and lower SHBG levels can increase free testosterone. Free testosterone can be diagnostically useful when testosterone does not correspond with the clinical presentation of hypogonadism, especially in aging men with borderline low levels of testosterone and men in whom levels of SHBG are suspected to be altered?

Measuring free testosterone is an indirect way to measure a far more important health indicator-the level of SHBG in the bloodstream. Free testosterone measurements have been very helpful in monitoring Alzheimer's disease for older men and women. Lower levels of free testosterone may have a higher risk of developing Alzheimer's disease.


Dr. Anuja Adarsh, MBBS, MD
(Consultant Biochemist)
Reg. No. DMC-90705


Dr. Prashant Goyal (DCP)
(Director & Chief Pathologist)
Reg. No. DMC-53016

Page 2 of 2

A Unit of Hexamed Healthcare Services LLP

Corporate Office: Flat no. 203, Sagacity heights, Plot No. 27, Kavuri Hills Road, CBI Colony, Madhapur, Telangana 500033

Jammu Address: 25-270/6, Main Bypass Road, NH-1A, Channi Rama, Jammu, J & K 180015

✉ feedback@hexamedhealthcare.com @ enquiry@hexamedhealthcare.com 🌐 support@hexamedhealthcare.com



Scan QR for Location

📞 +91 95419 01145

📞 1800 212 6547

🌐 www.hexamedhealthcare.com



Scan QR for Enquiry Form