

Two-Way Example as Factorial (Easy Way)

This example is for illustration! This data represents 2 Varieties and 3 Tillage methods for a total of 6 Treatment combinations.

```
library(car)
library(emmeans)
InData <- read.csv("~/Dropbox/STAT512/Lectures/ExpDesign2/ED2_2wayData.csv")
str(InData)
```

```
## 'data.frame':    24 obs. of  4 variables:
## $ trt : int  1 2 3 4 5 6 1 2 3 4 ...
## $ till: int  1 2 3 1 2 3 1 2 3 1 ...
## $ var : int  1 1 1 2 2 2 1 1 1 2 ...
## $ resp: num  9.2 4.1 4.1 7.3 5.1 8.2 8.1 6.8 6.1 6.1 ...
```

3 x 2 x 4 reps

```
table(InData$var, InData$till)
```


```
##
##      1 2 3
## 1 4 4 4
## 2 4 4 4
```

✓ widely
4 reps per trt combo

#Important: Need to define till, var as.factors!

```
InData$till <- as.factor(InData$till)
InData$var <- as.factor(InData$var)
with(interaction.plot(till, var, resp), data = InData)
```

Factors



```
aggregate(resp ~ till*var, data = InData, FUN = mean)
```

```
## till var resp
## 1    1    1 7.975
## 2    2    1 4.550
```

```
## 3    3    1 4.275
## 4    1    2 6.475
## 5    2    2 4.675
## 6    3    2 4.425
```

For Illustration: Comparing Type 1, 2, 3 ANOVA tables with Default Contrasts

The default contrasts in R do not sum to zero. Hence the Type3 tests are meaningless! For illustration, we look at the coefficients table from the `summary()` output as well as the `model.matrix`. But this information is not usually required to address typical research questions!

```
Model1 <- lm(resp ~ till*var, data = InData)
anova(Model1)
```

```
## Analysis of Variance Table
##
## Response: resp
##           Df Sum Sq Mean Sq F value    Pr(>F)
## till       2 40.426  20.2129   4.6167 0.02407 *
## var        1  1.000   1.0004   0.2285 0.63839
## till:var    2  3.576   1.7879   0.4084 0.67074
## Residuals 18 78.808   4.3782
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Handwritten notes:
 - given till in model
 - given till & var

```
Anova(Model1, type = 2)
```

```
## Anova Table (Type II tests)
##
## Response: resp
##           Sum Sq Df F value    Pr(>F)
## till       40.426  2   4.6167 0.02407 *
## var         1.000  1   0.2285 0.63839
## till:var    3.576  2   0.4084 0.67074
## Residuals 78.807 18
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Handwritten notes:
 - given var Type 1 = Type 2
 - given till

```
Anova(Model1, type = 3)
```

```
## Anova Table (Type III tests)
##
## Response: resp
##           Sum Sq Df F value    Pr(>F)
## (Intercept) 254.402  1 58.1067 4.842e-07 ***
## till        33.995  2   3.8823  0.03965 *
## var         4.500  1   1.0278  0.32411
## till:var    3.576  2   0.4084  0.67074
## Residuals   78.808 18
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Handwritten notes:
 - given var, var & till interaction
 - Type 3 ≠ Type 1

```
summary(Model1)
```

```
##
## Call:
## lm(formula = resp ~ till * var, data = InData)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.175 -1.188 -0.025  1.275  3.775
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    7.975      1.046   7.623 4.84e-07 ***
## till2         -3.425      1.480  -2.315  0.0326 *
## till3         -3.700      1.480  -2.501  0.0223 *
## var2          -1.500      1.480  -1.014  0.3241
## till2:var2     1.625      2.092   0.777  0.4475
## till3:var2     1.650      2.092   0.789  0.4406
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.092 on 18 degrees of freedom
## Multiple R-squared:  0.3635, Adjusted R-squared:  0.1867
## F-statistic: 2.056 on 5 and 18 DF, p-value: 0.1189
```

```
model.matrix(Model1)
```

```
##      (Intercept) till2 till3 var2 till2:var2 till3:var2
## 1              1      0      0      0              0
## 2              1      1      0      0              0
## 3              1      0      1      0              0
## 4              1      0      0      1              0
## 5              1      1      0      1              1
## 6              1      0      1      1              0
## 7              1      0      0      0              0
## 8              1      1      0      0              0
## 9              1      0      1      0              0
## 10             1      0      0      1              0
## 11             1      1      0      1              1
## 12             1      0      1      1              0
## 13             1      0      0      0              0
## 14             1      1      0      0              0
## 15             1      0      1      0              0
## 16             1      0      0      1              0
## 17             1      1      0      1              1
## 18             1      0      1      1              0
## 19             1      0      0      0              0
## 20             1      1      0      0              0
## 21             1      0      1      0              0
## 22             1      0      0      1              0
## 23             1      1      0      1              1
## 24             1      0      1      1              0
## attr("assign")
## [1] 0 1 1 2 3 3
```

```
## attr("contrasts")
## attr("contrasts")$till
## [1] "contr.treatment"
##
## attr("contrasts")$var
## [1] "contr.treatment"
```

```
getOption("contrasts")
```

```
## unordered ordered
## "contr.treatment" "contr.poly"
contrasts(InData$till)
```

```
## 2 3
## 1 0 0
## 2 1 0
## 3 0 1
```

default R parameterisation for categorical

not important (for cont)

default

Standard Analysis *

It is necessary to choose a contrasts setting that sums to zero (not the default used by R). For routine analysis, we are typically interested in the Type 3 tests and pairwise comparisons from emmeans. Now the Type3 ANOVA table matches the Type1 and 2 tables above (due to balance). Note: The warning from emmeans ("Results may be misleading due to involvement in interactions") will be displayed when considering comparisons of main effects in any model that includes an interaction.

```
options(contrasts = c("contr.sum", "contr.poly"))
contrasts(InData$till)
```

```
## [,1] [,2]
## 1 1 0
## 2 0 1
## 3 -1 -1
```

sum-to-zero parameterization

```
Model2 <- lm(resp ~ till*var, data = InData)
Anova(Model2, type = 3)
```

see slide 60

```
## Anova Table (Type III tests)
```

```
##
```

```
## Response: resp
```

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	698.76	1	159.6001	2.195e-10 ***
till	40.43	2	4.6167	0.02407 *
var	1.00	1	0.2285	0.63839
till:var	3.58	2	0.4084	0.67074
Residuals	78.81	18		

this

```
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
emmeans(Model2, pairwise ~ till)
```

NOTE: Results may be misleading due to involvement in interactions

```
## $emmeans
```

	till	emmean	SE	df	lower.CL	upper.CL
##	1	7.2250	0.7397799	18	5.67078	8.77922

```
## 2 4.6125 0.7397799 18 3.05828 6.16672
## 3 4.3500 0.7397799 18 2.79578 5.90422
##
## Results are averaged over the levels of: var
## Confidence level used: 0.95
##
## $contrasts
## contrast estimate SE df t.ratio p.value
## 1 - 2 2.6125 1.046207 18 2.497 0.0557
## 1 - 3 2.8750 1.046207 18 2.748 0.0337
## 2 - 3 0.2625 1.046207 18 0.251 0.9660
##
## Results are averaged over the levels of: var
## P value adjustment: tukey method for comparing a family of 3 estimates
```

```
emmeans(Model2, pairwise ~ var)
```

```
## NOTE: Results may be misleading due to involvement in interactions
```

```
## $emmeans
## var emmean SE df lower.CL upper.CL
## 1 5.600000 0.6040278 18 4.330985 6.869015
## 2 5.191667 0.6040278 18 3.922651 6.460682
##
## Results are averaged over the levels of: till
## Confidence level used: 0.95
##
## $contrasts
## contrast estimate SE df t.ratio p.value
## 1 - 2 0.4083333 0.8542243 18 0.478 0.6384
##
## Results are averaged over the levels of: till
```

```
emmeans(Model2, pairwise ~ till:var)
```

```
## $emmeans
## till var emmean SE df lower.CL upper.CL
## 1 1 7.975 1.046207 18 5.777001 10.172999
## 2 1 4.550 1.046207 18 2.352001 6.747999
## 3 1 4.275 1.046207 18 2.077001 6.472999
## 1 2 6.475 1.046207 18 4.277001 8.672999
## 2 2 4.675 1.046207 18 2.477001 6.872999
## 3 2 4.425 1.046207 18 2.227001 6.622999
##
```

```
## Confidence level used: 0.95
```

```
##
## $contrasts
## contrast estimate SE df t.ratio p.value
## 1,1 - 2,1 3.425 1.47956 18 2.315 0.2385
## 1,1 - 3,1 3.700 1.47956 18 2.501 0.1756
## 1,1 - 1,2 1.500 1.47956 18 1.014 0.9073
## 1,1 - 2,2 3.300 1.47956 18 2.230 0.2721
## 1,1 - 3,2 3.550 1.47956 18 2.399 0.2080
## 2,1 - 3,1 0.275 1.47956 18 0.186 1.0000
## 2,1 - 1,2 -1.925 1.47956 18 -1.301 0.7808
## 2,1 - 2,2 -0.125 1.47956 18 -0.084 1.0000
```

borderline

significant

FTR from ANOVA

not significant interaction

matches 6 trts
one-way ANOVA

15 pairwise
from one way

```

## 2,1 - 3,2    0.125 1.47956 18    0.084 1.0000
## 3,1 - 1,2   -2.200 1.47956 18   -1.487 0.6763
## 3,1 - 2,2   -0.400 1.47956 18   -0.270 0.9998
## 3,1 - 3,2   -0.150 1.47956 18   -0.101 1.0000
## 1,2 - 2,2    1.800 1.47956 18    1.217 0.8233
## 1,2 - 3,2    2.050 1.47956 18    1.386 0.7348
## 2,2 - 3,2    0.250 1.47956 18    0.169 1.0000
##
## P value adjustment: tukey method for comparing a family of 6 estimates

```