



Started on	Thursday, 1 October 2026, 8:40 AM
State	Finished
Completed on	Thursday, 1 October 2026, 8:43 AM
Time taken	3 mins 24 secs
Marks	1.00/1.00
Grade	10.00 out of 10.00 (100%)

Question 1 | Correct | Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2  int main() {
3      int n, k, l = 0, r = 1;
4      if (scanf("%d", &n) != 1) return 0;
5      int a[n];
6      for (int i = 0; i < n; i++) scanf("%d", &a[i]);
7      if (scanf("%d", &k) != 1) return 0;
8      while (l < n && r < n) {
9          if (l != r && a[r] - a[l] == k) return printf("1\n"), 0;
10         if (a[r] - a[l] < k) r++; else l++;
11     }
12     return printf("0\n"), 0;
13 }
14

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓

	Input	Expected	Got	
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

Back to Course