

resourcefulness. Of our sample, 53.8% (N=184) reported belonging to a support group. The majority of participants (138) belonged to online support groups on Facebook, 41 reported participating in in-person support groups, 3 belonged to both in-person and online support groups, and 8 did not respond. When asked whether and how well our participants knew other families like theirs, 31% (106) said they did not know any, 40.6% (139) knew of at least one, but not well, and 28.4% (97) said they had at least one friend with a family like her own. Knowing other families like their own affected self-appraised stress, but no other outcomes. Grandmothers participating in support groups had lower mindfulness scores, higher stress, and worse mental health scores than grandmothers not participating in support groups, possibly reflecting the higher need for support among those participants.

THE RELATIONSHIP BETWEEN SOCIAL SUPPORT, AGING EXPECTATIONS, AND HEALTH BEHAVIOR IN MIDDLE AGE

Hyun-E Yeom,¹ Misook Jung,² and Eunyong Park,¹

1. *Chungnam National University, Daejeon, Taejon-jikhalsi, Republic of Korea*, 2. *Chungnam National University, Daejeon, Republic of Korea*

Midlife is a critical period when individuals need to actively engage in healthy behaviors for healthy aging. Although both social relations and attitudes toward aging are factors related to health behavior, little is known about their relationships based on age-related differences. The purposes of this study were to investigate the influence of social support affecting health behavior through expectations regarding aging and to examine how age affects the relationship. A cross-sectional study was conducted with data from 245 midlife Koreans (mean age= 51.5) collected by a self-administered survey. Data were analyzed using the PROCESS macro in SPSS. Social support was significantly related to expectations regarding aging ($r = -.135$, $p = .034$) and health behavior ($r = .223$, $p < .000$). There was age-related difference in the relationship between social support and expectations regarding aging ($\beta = .007$, $p = .038$), indicating that the relationship was much stronger in the younger group. In addition, the influence of social support on health behavior through expectations regarding aging was significant in relatively young middle-aged individuals. Our findings emphasize the importance of supportive social relationships, which could affect expectations regarding aging linked to health behavior, especially for young middle-aged individuals. It is necessary to develop psycho-cognitive programs to activate social interaction and to improve positive attitudes toward aging for more active engagement in health behaviors in mid-life individuals.

SESSION 2956 (POSTER)

TECHNOLOGY II

A SYSTEMATIC REVIEW OF IN-HOME SMART TECHNOLOGY ADOPTION TO IMPROVE OLDER ADULT HEALTH AND FAMILY CAREGIVING

Geunhye Park, and Erin Robinson, *University of Missouri-Columbia, Columbia, Missouri, United States*

In-home and internet-based smart technologies to improve older adult health has been rapidly developing. Technologies such as in-home sensors and smart homes enable older adults to live independently and age in place. These technologies also assist informal caregivers in their roles, thus reducing caregiver burden. However, technology adoption among older adults and family caregivers has been relatively low and reasons for technology acceptance are complex. Therefore, the purpose of this study was to conduct a systematic review of the literature, examining acceptance and adoption of in-home, internet-based smart technologies that are designed to improve health outcomes of older adults and can assist family caregivers in providing supports. This study utilized the Rew method (2011) and included peer-reviewed research articles published between 1991 and 2019 and available in: ISI Web of Science; PubMed; Scopus; CINAHL; and PsycInfo. A total of 1,227 relevant articles were identified with the search strings used and a final sample of 48 articles were included after the title, abstract, and full article review processes. Findings highlight several facilitators and barriers to technology adoption. Some facilitators to adoption include: technology familiarity, safety/security, personally tailored, non-obtrusive design, easy access, and reduction of caregiver burden. A few barriers include: cost, difficulty to use, time, stigma, privacy, data accuracy, and confidence. Additional findings will also be presented. A more thorough understanding of these facilitators and barriers to acceptance/adoption is crucial for the successful dissemination of in-home, internet-based smart technologies. Increased adoption can improve older adult health and reduce caregiver burden.

ADDRESSING THE ROLE OF SMART ROBOTIC HEALTH ASSISTANTS WITHIN THE HUMAN-MACHINE FRONTIER OF GERIATRIC HEALTHCARE

Nadia Firdausya,¹ Alex Bishop,² Barbara Carlson,³ and Weihua Sheng,² 1. *Oklahoma State University, Stillwater, Oklahoma, United States*, 2. *Oklahoma State University, Stillwater, Oklahoma, United States*, 3. *University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, United States*

Data for this study was acquired from three separate stakeholder focus group sessions involving nurse case managers ($n = 5$), social agency caseworkers ($n = 5$), and rural outreach providers ($n = 5$). Participants across all groups were asked to address the question: "When it comes to your work, what would you want a smart robot assistant to do for you?" Data from the three sessions were combined, transcribed verbatim, coded, and analyzed for thematic content. Three shared themes emerged, including health monitoring, behavioral intervention, and healthcare literacy. Relative to health monitoring, participants desired a robot that possessed functions in the form of "taking vital signs," and "tracking water and food intake." There was also a thematic agreement regarding behavioral intervention capabilities. Most notably, advisory stakeholders acknowledged a need for a smart robotic assistant to provide geriatric care recipients with "an alert or reminder to take medication." This was viewed as an essential intervention for improving medication adherence. Healthcare literacy emerged as a final theme among advisory groups. In particular, participants noted that a smart robot