

CP214: Discrete Structures

Counting

Dr. Zia Ud Din

Outline

- Basic Counting Principles
- Inclusion-Exclusion Principle
- The Pigeonhole Principle
- Permutations and Combinations
- Pascal's Triangle and Binomial Coefficients

Basic Counting Principles

- **Counting problems** are of the following kind:
 - “**How many** different 8-letter passwords are there?”
 - “**How many** possible ways are there to pick 11 soccer players out of a 20-player team?”
- In addition, counting is the basis for computing **probabilities of discrete events**.
 - “What is the probability of winning the lottery?”

Basic Counting Principles

The sum rule:

- If a task can be done in n_1 ways and a second task in n_2 ways, and if these two tasks cannot be done at the same time, then there are $n_1 + n_2$ ways to do either task.

Example:

- The department will award a free computer to either a CS student or a CS professor. How many different choices are there, if there are 530 students and 15 professors?
- There are $530 + 15 = 545$ choices.

Basic Counting Principles

Generalized sum rule:

- If we have tasks T_1, T_2, \dots, T_m that can be done in n_1, n_2, \dots, n_m ways, respectively, and no two of these tasks can be done at the same time, then there are $n_1 + n_2 + \dots + n_m$ ways to do one of these tasks.

Basic Counting Principles

The product rule:

- Suppose that a procedure can be broken down into two successive tasks. If there are n_1 ways to do the first task and n_2 ways to do the second task **after the first task has been done**, then there are $n_1 n_2$ ways to do the procedure.

Generalized product rule:

- If we have a procedure consisting of sequential tasks T_1, T_2, \dots, T_m that can be done in n_1, n_2, \dots, n_m ways, respectively, then there are $n_1 \cdot n_2 \cdot \dots \cdot n_m$ ways to carry out the procedure.

Basic Counting Principles

Example:

- How many different license plates are there that contain exactly three English letters ?

Solution:

- There are 26 possibilities to pick the first letter, then 26 possibilities for the second one, and 26 for the last one.
- So, there are $26 \cdot 26 \cdot 26 = 17,576$ different license plates.

Basic Counting Principles

- The sum and product rules can also be phrased in terms of **set theory**.
- **Sum rule:** Let A_1, A_2, \dots, A_m be disjoint sets. Then the number of ways to choose any element from one of these sets is $|A_1 \cup A_2 \cup \dots \cup A_m| = |A_1| + |A_2| + \dots + |A_m|$.
- **Product rule:** Let A_1, A_2, \dots, A_m be finite sets. Then the number of ways to choose one element from each set in the order A_1, A_2, \dots, A_m is $|A_1 \times A_2 \times \dots \times A_m| = |A_1| \cdot |A_2| \cdot \dots \cdot |A_m|$.

Inclusion-Exclusion Principle

- How many bit strings of length 8 either start with a 1 or end with 00?
- **Task 1:** Construct a string of length 8 that starts with a 1.
There is one way to pick the first bit (1),
two ways to pick the second bit (0 or 1),
two ways to pick the third bit (0 or 1),
.
.
.
two ways to pick the eighth bit (0 or 1).
- **Product rule:** Task 1 can be done in $1 \cdot 2^7 = 128$ ways.

Inclusion-Exclusion Principle

□ **Task 2:** Construct a string of length 8 that ends with 00.

There are two ways to pick the first bit (0 or 1),
two ways to pick the second bit (0 or 1),

·
·
·

two ways to pick the sixth bit (0 or 1),
one way to pick the seventh bit (0), and
one way to pick the eighth bit (0).

□ **Product rule:** Task 2 can be done in $2^6 = 64$ ways.

Inclusion-Exclusion Principle

- Since there are 128 ways to do Task 1 and 64 ways to do Task 2, does this mean that there are 192 bit strings either starting with 1 or ending with 00 ?
- No, because here Task 1 and Task 2 can be done **at the same time**.
- When we carry out Task 1 and create strings starting with 1, some of these strings end with 00.
- Therefore, we sometimes do Tasks 1 and 2 at the same time, so **the sum rule does not apply**.

Inclusion-Exclusion Principle

- If we want to use the sum rule in such a case, we have to subtract the cases when Tasks 1 and 2 are done at the same time.
- How many cases are there, that is, how many strings start with 1 **and** end with 00?

There is one way to pick the first bit (1),
two ways for the second, ..., sixth bit (0 or 1),
one way for the seventh, eighth bit (0).


- **Product rule:** In $2^5 = 32$ cases, Tasks 1 and 2 are carried out at the same time.

Inclusion-Exclusion Principle

- Since there are 128 ways to complete Task 1 and 64 ways to complete Task 2, and in 32 of these cases Tasks 1 and 2 are completed at the same time, there are
- $128 + 64 - 32 = 160$ ways to do either task.
- In set theory, this corresponds to sets A_1 and A_2 that are **not** disjoint. Then we have:
$$|A_1 \cup A_2| = |A_1| + |A_2| - |A_1 \cap A_2|$$
- This is called the **principle of inclusion-exclusion**.

Tree Diagrams

- How many bit strings of length four do not have two consecutive 1s?



There are 8 strings.

The Pigeonhole Principle

- **The pigeonhole principle:** If $(k + 1)$ or more objects are placed into k boxes, then there is at least one box containing two or more of the objects.
- **Example 1:** If there are 11 players in a soccer team that wins 12-0, there must be at least one player in the team who scored at least twice.
- **Example 2:** If you have 6 classes from Monday to Friday, there must be at least one day on which you have at least two classes.

The Pigeonhole Principle

- **The generalized pigeonhole principle:** If N objects are placed into k boxes, then there is at least one box containing at least $\lceil N/k \rceil$ of the objects.
- **Example 1:** In a 60-student class, at least 12 students will get the same letter grade (A, B, C, D, or F).
- **Example 2:** In a 61-student class, at least 13 students will get the same letter grade.

The Pigeonhole Principle

- **Example 3:** Assume you have a drawer containing a random distribution of a dozen brown socks and a dozen black socks. It is dark, so how many socks do you have to pick to be sure that among them there is a matching pair?
- There are two types of socks, so if you pick at least 3 socks, there must be either at least two brown socks or at least two black socks.
- Generalized pigeonhole principle: $\lceil 3/2 \rceil = 2$.

Permutations and Combinations

- ❑ How many different sets of 3 people can we pick from a group of 6?
- ❑ There are 6 choices for the first person, 5 for the second one, and 4 for the third one, so there are $6 \cdot 5 \cdot 4 = 120$ ways to do this.
- ❑ **This is not the correct result!**
- ❑ For example, picking person C, then person A, and then person E leads to the **same group** as first picking E, then C, and then A.
- ❑ However, these cases are counted **separately** in the above equation.

Permutations and Combinations

- So how can we compute how many different subsets of people can be picked (that is, we want to disregard the order of picking) ?
- To find out about this, we need to look at **permutations and combinations**.

Permutations and Combinations

- A **permutation** of a set of distinct objects is an ordered arrangement of these objects.
- An ordered arrangement of r elements of a set is called an **r -permutation**.

Permutations and Combinations

- **Example:** Let $S = \{1, 2, 3\}$.
The arrangement 3, 1, 2 is a permutation of S .
The arrangement 3, 2 is a 2-permutation of S .
- The number of r -permutations of a set with n distinct elements is denoted by $P(n, r)$.
- We can calculate $P(n, r)$ with the product rule:
$$P(n, r) = n \cdot (n - 1) \cdot (n - 2) \cdot \dots \cdot (n - r + 1).$$

(n choices for the first element, $(n - 1)$ for the second one, $(n - 2)$ for the third one...)

Permutations and Combinations

□ Example:

$$\begin{aligned} P(8, 3) &= 8 \cdot 7 \cdot 6 = 336 \\ &= (8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1) / (5 \cdot 4 \cdot 3 \cdot 2 \cdot 1) \end{aligned}$$

□ General formula:

$$P(n, r) = n! / (n - r)!$$

Permutations and Combinations

- An **r-combination** of elements of a set is an **unordered selection** of r elements from the set.
- Thus, an r -combination is simply a subset of the set with r elements.
- **Example:** Let $S = \{1, 2, 3, 4\}$.
Then $\{1, 3, 4\}$ is a 3-combination from S .

The number of r -combinations of a set with n distinct elements is denoted by $C(n, r)$.
- **Example:** $C(4, 2) = 6$, since, for example, the 2-combinations of a set $\{1, 2, 3, 4\}$ are $\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}$.

Permutations and Combinations

- $C(n, r) = n! / (r!(n - r)!)$
- Now we can answer our initial question: How many ways are there to pick a set of 3 people from a group of 6 (disregarding the order of picking)?

$$C(6, 3) = 6! / (3! \cdot 3!) = 720 / (6 \cdot 6) = 720 / 36 = 20$$

There are 20 different ways, that is, 20 different groups to be picked.

Permutations and Combinations

Corollary:

□ Let n and r be nonnegative integers with $r \leq n$.

Then $C(n, r) = C(n, n - r)$.

□ Note that “picking a group of r people from a group of n people” **is the same as** “splitting a group of n people into a group of r people and another group of $(n - r)$ people”.

Permutations and Combinations

□ Example:

A soccer club has 8 female and 7 male members. For today's match, the coach wants to have 6 female and 5 male players on the grass. How many possible configurations are there?

$$\begin{aligned}C(8, 6) \cdot C(7, 5) &= 8!/(6! \cdot 2!) \cdot 7!/(5! \cdot 2!) \\&= 28 \cdot 21 \\&= 588\end{aligned}$$


Combinations

Pascal's Identity:

- Let n and k be positive integers with $n \geq k$.
Then $C(n + 1, k) = C(n, k - 1) + C(n, k)$.


Pascal's Triangle

- In Pascal's triangle, each number is the sum of the numbers to its upper left and upper right:



Pascal's Triangle

- Since we have $C(n + 1, k) = C(n, k - 1) + C(n, k)$ and $C(0, 0) = 1$, we can use Pascal's triangle to simplify the computation of $C(n, k)$:



Binomial Coefficients

- Expressions of the form $C(n, k)$ are also called **binomial coefficients**.
- A **binomial expression** is the sum of two terms, such as $(a + b)$.

Consider $(a + b)^2 = (a + b)(a + b)$.

When expanding such expressions, we have to form all possible products of a term in the first factor and a term in the second factor:

$$(a + b)^2 = a \cdot a + a \cdot b + b \cdot a + b \cdot b$$

Then we can sum identical terms:

$$(a + b)^2 = a^2 + 2ab + b^2$$

Binomial Coefficients

□ This leads us to the following formula:

$$(a + b)^n = \sum_{j=0}^n C(n, j) \cdot a^{n-j} b^j \quad (\text{Binomial Theorem})$$

With the help of Pascal's triangle, this formula can considerably simplify the process of expanding powers of binomial expressions.

For example, the fifth row of Pascal's triangle

(1 - 4 - 6 - 4 - 1) helps us to compute $(a + b)^4$:

$$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$$

Summary

- In this part, we discussed:
 - Basic Counting Principles
 - Inclusion-Exclusion Principle
 - The Pigeonhole Principle
 - Permutations and Combinations
 - Pascal's Triangle and Binomial Coefficients

References

- These slides are largely based on the following two sources:
 - Slides by Marc Pomplun, UMass Boston
 - Official McGraw Hill's Slides for our textbook