

Drug Facts (continued)

3

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stool • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present
- any new symptoms appear • ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Directions

• **do not take more than directed**

(see overdose warning)

• **adults and children 12 years and over:**

place 1 powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose or may stir powder into a glass of water or other liquid.

• **do not take more than 4 powders in 24 hours unless directed by a doctor.**

• **Children under 12 years of age:** ask a doctor.

Other Information

- each powder contains: **potassium 51 mg**
- store below 25°C (77°F)

Inactive ingredient

lactose monohydrate, potassium chloride

Questions?

1-833-663-2933 (applies to all places)

1

RETAIN LEAFLET UPON OPENING

AERO

EXTRA STRENGTH
HEADACHE POWDER

Drug Facts

Active Ingredients (in each pack).....Purposes

Acetaminophen 260mgPain reliver

Aspirin (NSAID*) 520mgPain reliver

Caffeine 32.5 mgPain reliver aid

*nonsteroidal anti-inflammatory drug

Drug Facts (continued)

2

Uses

- temporarily relieves minor aches and pains due to: • headache • minor arthritis pain • muscular aches • colds • temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause severe allergic reaction which may include: • hives • facial swelling • shock • asthma (wheezing).

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4 powders in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains a NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed.

Caffeine warning: This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use • if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if • you have liver disease stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis the blood thinning drug warfarin