

AX-17 Germany - Approved Product Information v4.2

Document ID: DE-AX17-LBL-042


Status: APPROVED

Effective: 15-Sep-2026

Jurisdiction: Germany

Supersedes: DE-AX17-LBL-038

Controlled fact: moderate renal impairment starting dose = 25 mg once daily.



Indications and Population

Indications and Population. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 10.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 13.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 10.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 16.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 8.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 18.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 12.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 14.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Indications and Population. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 6.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration

Dosage and Administration. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 3.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 15.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 13.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 16.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 3.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 17.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 5.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 13.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 14.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 4.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Dosage and Administration. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 17.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment

Renal Impairment. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 4.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 14.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 8.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 4.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 3.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 5.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 13.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 15.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 16.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 15.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 5.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Impairment. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 15.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Table 3. Renal dose modification

Renal category	eGFR	Starting dose	Maximum	Monitoring
Normal	>=90	50 mg OD	100 mg/day	Routine
Mild	60-89	50 mg OD	100 mg/day	Routine
Moderate	30-59	25 mg OD	50 mg/day	Recheck 2-4 weeks
Severe	15-29	Specialist review	25 mg/day	Enhanced

Hepatic Impairment

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 15.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 9.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 6.8% of the described subgroup. This value is

descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 8.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 16.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 17.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 3.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 17.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 13.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Hepatic Impairment. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 5.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 4.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 18.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 14.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 16.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 15.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 10.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 11.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 3.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 11.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 3.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Warnings and Precautions. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 8.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 8.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 12.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 11.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 6.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 6.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 12.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 17.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 11.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 5.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 7.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 4.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Clinical Pharmacology. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 7.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.




Figure caption: controlled evidence; interpret with the surrounding section.

Adverse Reactions

Adverse Reactions. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 12.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 7.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 6.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 10.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 9.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 3.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 6.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 12.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 15.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Adverse Reactions. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 10.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 8.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 10.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 7.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 16.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 16.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 7.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 6.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 8.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 18.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Evidence Interpretation. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 12.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling

Traceability and Handling. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 13.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 12.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 7.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 5.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 10.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 4.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 8.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 12.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 16.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 13.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Traceability and Handling. The controlled AX-17 evidence package for Germany included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 15.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.