

# HN-204 Clinical Study Report - AX-17

Study ID: HN-204

Status: FINAL

Randomized population: 240

Tables and figures are evidence-bearing.

## Study Synopsis

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 4.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 6.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 11.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 17.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 11.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 12.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 10.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 13.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Study Synopsis. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 12.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

## Objectives and Endpoints

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 4.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 5.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 9.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 7.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 18.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 14.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 12.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 12.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Objectives and Endpoints. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 3.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

## Population and Disposition

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 8.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 10.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 7.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 15.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 7.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 4.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 11.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 11.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 17.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Population and Disposition. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 13.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

## Baseline Characteristics

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 4.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 14.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 12.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 16.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 3.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 4.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 15.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 18.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 10.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Baseline Characteristics. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 7.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

## Efficacy Results

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 5.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 9.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 6.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 9.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 13.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 12.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 12.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 14.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 4.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 12.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Efficacy Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 7.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

## Safety Results

Safety Results. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 15.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 11.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 18.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 4.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 14.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 7.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 11.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 9.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 10.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 9.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 14.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 16.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Safety Results. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 6.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Table 14. Treatment-emergent adverse events

| Adverse event          | AX-17 % | Control % | Serious? |
|------------------------|---------|-----------|----------|
| Nausea                 | 12.7    | 6.0       | No       |
| Headache               | 14.0    | 6.6       | No       |
| Fatigue                | 15.3    | 7.2       | No       |
| Dizziness              | 16.6    | 7.8       | No       |
| Rash                   | 17.8    | 8.4       | No       |
| Transaminase elevation | 19.1    | 9.0       | Yes      |




Figure caption: controlled evidence; interpret with the surrounding section.

## Renal Subgroup Analysis

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 14.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 14.4% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 7.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 7.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 16.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 15.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 16.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 10.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 14.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 17.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 16.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Renal Subgroup Analysis. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 5.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Table 18. Renal subgroup summary

| Renal stratum | N  | Exposure ratio | Interpretation                   |
|---------------|----|----------------|----------------------------------|
| Normal        | 80 | 1.0            | Reference                        |
| Mild          | 60 | 1.12           | Small increase                   |
| Moderate      | 48 | 1.48           | Supports reduced-dose evaluation |
| Severe        | 20 | 1.91           | Limited sample                   |

Interpretation and Limitations

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 8.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Rash was reported in 14.2% of the described subgroup. This value is

descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 4.0% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 14.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 9.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 15.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 9.6% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 5.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 16.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Fatigue was reported in 5.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which

it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Interpretation and Limitations. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 18.1% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

## Appendices and Definitions

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 13.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 180 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 16.7% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 96 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 15.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 48 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Headache was reported in 8.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 12.3% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Transaminase elevation was reported in 5.9% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 72 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 13.5% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 120 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Dizziness was reported in 6.8% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.

Appendices and Definitions. The controlled AX-17 evidence package for GLOBAL included 240 evaluable participants in this analysis. Investigators recorded treatment-emergent findings, laboratory changes, dose interruptions, concomitant therapy, and clinically relevant renal and hepatic observations. Nausea was reported in 9.2% of the described subgroup. This value is descriptive and must be interpreted only within the population, study, jurisdiction, version, and effective-date context in which it appears. Tables, warnings, captions, footnotes, and parent headings may modify the meaning of this paragraph and therefore form part of the retrievable evidence. HelixNova medical review requires source identity, document status, page context, and authority to remain traceable through downstream retrieval and generation.