Fine-Grained Buy-Many Mechanisms are Not Much Better than Bundling

Sepehr Assadi¹ Vikram Kher² George Li³ Ariel Schvartzman⁴

¹Rutgers University

²University of Southern California

³University of Maryland, College Park

⁴Google Research





Table of Contents

Basic Auction Theory

2 Our Contribution: Buy-k Mechanisms

Conclusion

Vikram Kher (USC)

Table of Contents

Basic Auction Theory

2 Our Contribution: Buy-k Mechanisms

Conclusion

Single Item, Single Buyer Auctions

- Single item and single buyer.
- ullet Buyer samples their value for the item from a distribution \mathcal{D} .
- Buyer will purchase the item if they value it at least its posted price.

Single Item, Single Buyer Auctions

- Single item and single buyer.
- ullet Buyer samples their value for the item from a distribution \mathcal{D} .
- Buyer will purchase the item if they value it at least its posted price.

Question

How should the seller price the item to maximize expected revenue?

Example of Single Item Auction

Historically, people are willing to pay either \$2, \$4, or \$6 for a cup of coffee.



Example of Single Item Auction

Historically, people are willing to pay either \$2, \$4, or \$6 for a cup of coffee.

	pr(sell)	Rev
P(coffee) = 2	1	2
P(coffee) = 4	2/3	8/3
P(coffee) = 6	1/3	2



Example of Single Item Auction

Historically, people are willing to pay either \$2, \$4, or \$6 for a cup of coffee.

pay	Juuru	
	<u>₩</u>	
′		

Coffee Shop

	pr(sell)	Rev
P(coffee) = 2	1	2
P(coffee) = 4	2/3	8/3
P(coffee) = 6	1/3	2

Remark

For this example, the optimal price is \$4. Myerson [1981] proved this general "algorithm" is optimal. Namely, the optimal selling price is $p^* = argmax_p(p \cdot (1 - F(p)))$, where F is the CDF of \mathcal{D} .

Two natural generalizations of Myerson's idea to a multi-parameter space. Note, we allow \mathcal{D} to be arbitrarily correlated. We additionally assume that buyers can only purchase at most one menu option (buy-one model).

Two natural generalizations of Myerson's idea to a multi-parameter space. Note, we allow \mathcal{D} to be arbitrarily correlated. We additionally assume that buyers can only purchase at most one menu option (buy-one model).

 Item Pricing: Compute "Myerson price" for each item individually and sell each item at this respective price. Every subset of items is also listed at the sum of its component prices.

Two natural generalizations of Myerson's idea to a multi-parameter space. Note, we allow $\mathcal D$ to be arbitrarily correlated. We additionally assume that buyers can only purchase at most one menu option (buy-one model).

- Item Pricing: Compute "Myerson price" for each item individually and sell each item at this respective price. Every subset of items is also listed at the sum of its component prices.
- Bundling: Group all items together and calculate the collective "Myerson price". Sell this grand bundle at the calculated price.

Two natural generalizations of Myerson's idea to a multi-parameter space. Note, we allow $\mathcal D$ to be arbitrarily correlated. We additionally assume that buyers can only purchase at most one menu option (buy-one model).

- Item Pricing: Compute "Myerson price" for each item individually and sell each item at this respective price. Every subset of items is also listed at the sum of its component prices.
- Bundling: Group all items together and calculate the collective "Myerson price". Sell this grand bundle at the calculated price.
- These are what are known as simple mechanisms (roughly, polynomial number of menu items).

Two natural generalizations of Myerson's idea to a multi-parameter space. Note, we allow $\mathcal D$ to be arbitrarily correlated. We additionally assume that buyers can only purchase at most one menu option (buy-one model).

- Item Pricing: Compute "Myerson price" for each item individually and sell each item at this respective price. Every subset of items is also listed at the sum of its component prices.
- Bundling: Group all items together and calculate the collective "Myerson price". Sell this grand bundle at the calculated price.
- These are what are known as simple mechanisms (roughly, polynomial number of menu items).

Theorem [BCKW 2015, HN 2017]

There exist distributions over two items such that the optimal revenue is unbounded, but the revenue from item pricing/bundling is finite.

Multi-item auctions are hard!

- Computational Barriers
 - Hard to compute optimal mechanisms in some simple settings [CDOPSY 2015, DDT 2014].

Multi-item auctions are hard!

- Computational Barriers
 - Hard to compute optimal mechanisms in some simple settings [CDOPSY 2015, DDT 2014].
- Structural Barriers
 - Optimal mechanisms may require infinitely many menu options [DDT 2014].

Multi-item auctions are hard!

- Computational Barriers
 - Hard to compute optimal mechanisms in some simple settings [CDOPSY 2015, DDT 2014].
- Structural Barriers
 - Optimal mechanisms may require infinitely many menu options [DDT 2014].
- Mathematical Barriers
 - Simple mechanisms can't approximate optimal mechanisms ([BCKW 2015, HN 2017].

Example of Unintuitive Optimal mechanism



	Coffee	Scone
Bob	2	0
Sally	0	4
Alice	4	6

Optimal Mechanism:

Price of Coffee: \$2

• Price of Scone: \$4

Price of Coffee and Scone: \$7.99

Example of Unintuitive Optimal mechanism



	Coffee	Scone
Bob	2	0
Sally	0	4
Alice	4	6

Optimal Mechanism:

• Price of Coffee: \$2

• Price of Scone: \$4

Price of Coffee and Scone: \$7.99

Remark

Optimal Mechanism achieves expected revenue (2+4+7.99)/3=\$4.66. But, crucially, this mechanism relies on a super-additive pricing scheme. The seller is taking advantage of the fact that the buyer can only interact with the mechanism once! If the buyer could purchase multiple menu items then the expected revenue drops to \$4!

8/31

• A mechanism is buy-many incentive-compatible (IC) if the buyer prefers to purchase one menu option from the mechanism as opposed to any multi-set of allocations [BCKW 2015].

- A mechanism is buy-many incentive-compatible (IC) if the buyer prefers to purchase one menu option from the mechanism as opposed to any multi-set of allocations [BCKW 2015].
- Any deterministic sub-additive mechanism is buy-many incentive compatible

- A mechanism is buy-many incentive-compatible (IC) if the buyer prefers to purchase one menu option from the mechanism as opposed to any multi-set of allocations [BCKW 2015].
- Any deterministic sub-additive mechanism is buy-many incentive compatible
- Does this restricted class of mechanisms allow for optimal mechanisms to be approximated by simple ones?

- A mechanism is buy-many incentive-compatible (IC) if the buyer prefers to purchase one menu option from the mechanism as opposed to any multi-set of allocations [BCKW 2015].
- Any deterministic sub-additive mechanism is buy-many incentive compatible
- Does this restricted class of mechanisms allow for optimal mechanisms to be approximated by simple ones?

Theorem [BCKW 2015, CTT 2019]

For arbitrary distributions with arbitrary buyer valuations, item pricing gets within $O(\log n)$ of the revenue of the optimal buy-many IC mechanism.

Vikram Kher (USC) Buy-k Mechanisms October 2022 9 / 31

The Previous State of Affairs

Buy-one mechanisms: Sellers have too much power to extract revenue! Buy-many mechanisms: Buyers have too much power!

10/31

The Previous State of Affairs

Buy-one mechanisms: Sellers have too much power to extract revenue! Buy-many mechanisms: Buyers have too much power!

Question

Is there a good middle ground between the classical buy-one model and newer buy-many model? More specifically, to what extent does the number of menu option deviations that the seller must guard against dictate whether simple mechanisms can approximate optimal ones.

Table of Contents

Basic Auction Theory

2 Our Contribution: Buy-k Mechanisms

Conclusion

Introducing Buy-k IC Mechanisms

A mechanism is buy-k incentive-compatible (IC) if the buyer prefers
to purchase a single menu option from mechanism compared to any
other multi-set of menu options of size at most k.

Introducing Buy-k IC Mechanisms

- A mechanism is buy-k incentive-compatible (IC) if the buyer prefers to purchase a single menu option from mechanism compared to any other multi-set of menu options of size at most k.
- Setting k = 1 returns notion of buy-one IC.

12 / 31

Vikram Kher (USC) Buy-k Mechanisms October 2022

Introducing Buy-k IC Mechanisms

- A mechanism is buy-k incentive-compatible (IC) if the buyer prefers to purchase a single menu option from mechanism compared to any other multi-set of menu options of size at most k.
- Setting k = 1 returns notion of buy-one IC.
- Setting $k = \infty$ returns the notion of buy-many IC.

Vikram Kher (USC)

Benchmarks of Interests

For a given distribution \mathcal{D} , let

- $\mathrm{BRev}(\mathcal{D})$ be the revenue of the mechanism which sells the grand bundle for its optimal price.
- Buy^kRev(\mathcal{D}) be the revenue of the optimal buy-k incentive-compatible mechanism.
- $\operatorname{Buy}^k\operatorname{Rev}(\mathcal{D},\mathcal{M})$ be the revenue of (not necessarily buy-k incentive-compatible) mechanism \mathcal{M} when the buyer is allowed to buy up to k menu entries from \mathcal{M} .

Theorem 1 [AKLS 2022]

For buyers with additive valuations and allowing arbitrary n-dimensional distributions, $BRev(D)O(n^2) \ge Buy^n Rev(D)$

14 / 31

Vikram Kher (USC) Buy-k Mechanisms

Theorem 2 [AKLS 2022]

For buyers with general monotone valuations and allowing arbitrary n-dimensional distributions, $BRev(\mathcal{D})O(2^n n^2) \geq Buy^n Rev(D)$

14 / 31

Vikram Kher (USC) Buy-k Mechanisms Octo

Theorem 3 [AKLS 2022]

For buyers with additive valuations, if $k \leq n^{1/2-\epsilon}$ for some $\epsilon > 0$, there exists a n-dimensional distribution \mathcal{D} such that $Buy^k Rev(\mathcal{D}) \geq BRev(\mathcal{D}) exp(\Omega(n^{\epsilon}))$

14 / 31

Vikram Kher (USC) Buy-k Mechanisms October 2022

Theorem 1 [AKLS 2022]

For buyers with additive valuations and allowing arbitrary n-dimensional distributions, $BRev(D)O(n^2) \ge Buy^n Rev(D)$

Theorem 2 [AKLS 2022]

For buyers with general monotone valuations and allowing arbitrary n-dimensional distributions, $BRev(\mathcal{D})O(2^n n^2) \geq Buy^n Rev(D)$

Theorem 3 [AKLS 2022]

For buyers with additive valuations, if $k \leq n^{1/2-\epsilon}$ for some $\epsilon > 0$, there exists a *n*-dimensional distribution \mathcal{D} such that $Buy^k Rev(\mathcal{D}) \geq BRev(\mathcal{D}) exp(\Omega(n^{\epsilon}))$

15 / 31

Vikram Kher (USC) Buy-k Mechanisms October 2022

Preliminaries

• Buyer is utility-maximizing and draws their valuation x from a known, possibly correlated n-dimensional distribution \mathcal{D} , with support set $\mathcal{X} = \text{SUPP}(\mathcal{D})$

Preliminaries

- Buyer is utility-maximizing and draws their valuation x from a known, possibly correlated n-dimensional distribution \mathcal{D} , with support set $\mathcal{X} = \text{SUPP}(\mathcal{D})$
- Valuations are additive (i.e., for any subset of items $S \subseteq [n]$, $x(S) = \sum_{i \in S} x_i$)

Vikram Kher (USC)

Preliminaries

- Buyer is utility-maximizing and draws their valuation x from a known, possibly correlated n-dimensional distribution \mathcal{D} , with support set $\mathcal{X} = \text{SUPP}(\mathcal{D})$
- Valuations are additive (i.e., for any subset of items $S \subseteq [n]$, $x(S) = \sum_{i \in S} x_i$)
- A mechanism $\mathcal{M} = (p, q)$ is then defined by a pair of functions:
 - Payment Function: $p: \mathcal{X} \to \mathcal{R}_{\geq 0}$
 - Allocation Function: $q:\mathcal{X} \to [0,1]^n$

Preliminaries

- Buyer is utility-maximizing and draws their valuation x from a known, possibly correlated n-dimensional distribution \mathcal{D} , with support set $\mathcal{X} = \text{SUPP}(\mathcal{D})$
- Valuations are additive (i.e., for any subset of items $S \subseteq [n]$, $x(S) = \sum_{i \in S} x_i$)
- A mechanism $\mathcal{M} = (p, q)$ is then defined by a pair of functions:
 - Payment Function: $p: \mathcal{X} \to \mathcal{R}_{\geq 0}$
 - Allocation Function: $q: \mathcal{X} \to [\overline{0}, 1]^n$
- Given a possibly randomized allocation of items $\vec{q} \in [0, 1]^n$, we use $x(\vec{q}) p(x)$ to denote the buyer's expected utility.

Purchasing Multiple Allocations

• Let $\Lambda = \{\vec{q}_1, \vec{q}_2, \dots, \vec{q}_k, \dots\}$ be a multi-set of allocations (of possibly unbounded size).

Vikram Kher (USC) Buy-k Mechanisms October 2022 17 / 31

Purchasing Multiple Allocations

- Let $\Lambda = \{\vec{q}_1, \vec{q}_2, \dots, \vec{q}_k, \dots\}$ be a multi-set of allocations (of possibly unbounded size).
- We let $Lot(\Lambda) \in [0,1]^n$ denote the expected allocation that results from being allocated every $\vec{q_i} \in \Lambda$ independently and at once.
 - In particular, for all $j \in [n]$, $\vec{\text{Lot}}(\Lambda)_j = 1 \prod_{i=1} (1 q_{ij})$.

Purchasing Multiple Allocations

- Let $\Lambda = \{\vec{q}_1, \vec{q}_2, \dots, \vec{q}_k, \dots\}$ be a multi-set of allocations (of possibly unbounded size).
- We let $Lot(\Lambda) \in [0,1]^n$ denote the expected allocation that results from being allocated every $\vec{q_i} \in \Lambda$ independently and at once.
 - In particular, for all $j \in [n]$, $\vec{\mathrm{Lot}}(\Lambda)_j = 1 \prod_{i=1} (1-q_{ij})$.

Buy-k IC Definition

 $\mathcal M$ is buy-k incentive compatible if for every valuation $x \in \mathcal X$ there exists some $(\vec q,p) \in \mathcal M$ such that $x(\vec q)-p(x) \geq x(\mathop{\rm Lot}(\Lambda))-\sum_{i=1}p(\vec q_i)$ for any possible multi-set of menu options Λ of size at most k.

4□ > 4□ > 4 = > 4 = > = 90

Menu Gap: An Intermediary Measure

The definitions below are generalizations of ones proposed in [HN 2017].

Gap Definition

Let $X=\{\vec{x_i}\}_{i=1}^N\in\mathbb{R}^n_{\geq 0},$ $Q=\{\vec{q_i}\}_{i=0}^N\in[0,1]^n$ be sequences of vectors with $\vec{q_0}=\vec{0}^n$. Then,

$$\operatorname{gap}_{i}^{k}(X,Q) = \min_{j_{1},j_{2},\ldots,j_{k} < i} \vec{x}_{i} \cdot (\vec{q}_{i} - \operatorname{Lot}(\vec{q}_{j_{1}}, \vec{q}_{j_{2}}, \ldots, \vec{q}_{j_{k}}))$$

4□ > 4ⓓ > 4≧ > 4≧ > ½

18 / 31

Menu Gap: An Intermediary Measure

The definitions below are generalizations of ones proposed in [HN 2017].

Gap Definition

Let $X = {\vec{x_i}}_{i=1}^N \in \mathbb{R}_{>0}^n, Q = {\vec{q_i}}_{i=0}^N \in [0,1]^n$ be sequences of vectors with $\vec{q}_0 = \vec{0}^n$. Then,

$$\operatorname{gap}_{i}^{k}(X,Q) = \min_{j_{1},j_{2},\ldots,j_{k} < i} \vec{x}_{i} \cdot (\vec{q}_{i} - \operatorname{Lot}(\vec{q}_{j_{1}}, \vec{q}_{j_{2}}, \ldots, \vec{q}_{j_{k}}))$$

- Think of X as holding a sequence of possible valuations vectors and Q holding their corresponding allocations
- $\operatorname{gap}_{i}^{k}(X,Q)$ is the largest price a seller can post on menu entry (\vec{q}_i, p_i) such that x_i will prefer it any subset of the previous allocations for free.

Menu Gap: An Intermediary Measure (cont.)

Gap Definition

Let $X=\{\vec{x_i}\}_{i=1}^N\in\mathbb{R}_{\geq 0}^n,$ $Q=\{\vec{q_i}\}_{i=0}^N\in[0,1]^n$ be sequences of vectors with $\vec{q_0}=\vec{0}^n$. Then,

$$\operatorname{gap}_i^k(X,Q) = \min_{j_1,j_2,\ldots,j_k < i} \vec{x}_i \cdot (\vec{q}_i - \operatorname{Lot}(\vec{q}_{j_1},\vec{q}_{j_2},\ldots,\vec{q}_{j_k}))$$

MenuGap Definition

$$\operatorname{MenuGap}^k(X,Q) = \sum_{i=1}^N \operatorname{gap}_i^k(X,Q) / ||\vec{x_i}||_1$$

Proof Strategy for Theorem 1

Theorem 1 [AKLS 2022]

For buyers with additive valuations and allowing arbitrary n-dimensional distributions, $BRev(D)O(n^2) \ge Buy^nRev(D)$

Proof Strategy

- First, show that there exists an X, Q such that $\operatorname{MenuGap}^k(X, Q)$ upperbounds the revenue gap between $\operatorname{Buy}^k \operatorname{Rev}(\mathcal{D}, \mathcal{M})$ and $\operatorname{BRev}(\mathcal{D})$, up to a O(k) factor.
- ② Then, show that for k = n, $n \ge \text{MenuGap}^n(X, Q)$ for all X, Q.

Vikram Kher (USC) Buy-k Mechanisms October 2022 20 / 31

Lemma 1

Lemma 1


For any distribution \mathcal{D} , additive valuation over n items and any buy-k incentive compatible mechanism \mathcal{M} for \mathcal{D} , there exists a sequence of valuations $X = \{x_i\}_{i=1} \subseteq \mathcal{X}$, and a sequence of allocations $Q = \{\vec{q}_i\}_{i=0} \subseteq \mathcal{M}$ (starting with $\vec{q}_0 = (0, \ldots, 0)$) such that

$$\operatorname{MenuGap}^k(X,Q) \geq \frac{\operatorname{Buy}^k \operatorname{Rev}(\mathcal{D},\mathcal{M})}{8k \cdot \operatorname{BRev}(\mathcal{D})}.$$

Proof Outline for Lemma 1

Step One

- **1** Take \mathcal{M} and create a sub-menu \mathcal{M}' with the following properties:
 - All prices are at least c.
 - All prices belong to the set of intervals $\bigcup_{i=0}^{\infty} [c \cdot (2k)^{2i+a}, c \cdot (2k)^{2i+a+1})$ for an $a \in \{0,1\}$.
 - Buy^kRev $(\mathcal{D}, \mathcal{M}') \ge \frac{\text{Buy}^k \text{Rev}(\mathcal{D}, \mathcal{M}) c}{2}$.



Proof Outline for Lemma 1 (cont.)

Step Two

- Let \mathcal{M}' be the menu which satisfies step one. We will use this menu to find our X, Q:
 - Let $\mathcal{B}_j \subseteq \mathcal{M}$ be the sub-menu that has all menu entries priced in $[c \cdot (2k)^{2j+a}, c \cdot (2k)^{2j+a+1})$ for the same $a \in \{0, 1\}$.
 - Let $\vec{x_j}$ be the valuation on \mathcal{B}_j such that $||\vec{x_j}||_1 \leq (1+\delta)||\vec{x}||_1 \quad \forall \vec{x} \in \mathcal{B}_j$.
 - Then,

$$\Pr(x \in \mathcal{B}_j) \le \frac{\mathrm{BRev}(\mathcal{D})(1+\delta)}{||\vec{x}_j||_1}$$

Selecting our choice of X, Q

Let (X, Q) be defined by these \vec{x}_j and their corresponding \vec{q}_j .

◆ロト ◆御 ト ◆恵 ト ◆恵 ト ・恵 ・ 夕久で

Proof Outline for Lemma 1 (cont.)

Step Three

We can now lowerbound $gap_i^k(X,Q)/||\vec{x_i}||_1$. Recall,

$$\operatorname{gap}_i^k(X,Q) = \min_{j_1,j_2,\dots,j_k < i} \vec{x}_i \cdot (\vec{q}_i - \operatorname{Lot}(\vec{q}_{j_1},\vec{q}_{j_2},\dots,\vec{q}_{j_k}))$$

• Since \mathcal{M} was buy-k incentive compatible, we have that for each x_j and any previous set of k options $\vec{q}_{j_1}, \ldots, \vec{q}_{j_k}$,

$$x_j(\vec{q}_j)-p_j\geq x_j(\operatorname{Lot}(\vec{q}_{j_1},\ldots,\vec{q}_{j_k}))-\sum_{i=1}^k p_{j_i}.$$

By rearrangment, we have that

$$\frac{{\rm gap}_j^k(X,Q)}{||\vec{x_j}||_1} \ge \frac{p_j - \sum_{i=1}^k p_{j_i}}{||\vec{x_j}||_1}.$$

Proof for Lemma 1 (cont.)

Step Three (cont.)

Continuing from before

① Since for $j_i < j$, $p_j \ge 2k \cdot p_{j_i}$. Additionally, recall that

$$\Pr(x \in \mathcal{B}_j) \leq \frac{\operatorname{BRev}(\mathcal{D})(1+\delta)}{||\vec{x_j}||_1}.$$

It follows that

$$\frac{\operatorname{gap}_j^k(X,Q)}{||\vec{x}_j||_1} \geq \frac{p_j}{2 \cdot ||\vec{x}_j||_1} \geq \frac{\Pr(x \in \mathcal{B}_j)p_j}{2 \cdot \operatorname{BRev}(\mathcal{D})(1+\delta)}.$$

Proof for Lemma 1

Lemma 1

For any distribution \mathcal{D} , additive valuation over n items and any buy-kincentive compatible mechanism \mathcal{M} for \mathcal{D} , there exists a sequence of valuations $X = \{x_i\}_{i=1} \subseteq \mathcal{X}$, and a sequence of allocations $Q = \{\vec{q}_i\}_{i=0} \subseteq \mathcal{M}$ (starting with $\vec{q}_0 = (0, ..., 0)$) such that

$$\operatorname{MenuGap}^k(X,Q) \geq \frac{\operatorname{Buy}^k\operatorname{Rev}(\mathcal{D},\mathcal{M})}{8k\cdot\operatorname{BRev}(\mathcal{D})}.$$

Proof.

- **1** Take as a fact that $\sum_{j} \Pr(x \in \mathcal{B}_{j}) p_{j} \geq \frac{(\operatorname{Buy}^{k}\operatorname{Rev}(\mathcal{D},\mathcal{M}) c)}{4k}$.
- 2 Then, MenuGap^k $(X,Q) = \sum_{i} \frac{\operatorname{gap}_{i}^{k}(X,Q)}{||\vec{\mathbf{x}}||_{1}} \ge \sum_{i} \frac{\operatorname{Pr}(\mathbf{x} \in \mathcal{B}_{i})p_{i}}{2 \cdot \operatorname{BRey}(\mathcal{D})(1+\delta)} \ge$ $\frac{(\operatorname{Buy}^k\operatorname{Rev}(\mathcal{D},\mathcal{M})-c)}{8k\cdot\operatorname{BRev}(\mathcal{D})(1+\delta)}.$

Lemma 2

Lemma 2

For all sequences X, Q, where $X = \{\vec{x_i}\}_{i=1}^N \in \mathbb{R}_{>0}^n, Q = \{\vec{q_i}\}_{i=0}^N \in [0,1]^n$ with $\vec{q}_0 = \vec{0}^n$ and when buyer's have additive valuations, it holds that $\operatorname{MenuGap}^n(X,Q) \leq n.$

Proof.

Proof omitted. The (vague) intuition:

- Break down the sum of $gap_i^k(X,Q)$ into summing across ncoordinates of a vector.
- Show that the sum across each individual coordinate telescopes to 1.
- Since there are *n* coordinates, $\operatorname{MenuGap}^n(X, Q) \leq n$.
- Note, setting both sequences (X, Q) equal to the standard basis of \mathbb{R}^n makes this inequality tight.

Vikram Kher (USC) Buy-k Mechanisms October 2022 27 / 31

Proof for Theorem 1

Theorem 1 [AKLS 2022]

For buyers with additive valuations and allowing arbitrary distributions, $BRev(D)O(n^2) \ge Buy^n Rev(D)$

Proof.

Follows directly from applying Lemma 1 and Lemma 2.



Table of Contents

Basic Auction Theory

2 Our Contribution: Buy-k Mechanisms

Conclusion

29/31

Vikram Kher (USC) Buy-k Mechanisms October 2022

Big Take Aways

 Multi-item auctions are exhibit suprisingly unintuitive features. In particular, simple mechanisms cannot approximate optimal ones.

Big Take Aways

- Multi-item auctions are exhibit suprisingly unintuitive features. In particular, simple mechanisms cannot approximate optimal ones.
- In the buy-many world, many of these bad properties (like revenue inapproximability) disappear.

Big Take Aways

- Multi-item auctions are exhibit suprisingly unintuitive features. In particular, simple mechanisms cannot approximate optimal ones.
- In the buy-many world, many of these bad properties (like revenue inapproximability) disappear.
- Our work shows that when k = n, the optimal buy-k IC mechanism can be well-approximated by bundling.

Future Directions

• What happens to the revenue gap between optimal and simple mechanisms when $\sqrt{n} < k < n$?

◆ロト ◆問 ト ◆ 恵 ト ◆ 恵 ・ 釣 へ ご

Future Directions

- What happens to the revenue gap between optimal and simple mechanisms when $\sqrt{n} < k < n$?
- ullet Do there exist distributions ${\mathcal D}$ such that $\operatorname{Buy}^k \operatorname{Rev}(\mathcal{D}) > \operatorname{Buy}^{k+1} \operatorname{Rev}(\mathcal{D})$?

Future Directions

- What happens to the revenue gap between optimal and simple mechanisms when $\sqrt{n} < k < n$?
- Do there exist distributions \mathcal{D} such that $\operatorname{Buy}^k \operatorname{Rev}(\mathcal{D}) > \operatorname{Buy}^{k+1} \operatorname{Rev}(\mathcal{D})$?
- Not obvious how to efficiently test whether a mechanism is buy-many or buy-k IC for some k.

Thank you! Questions?