

**ABOUT THE TEST** FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

## PATIENT

**DISEASE** Lung adenocarcinoma

**DATE OF BIRTH** 20 November 1953

**SEX** Male

**MEDICAL RECORD #** Not given

## PHYSICIAN

**MEDICAL FACILITY** Arias Stella

**ADDITIONAL RECIPIENT** None

**MEDICAL FACILITY ID** 317319

**PATHOLOGIST** Not Provided

## SPECIMEN

**SPECIMEN SITE** Lung

**SPECIMEN ID** BP20-00860-1-A

**SPECIMEN TYPE** Block

**DATE OF COLLECTION** 21 December 2020

**SPECIMEN RECEIVED** 23 January 2021

## Biomarker Findings

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 3 Muts/Mb

## Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

**EGFR** amplification

**MET** amplification - equivocal<sup>†</sup>

**HGF** amplification - equivocal<sup>†</sup>

**RICTOR** amplification

**FGF10** amplification - equivocal<sup>†</sup>

**PARP1** amplification

**TP53** splice site 376-1G>A

<sup>†</sup> See About the Test in appendix for details.

9 Therapies with Clinical Benefit

28 Clinical Trials

0 Therapies with Lack of Response

## BIOMARKER FINDINGS

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 3 Muts/Mb

## GENOMIC FINDINGS

**MET** - amplification - equivocal

10 Trials see p. 18

**EGFR** - amplification

10 Trials see p. 14

**HGF** - amplification - equivocal

10 Trials see p. 16

**RICTOR** - amplification

9 Trials see p. 20

## ACTIONABILITY

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

### THERAPIES WITH CLINICAL BENEFIT (IN PATIENT'S TUMOR TYPE)

Capmatinib 2A

Crizotinib 2A

Afatinib

Dacomitinib

Erlotinib

Gefitinib

none

none

### THERAPIES WITH CLINICAL BENEFIT (IN OTHER TUMOR TYPE)

Cabozantinib

Cetuximab

Panitumumab

none

none

NCCN category

**GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIALS OPTIONS**

*For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.*

**FGF10 - amplification - equivocal** ..... p. 7    **TP53 - splice site 376-1G>A** ..... p. 8  
**PARP1 - amplification** ..... p. 7

**NOTE** Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

ORDERED TEST # ORD-1002521-02

## BIOMARKER FINDINGS

## BIOMARKER

## Microsatellite status

## RESULT

MS-Stable

### POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors<sup>1-3</sup>, including approved therapies nivolumab and pembrolizumab<sup>4</sup>. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared

with non-MSI-H cases (70% vs. 12%,  $p=0.001$ )<sup>5</sup>.

### FREQUENCY & PROGNOSIS

MSI-H is generally infrequent in NSCLC, reported in fewer than 1% of samples across several large studies<sup>6-11</sup>, whereas data on the reported incidence of MSI-H in SCLC has been limited and conflicting<sup>12-15</sup>. One study reported MSI-H in lung adenocarcinoma patients with smoking history, and 3 of 4 MSI-H patients examined also had metachronous carcinomas in other organs, although this has not been investigated in large scale studies<sup>6</sup>. The prognostic implications of MSI in NSCLC have not been extensively studied (PubMed, Oct 2020).

### FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor<sup>16</sup>. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2<sup>16-18</sup>. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers<sup>19-21</sup>. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins<sup>16,18,20-21</sup>.

## BIOMARKER

## Tumor Mutational Burden

## RESULT

3 Muts/Mb

### POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1<sup>22-24</sup>, anti-PD-1 therapies<sup>22-25</sup>, and combination nivolumab and ipilimumab<sup>26-30</sup>. Multiple clinical trials of PD-1- or PD-L1-targeting immune checkpoint inhibitors or combination of PD-1 and CTLA-4 inhibitors in NSCLC have reported that patients with tumors harboring TMB  $\geq 10$  Muts/Mb derive greater clinical benefit from these therapies than those with TMB  $< 10$  Muts/Mb (based on this assay or others); similarly, higher efficacy of anti-PD-1 or anti-PD-L1 immunotherapy for treatment of patients with NSCLC, compared with the use of chemotherapy, has been observed more significantly in cases of TMB  $\geq 10$  Muts/Mb (based on this assay or others)<sup>22-23,26-28,31-38</sup>. Improved OS of patients with NSCLC treated with pembrolizumab plus

chemotherapy relative to chemotherapy only<sup>39</sup>, or those treated with nivolumab plus ipilimumab also relative to chemotherapy<sup>40</sup>, has been observed across all TMB levels.

### FREQUENCY & PROGNOSIS

A large-scale genomic analysis found that unspecified lung non-small cell lung carcinoma (NSCLC), lung adenocarcinoma, and lung squamous cell carcinoma (SCC) samples harbored median TMBs between 6.3 and 9 Muts/Mb, and 12% to 17% of cases had an elevated TMB of greater than 20 Muts/Mb<sup>41</sup>. Lower TMB is observed more commonly in NSCLCs harboring known driver mutations (EGFR, ALK, ROS1, or MET) with the exception of BRAF or KRAS mutations, which are commonly observed in elevated TMB cases<sup>42</sup>. Although some studies have reported a lack of association between smoking and mutational burden in NSCLC<sup>43-44</sup>, several other large studies did find a strong association with increased TMB<sup>45-48</sup>. TMB  $> 10$  muts/Mb was found to be more frequent in NSCLC metastases compared with primary tumors for both adenocarcinoma (38% vs. 25%) and SCC (41% vs. 35%) subtypes<sup>49</sup>. A large study of Chinese patients with lung adenocarcinoma reported a shorter median OS for tumors with a higher number of mutations in a limited gene set compared with a lower mutation number (48.4 vs. 61.0 months)<sup>43</sup>.

Another study of patients with NSCLC correlated elevated TMB with poorer prognosis and significantly associated lower TMB in combination with PD-L1 negative status with longer median survival in patients with lung adenocarcinoma<sup>50</sup>. However, no significant prognostic association of TMB and/or PD-L1 status with survival has been reported in patients with lung SCC<sup>50-51</sup>.

### FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma<sup>52-53</sup> and cigarette smoke in lung cancer<sup>31,54</sup>, treatment with temozolomide-based chemotherapy in glioma<sup>55-56</sup>, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes<sup>57-61</sup>, and microsatellite instability (MSI)<sup>57,60-61</sup>. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents<sup>22-23,26-28,31-38,62</sup>.

ORDERED TEST # ORD-1002521-02

## GENOMIC FINDINGS

## GENE

**EGFR**

## ALTERATION

amplification

**POTENTIAL TREATMENT STRATEGIES**

EGFR amplification or expression may be associated with benefit from anti-EGFR antibodies, such as cetuximab<sup>63-66</sup>, panitumumab<sup>64</sup>, or necitumumab<sup>67</sup>, or EGFR TKIs that target wild-type EGFR<sup>68-72</sup>. Necitumumab is an anti-EGFR antibody that is approved to treat metastatic squamous NSCLC in combination with gemcitabine and cisplatin<sup>73-74</sup> that has also shown benefit in patients with CRC and melanoma<sup>75-76</sup>. Irreversible EGFR inhibitors, as well as HSP90 inhibitors, may be appropriate for patients with de novo or acquired resistance to EGFR-targeted therapy<sup>77-80</sup>. Preclinical studies have reported that EGFR-mutant cells<sup>77-79</sup>, including cells with exon 20 insertions<sup>81</sup>, are sensitive to HSP90 inhibitors. Consistent with preclinical data demonstrating that the EGFR inhibitor AZD3759 is capable of penetrating the blood-brain barrier and reducing the volume of brain and leptomeningeal metastases, preliminary results from a Phase 1 trial evaluating single-agent AZD3759 reported a

reduction in the volume of brain metastases in 40.0% (8/20) of patients with previously treated NSCLC harboring either EGFR L858R or EGFR exon 19 deletion, including 3 confirmed PRs and 3 unconfirmed PRs<sup>82-83</sup>. In a Phase 1/2 trial for advanced NSCLC, the brain-penetrant third-generation EGFR TKI lazertinib enabled ORRs of 54.3% (69/127) for all evaluable patients and 44.4% (8/18, intracranial) for patients with brain metastases<sup>84</sup>. The reovirus Reolysin targets cells with activated RAS signaling<sup>85-87</sup> and is in clinical trials in patients with some tumor types. Reolysin has demonstrated mixed clinical efficacy, with the highest rate of response reported for patients with head and neck cancer<sup>88-96</sup>. The role of EGFR or KRAS mutations as biomarkers for response to Reolysin in NSCLC is unclear<sup>97</sup>. Clinical and preclinical studies of lung cancer have shown that MET amplification is a common mechanism of resistance to EGFR inhibitors in first-line and later treatment settings<sup>98-104</sup>. Multiple studies have demonstrated that patients with a concurrent EGFR mutation and MET amplification, as recurrently observed at progression in patients with EGFR-mutated NSCLC on EGFR TKI, have benefited from a combination of MET- and EGFR-targeted therapies<sup>98,103-106</sup>.

**FREQUENCY & PROGNOSIS**

Amplification of EGFR has been variously

reported in 4-42% of non-small cell lung carcinoma (NSCLC) samples<sup>107-111</sup>. EGFR protein expression/overexpression has been reported in up to 70% of NSCLC cases<sup>109-114</sup>. In addition, expression of EGFR protein has been shown to be higher in lung squamous cell carcinoma samples as compared to lung adenocarcinoma<sup>115-116</sup>. In lung adenocarcinoma, EGFR gene amplification was a predictor of poor disease-free survival in all patients and of poor overall survival in patients with EGFR mutations<sup>117-118</sup>. Nuclear expression of EGFR in NSCLC has been reported to associate with higher disease stage, shorter progression-free survival, and shorter overall survival<sup>119</sup>. However, EGFR mutations have been reported to predict improved survival in patients with resected Stage 1-3 lung adenocarcinoma<sup>120</sup> or resected Stage 1 NSCLC<sup>121</sup>.

**FINDING SUMMARY**

EGFR encodes the epidermal growth factor receptor, which belongs to a class of proteins called receptor tyrosine kinases. In response to signals from the environment, EGFR passes biochemical messages to the cell that stimulate it to grow and divide<sup>122</sup>. Amplification of EGFR has been associated with increased expression of EGFR mRNA and protein in several cancer types<sup>110,123-124</sup>.

ORDERED TEST # ORD-1002521-02

## GENOMIC FINDINGS

## GENE

# MET

## ALTERATION

amplification - equivocal

## POTENTIAL TREATMENT STRATEGIES

On the basis of extensive clinical evidence, MET amplification or activating mutations may predict sensitivity to MET-targeting therapies such as kinase inhibitors crizotinib, capmatinib, tepotinib, and cabozantinib. A Phase 1 study for patients with MET-altered NSCLC treated with MET inhibitor bosutinib monotherapy reported an overall ORR of 30.6% (11/36) and DCR of 97.2% (35/36) with MET overexpression, amplification, and exon 14 skipping demonstrating ORRs of 35.7% (5/14), 41.2% (7/17), and 66.7% (10/15), respectively; increased ORRs were observed in patients with both exon 14 skipping and amplification (100%, 4/4) and with both amplification and overexpression (50%, 3/6)<sup>125</sup>. Crizotinib has benefited patients with MET-amplified NSCLC of varied histologies<sup>126-129</sup>, gastroesophageal cancer<sup>130</sup>, glioblastoma<sup>131</sup>, and carcinoma of unknown primary<sup>132</sup>. Capmatinib has demonstrated clinical efficacy for patients with MET-amplified NSCLC both as a monotherapy<sup>133-134</sup> and in combination with an EGFR-TKI for patients with concurrent activating EGFR mutations<sup>135-136</sup>. Tepotinib has demonstrated efficacy for patients with MET-amplified HCC<sup>137</sup> as a monotherapy, and in combination with gefitinib for patients with MET-amplified and EGFR-mutated NSCLC<sup>138-140</sup>.

Savolitinib elicited responses in patients with MET-amplified papillary renal cell carcinoma<sup>141</sup> and gastric cancer either alone or in combination with docetaxel<sup>142-143</sup>. AMG 337 elicited an ORR of 50% (5/10), including 1 CR, for patients with MET-amplified gastric, esophageal, or gastroesophageal junction cancer<sup>144</sup>. Patients with MET-amplified NSCLC<sup>145</sup> and gastric cancer<sup>146</sup> treated with the MET-targeting antibody onartuzumab (MetMab) achieved clinical responses. In addition, high MET expression has been suggested to predict patient response to therapy regimens including rilotumumab, a monoclonal HGF-targeting antibody, as well as emibetuzumab, a monoclonal MET-targeting antibody, combined with ramucirumab<sup>147</sup>. Telisotuzumab vedotin, a MET antibody-drug conjugate, was reported to be active in MET-positive NSCLC with an ORR of 18.8% (3/16) and a DCR of 56.3%<sup>148</sup>. Clinical and preclinical studies of lung cancer have shown that MET amplification is a common mechanism of resistance to EGFR inhibitors in first-line and later treatment settings<sup>98-104</sup>. Multiple studies have demonstrated that patients with a concurrent EGFR mutation and MET amplification, as recurrently observed at progression in patients with EGFR-mutated NSCLC on EGFR TKI, have benefited from a combination of MET- and EGFR-targeted therapies<sup>98,103-106</sup>.

## FREQUENCY & PROGNOSIS

MET amplification has been reported at incidences of 14-48% in non-small cell lung cancer (NSCLC), is correlated with increased MET protein expression, and occurs more frequently following treatment with EGFR

inhibitors<sup>109,145,149-155</sup>. In the Phase 2 VISION study of patients with NSCLC, MET amplification was reported in 4.9% of samples<sup>156</sup>. Studies on the effect of MET amplification on prognosis in NSCLC have yielded conflicting results<sup>109,149,153,157-161</sup>, although concurrent MET amplification and EGFR mutation have been correlated with reduced disease-free survival<sup>162</sup>. MET exon 14 splice alteration, which has predominantly been observed in lung cancer, was found to be an independent poor prognostic factor in a study of 687 patients with NSCLC<sup>163</sup>. However, other studies did not find MET exon 14 splice alteration as a major risk factor for overall survival for NSCLC patients, although recurrence rate was significantly higher in patients with exon 14 splice alteration compared to those with ALK fusion<sup>164-165</sup>. Among NSCLC patients with exon 14 alterations that had not been previously treated with a MET inhibitor, a non-significant trend for reduced survival was noted in the context of concurrent MET amplification (5.2 vs 10.5 months,  $p = 0.06$ )<sup>166</sup>.

## FINDING SUMMARY

MET encodes a receptor tyrosine kinase, also known as c-MET or hepatocyte growth factor receptor (HGFR), that is activated by the ligand HGF; MET activation results in signaling mediated partly by the RAS-RAF-MAPK and PI3K pathways to promote proliferation<sup>167-168</sup>. MET has been reported to be amplified in cancer<sup>169</sup>, with amplification positively correlating with protein expression in some cancer types<sup>149,170-173</sup> and associating with therapeutic response to MET inhibitors in a variety of cancer types<sup>126-128,130-132,174-175</sup>.

ORDERED TEST # ORD-1002521-02

## GENOMIC FINDINGS

## GENE

## HGF

## ALTERATION

amplification - equivocal

## POTENTIAL TREATMENT STRATEGIES

On the basis of several preclinical studies in different cancer types, high HGF gene expression may associate with sensitivity to MET-targeted therapies, such as the approved multikinase inhibitors crizotinib and cabozantinib<sup>176-180</sup>. However, this hypothesis has not been extensively tested in clinical studies. Whereas patients with glioblastoma and high tumor HGF gene expression experienced longer survival and a higher objective response rate (5/14 vs. 0/16) on the MET-targeting antibody onartuzumab combined with the anti-VEGF antibody

bevacizumab than with placebo plus bevacizumab<sup>181</sup>, tumor HGF gene expression did not predict significant benefit from onartuzumab added to the EGFR-inhibitor erlotinib for patients with non-small cell lung cancer<sup>182</sup>. Anti-HGF antibodies, such as ficlatuzumab, are also under clinical investigation<sup>183-184</sup>. Preclinical studies have shown that increased HGF protein levels can induce resistance of EGFR-mutant lung tumors to EGFR inhibitors and of BRAF-mutant melanoma cells to RAF inhibitors; this resistance could be overcome by combination therapy with MET inhibitors<sup>185-190</sup>.

## FREQUENCY &amp; PROGNOSIS

HGF mutation or amplification has been reported in 9% and 2% of lung adenocarcinomas, respectively<sup>107</sup>, and in 3% and 1% of lung squamous cell carcinomas, respectively<sup>108</sup>. HGF protein expression has been identified in 57% of lung adenocarcinoma samples in one study, and

associated with poor survival<sup>191</sup>. In patients with non-small cell lung cancer (NSCLC), including lung adenocarcinoma, low HGF protein levels and high HGF serum concentrations have been associated with longer and shorter overall survival, respectively<sup>192-193</sup>.

## FINDING SUMMARY

HGF encodes hepatocyte growth factor, also known as scatter factor, an activating ligand of the receptor tyrosine kinase MET. Certain splice isoforms of HGF may also act as MET antagonists<sup>194-195</sup>. HGF plays an important role in normal development, acting as a growth factor in a number of different tissues<sup>194-195</sup>. HGF and its receptor, MET, have been implicated in growth, invasion, and metastasis of many solid tumors<sup>195</sup>. HGF has been reported to be amplified in cancer<sup>169</sup>, and may be biologically relevant in this context<sup>196-197</sup>.

## GENE

## RICTOR

## ALTERATION

amplification

## POTENTIAL TREATMENT STRATEGIES

RICTOR amplification may indicate sensitivity to mTORC1/2 inhibitors<sup>198</sup> or dual PI3K/mTOR inhibitors<sup>199</sup>. A patient with RICTOR-amplified lung adenocarcinoma experienced SD for >18 months upon treatment with the dual mTORC1/2 inhibitor CC-223<sup>198</sup>, and a patient with RICTOR-amplified metastatic thymic carcinoma achieved a PR upon treatment with a pan-PI3K/mTORC1/2

inhibitor PQR309<sup>199</sup>. However, 4/4 patients with small cell lung cancer and RICTOR amplification did not achieve an objective response or SD (PFS of 1.25 months) from treatment with vistusertib<sup>200</sup>, and additional trials of vistusertib were terminated due to lack of efficacy<sup>142</sup>. RICTOR alterations, including amplification, have been implicated in resistance to the EGFR tyrosine kinase inhibitor erlotinib in patients with non-small cell lung carcinoma<sup>201</sup>.

## FREQUENCY &amp; PROGNOSIS

In a genomic study of 1,070 lung cancer cases, focal amplification of RICTOR was detected in 14.6% of small cell lung cancers (7/48), 8.7% of large cell neuroendocrine carcinomas (2/23), 8.4% of adenocarcinomas (61/724), and 7.4% of

squamous cell carcinomas (8/108)<sup>198</sup>. Published data investigating the prognostic implications of RICTOR alterations in lung cancer are limited (PubMed, Dec 2020). RICTOR amplification in lung cancer often co-occurs with mutations in KRAS, EGFR, or the PI3K-AKT-mTOR pathway, but has also been characterized as a driver alteration in lung cancer<sup>198</sup>.

## FINDING SUMMARY

RICTOR encodes an mTOR-binding protein that forms part of the rapamycin-insensitive mTORC2 complex, a regulator of cell metabolism and the cytoskeleton<sup>202-204</sup>. RICTOR amplification has been reported in cancer<sup>205</sup> and has been associated with clinical response to mTORC1/2 inhibition<sup>206-207</sup>.



ORDERED TEST # ORD-1002521-02

## GENOMIC FINDINGS

## GENE

## FGF10

## ALTERATION

amplification - equivocal

### POTENTIAL TREATMENT STRATEGIES

A preclinical study reported that FGF10-driven migration and invasion of pancreatic cancer cell lines could be blocked by inhibitory antibodies targeting FGFR2<sup>208</sup>, and a second study found that expression of dominant-negative FGFR1 or FGFR2 led to a decrease in tumor size in a prostate cancer xenograft model driven by FGF10, although the decrease was not statistically significant<sup>209</sup>. Clinical trials are ongoing for

multiple inhibitors that target FGFR2 and other kinases, including the approved agents pazopanib, ponatinib, and lenvatinib, as well as pan-FGFR inhibitors such as AZD4547, infigratinib, CH5183284, and TAS-120; however, these agents have not been comprehensively tested in the context of FGF10 amplification or overexpression.

### FREQUENCY & PROGNOSIS

Infrequent but recurrent amplification of FGF10 has been reported in multiple solid tumor types, including gallbladder cancer<sup>210</sup>, gastric cancer<sup>211</sup>, and esophageal squamous cell carcinoma (SCC)<sup>212</sup>; one small-scale study reported FGF10 amplification in 7/7 oral SCC cases<sup>213</sup>. Preclinical studies have shown that increased FGF10 expression and FGF10-FGFR1/2 signaling promotes cancer cell proliferation, invasion,

migration, and tumorigenesis in a variety of tumor models<sup>208-209,214-215</sup>.

### FINDING SUMMARY

FGF10 encodes fibroblast growth factor 10, a ligand that primarily binds to FGFR2, but also FGFR1<sup>216</sup>, with a broad range of functions in development and wound healing. FGF10 has been implicated in regulating the epithelial-mesenchymal transition in cancer cells<sup>217</sup> and during normal development<sup>218</sup>. Germline mutations in FGF10 have been implicated in aplasia of the lacrimal and salivary glands, an autosomal dominant developmental disorder<sup>219</sup>. Amplification of FGF10 has been reported in cancer<sup>169</sup> and may be biologically relevant in this context<sup>196-197</sup>.

## GENE

## PARP1

## ALTERATION

amplification

### POTENTIAL TREATMENT STRATEGIES

Multiple PARP inhibitors with activity against PARP1, including the approved therapies olaparib, niraparib, and rucaparib, are in clinical trials in solid and hematologic cancers, and have shown efficacy against cancers harboring inactivating mutations in DNA repair genes such as BRCA1/2<sup>220-228</sup>. A Phase 1/2 trial of olaparib and temozolomide in SCLC, which is characterized by strong expression of PARP1<sup>229</sup>, showed a superior response rate compared to historical data for temozolomide alone<sup>230</sup>. However, further work is required to ascertain the strength of association

between PARP1 amplification and expression, as well as between PARP1 amplification or expression and efficacy of PARP inhibitors. On the basis of preclinical studies, PARP1 mutations are not predicted to confer sensitivity to PARP inhibitors<sup>231-233</sup>.

### FREQUENCY & PROGNOSIS

PARP1 mutations have been reported in 1% of solid tumors, including in 8% of nonmelanoma skin cancers and 3% each of endometrial cancer, anal cancer, melanoma, and small bowel cancer samples<sup>234</sup>. PARP1 amplification is less frequent, reported in 0.3% of cases, with highest incidence of 3% in anal cancer and 2% in endometrial cancer<sup>234</sup>. In one study, high expression of PARP1 was associated with shorter overall survival of patients with classical GBM, but not in other GBM subtypes<sup>235</sup>. In the context of lung cancer, neuroendocrine tumors have been shown to express PARP1 at higher levels than

adenocarcinomas and squamous cell carcinomas, with highest expression seen in small cell lung cancer (SCLC)<sup>236</sup>.

### FINDING SUMMARY

PARP1 encodes the dominant member of the poly(ADP-ribose) polymerase (PARP) family that plays roles in DNA damage repair (DDR) and cell cycle progression<sup>237-238</sup>. Several missense mutations in PARP1 have been reported in cancer, including the activating mutation L713F<sup>231</sup> and the hypomorphic variants F304L, V762A, and E988K<sup>231,239-240</sup>. PARP1 amplification has been reported as a rare but recurrent event in glioblastoma multiforme (GBM), where it was found to be associated with increased expression of PARP1 as well as with higher tumor grade<sup>235</sup>. Limited and conflicting data have been reported on the potential roles of PARP1 germline mutations in cancer predisposition<sup>231,239-240</sup>.

ORDERED TEST # ORD-1002521-02

## GENOMIC FINDINGS

## GENE

## TP53

## ALTERATION

splice site 376-1G&gt;A

## TRANSCRIPT ID

NM\_000546

## CODING SEQUENCE EFFECT

376-1G&gt;A

## VARIANT ALLELE FREQUENCY (% VAF)

47.9%

## POTENTIAL TREATMENT STRATEGIES

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib<sup>241-244</sup>, or p53 gene therapy and immunotherapeutics such as SGT-53<sup>245-249</sup> and ALT-801<sup>250</sup>. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% (17/176) and SDs in 53.4% (94/176) of patients with solid tumors; the response rate was 21.1% (4/19) in patients with TP53 mutations versus 12.1% (4/33) in patients who were TP53 wild-type<sup>251</sup>. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 31.9% (30/94, 3 CR) ORR and a 73.4% (69/94) DCR in patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer<sup>252</sup>. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 42.9% (9/21, 1 CR) ORR and a 76.2% (16/21) DCR in patients with platinum-refractory TP53-mutated ovarian cancer<sup>253</sup>. The combination of adavosertib with paclitaxel and carboplatin in patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone<sup>254</sup>. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24.0% (6/25) ORR with adavosertib combined with paclitaxel<sup>142</sup>. A Phase 1 trial of neoadjuvant

adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71.4% (5/7) response rate for patients with TP53 alterations<sup>255</sup>. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel in patients with solid tumors, 75.0% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage<sup>249</sup>. Additionally, the combination of a CHK1 inhibitor and irinotecan reportedly reduced tumor growth and prolonged survival in a TP53-mutated, but not TP53-wild-type, breast cancer xenotransplant mouse model<sup>256</sup>. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies<sup>257-258</sup>; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies<sup>259-260</sup>. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

## FREQUENCY &amp; PROGNOSIS

TP53 is one of the most commonly mutated genes in lung cancer; mutations have been reported in 43-80% of non-small cell lung cancers (NSCLCs)<sup>107-108,261-266</sup>, including 38-54% of lung adenocarcinomas and 47-83% of lung squamous cell carcinomas (cBioPortal, COSMIC, Sep 2020)<sup>47-48,107-108</sup>. In one study of 55 patients with lung adenocarcinoma, TP53 alterations correlated with immunogenic features including PD-L1 expression, tumor mutation burden and neoantigen presentation; likely as a consequence of this association TP53 mutations correlated with improved clinical outcomes to PD-1 inhibitors pembrolizumab and nivolumab in this study<sup>267</sup>. Mutations in TP53 have been associated with lymph node metastasis in patients with lung adenocarcinoma<sup>268</sup>. Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic

mutations that allow for clonal expansion<sup>269-274</sup>. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy<sup>269-270</sup>. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease<sup>275</sup>. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH<sup>273,276-277</sup>. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

## FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers<sup>278</sup>. Alterations such as seen here may disrupt TP53 function or expression<sup>279-283</sup>.

## POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters with no conflicts) associated with Li-Fraumeni syndrome (ClinVar, Sep 2020)<sup>284</sup>. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers<sup>285-287</sup>, including sarcomas<sup>288-289</sup>. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000<sup>290</sup> to 1:20,000<sup>289</sup>. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30<sup>291</sup>. In the appropriate clinical context, germline testing of TP53 is recommended.



ORDERED TEST # ORD-1002521-02

## THERAPIES WITH CLINICAL BENEFIT

## IN PATIENT'S TUMOR TYPE

## Afatinib

*Assay findings association*
**EGFR**  
amplification

### AREAS OF THERAPEUTIC USE

Afatinib is an irreversible kinase inhibitor that targets the kinase domains of EGFR, ERBB2/HER2, and ERBB4. It is FDA approved for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) and nonresistant EGFR mutations and for the treatment of patients with metastatic, squamous NSCLC after progression on platinum-based chemotherapy. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

EGFR activating mutations or amplification may indicate sensitivity to afatinib. In Phase 2 studies of afatinib, patients with EGFR-amplified NSCLC achieved an objective response rate of 20% (5/25) and a disease-control rate of 64% (16/25)<sup>71</sup>, and 2/5 patients with EGFR amplification in other solid tumors experienced stable disease<sup>72</sup>.

### SUPPORTING DATA

Afatinib enabled 1 PR and 1 SD for 2 patients with EGFR-amplified NSCLC in a Phase 2 study<sup>292</sup>. Extensive clinical data have demonstrated that afatinib is effective for patients with EGFR-mutated advanced NSCLC, including exon 19 deletions and L858 mutations, as well as uncommon sensitizing mutations in exons 18 or 20<sup>293-299</sup>. Afatinib has also shown activity for patients with advanced NSCLC and ERBB2 mutations, most of which were exon 20 insertions<sup>292,300-308</sup>. The randomized Phase 3 LUX-Lung 8 trial comparing afatinib with erlotinib as second-line therapy for advanced lung squamous cell carcinoma reported significantly longer median OS (7.9 vs. 6.8 months, HR=0.81), significantly longer median PFS (2.6 vs. 1.9 months, HR=0.81), and higher DCR (51% vs. 40%, p=0.002) for patients treated with afatinib<sup>298</sup>. For patients who progressed on afatinib monotherapy, additional clinical benefit has been reported from afatinib combined with paclitaxel<sup>309</sup>.

## Capmatinib

*Assay findings association*
**MET**  
amplification - equivocal

### AREAS OF THERAPEUTIC USE

Capmatinib is a Type Ib MET inhibitor that is FDA approved to treat patients with metastatic non-small cell lung cancer harboring MET exon 14 skipping-associated alterations.

### GENE ASSOCIATION

On the basis of clinical data in NSCLC<sup>133,138-140,310</sup>, HCC<sup>137</sup>, RCC<sup>141</sup>, and gastric cancer<sup>142</sup>, MET amplification may predict sensitivity to Type 1b MET inhibitors.

### SUPPORTING DATA

In the Phase 2 GEOMETRY mono-1 study for patients with advanced NSCLC and MET gene copy number (GCN) ≥10, capmatinib elicited ORRs of 29-40%, median PFS of 4.1-4.2 months, and median OS of 9.6-10.6 months across treatment-naïve and previously treated cohorts<sup>311</sup>.

A Phase 1 study of capmatinib monotherapy for advanced EGFR- and ALK-wild-type NSCLC reported ORRs of 46.7% (7/15) for patients with MET GCN ≥6, 25% (3/12) for patients with MET GCN 4-6, and 5.9% (1/17) for patients with MET GCN <4; median PFS was 3.7 months overall, and 7.9 months for patients with MET GCN ≥6<sup>312</sup>. Phase 1b/2 trial of capmatinib and nazartinib for patients with EGFR-mutated, EGFR-TKI-resistant NSCLC and unknown MET status reported a 42% (14/33, 2 CRs) ORR, with no correlation observed between responses and T790M status<sup>313</sup>. Multiple Phase 1 and 2 clinical studies have reported limited efficacy for capmatinib monotherapy in non-NSCLC indications, with no responses observed for patients with glioblastoma (n=10)<sup>314</sup>, gastric cancer (n=9), or other advanced solid tumors (n=24)<sup>315-316</sup>.

ORDERED TEST # ORD-1002521-02

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

## Crizotinib

*Assay findings association*
**MET**  
amplification - equivocal

### AREAS OF THERAPEUTIC USE

Crizotinib is an inhibitor of the kinases MET, ALK, ROS1, and RON. It is FDA approved to treat patients with ALK rearrangement- or ROS1 rearrangement-positive non-small cell lung cancer (NSCLC), and to treat pediatric and young adult patients with ALK rearrangement-positive anaplastic large cell lymphoma (ALCL). Please see the drug label for full prescribing information.

### GENE ASSOCIATION

Sensitivity of MET alterations to crizotinib is suggested by extensive clinical data in patients with MET-amplified cancers, including non-small cell lung cancer (NSCLC)<sup>126-128,317-318</sup>, gastric cancer<sup>174</sup>, gastroesophageal cancer<sup>130</sup>, glioblastoma<sup>131</sup>, and carcinoma of unknown primary<sup>132</sup>, as well as in patients with MET-mutated cancers, including NSCLC<sup>319-324</sup>, renal cell carcinoma (RCC)<sup>325</sup>, and histiocytic sarcoma<sup>319</sup>. Crizotinib has also benefited patients with NSCLC or histiocytic sarcoma tumors harboring various alterations associated with MET exon 14 skipping<sup>166,319-320,322-324</sup>.

### SUPPORTING DATA

In a small study for patients with NSCLC and MET

overexpression with or without gene amplification, crizotinib elicited 11 PRs and 3 SDs in 19 evaluable patients<sup>318</sup>. Crizotinib has demonstrated efficacy in patients with NSCLC and ALK rearrangements<sup>326-330</sup>, ROS1 rearrangements<sup>331-335</sup>, an NTRK1 fusion<sup>336</sup>, or MET activation<sup>126-128,317-318,320-324,337-343</sup>. The Phase 2 METROS and AcSe trials have reported ORRs of 31.3% to 32.0%, median PFS of 3.2 to 5.0 months, and median OS of 5.4 to 7.7 months for patients with MET amplified advanced non-small cell lung cancer (NSCLC); a higher level of amplification was predictive of better response in the AcSe trial ( $P=0.04$ )<sup>331,344</sup>. Additional patients with MET amplified NSCLC have been reported to experience clinical benefit from crizotinib in several case studies<sup>126-128,340,343,345</sup>. A patient with lung adenocarcinoma harboring K860I and L858R EGFR mutations, who acquired both EGFR T790M and MET amplification upon various treatments, experienced clinical benefit from subsequent combination treatment of osimertinib and crizotinib<sup>106</sup>. Two patients with ALK-positive NSCLC and acquired MET amplification experienced benefit from crizotinib monotherapy and crizotinib in combination with lorlatinib<sup>346</sup>.

## Dacomitinib

*Assay findings association*
**EGFR**  
amplification

### AREAS OF THERAPEUTIC USE

Dacomitinib is a second generation irreversible tyrosine kinase inhibitor that targets the kinase domains of EGFR, ERBB2/HER2, and ERBB4/HER4. It is FDA approved for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R substitution mutations. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

On the basis of clinical<sup>347-349</sup> and preclinical<sup>350-351</sup> data, EGFR amplification or activating mutation may indicate sensitivity to dacomitinib.

### SUPPORTING DATA

A randomized Phase 3 trial in patients with NSCLC with activating EGFR mutations (primarily L858R or exon 19 deletions) reported improved clinical benefit with first-line dacomitinib compared with gefitinib (median OS, 34.1 vs. 26.8 months, HR=0.760; median PFS, 14.7 vs. 9.2 months, HR=0.59)<sup>347,352</sup>; median OS was 34.1 to 36.7

months and ORR was 74.9% to 79.3%, depending on the dosing regimen<sup>353</sup>. A pooled subgroup analysis of patients with NSCLC with activating EGFR mutations reported improved clinical efficacy with dacomitinib treatment compared with erlotinib (median PFS, 14.6 vs. 9.6 months, HR=0.717; median OS, 26.6 vs. 23.2 months, HR=0.737)<sup>354</sup>. Reduced efficacy of dacomitinib treatment in patients with NSCLC harboring the EGFR T790M mutation has been reported in multiple studies<sup>355-357</sup>. A Phase 1 trial of combination dacomitinib and a MEK1/2 inhibitor for patients with KRAS-mutated CRC, NSCLC, or pancreatic cancer reported 20/36 SDs and 16 PDs, however toxicity from this combination prevented long-term treatment in this patient population<sup>358</sup>. A Phase 2 study of dacomitinib in patients with NSCLC who had been previously treated with chemotherapy or erlotinib and were not selected for EGFR mutations reported an ORR of 4.5% (3/66)<sup>356</sup>. In one study, the combination of dacomitinib and crizotinib was ineffective and associated with high toxicity in patients with NSCLC<sup>359</sup>.

ORDERED TEST # ORD-1002521-02

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

## Erlotinib

*Assay findings association*
**EGFR**  
amplification

### AREAS OF THERAPEUTIC USE

Erlotinib is a small-molecule inhibitor of EGFR. It is FDA approved as a monotherapy or in combination with ramucirumab for patients with metastatic non-small cell lung cancer (NSCLC) harboring EGFR exon 19 deletions or exon 21 (L858R) mutations. Erlotinib is also FDA approved in combination with gemcitabine as a first-line treatment for advanced pancreatic cancer. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

Amplification or activation of EGFR may predict sensitivity to therapies such as erlotinib. In a prospective study of advanced NSCLC treated with gefitinib (n=102),

EGFR copy gain was significantly associated with improved survival (HR=0.44)<sup>70</sup>. Several meta-analyses spanning 14 to 20 studies of patients with advanced NSCLC receiving single-agent erlotinib or gefitinib (n=1725 to 1854) reported the association of increased EGFR copy number with improved OS (HR=0.72 to 0.77), although the survival benefit was not observed for East Asian populations (HR=0.79 to 1.11)<sup>68-69,360</sup>.

### SUPPORTING DATA

The Phase 3 BR.21 trial demonstrated prolonged OS for genomically unselected patients with NSCLC treated with erlotinib compared with those treated with standard chemotherapy<sup>361</sup>.

## Gefitinib

*Assay findings association*
**EGFR**  
amplification

### AREAS OF THERAPEUTIC USE

Gefitinib targets the tyrosine kinase EGFR and is FDA approved to treat non-small cell lung cancer (NSCLC) harboring exon 19 deletions or exon 21 (L858R) substitution mutations in EGFR. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

Amplification or activation of EGFR may predict sensitivity to therapies such as gefitinib. Clinical studies have consistently shown significant improvement in response rates and progression-free survival for patients with EGFR-mutated NSCLC treated with gefitinib, compared to chemotherapy<sup>362-368</sup>. In a prospective study of advanced NSCLC treated with gefitinib (n=102), EGFR copy gain was significantly associated with improved survival (HR=0.44)<sup>70</sup>. Several meta-analyses spanning 14 to 20 studies of patients with advanced NSCLC receiving single-agent erlotinib or gefitinib (n=1725 to 1854) reported the association of increased EGFR copy number with improved OS (HR=0.72 to 0.77), although the survival benefit was not observed for East Asian populations (HR=0.79 to 1.11)<sup>68-69,360</sup>. Patients with refractory advanced esophageal carcinoma and EGFR amplification derived significant overall survival benefit from gefitinib compared to placebo (HR = 0.21)<sup>369-370</sup>.

### SUPPORTING DATA

In patients with EGFR-mutated NSCLC who progressed on 1st or 2nd generation EGFR TKIs, combination of gefitinib with the MET inhibitor capmatinib achieved ORRs of 32-47% and DCRs of 74-75% in cohorts with MET amplification or overexpression<sup>105</sup>. In this same setting, gefitinib in combination with the MET inhibitor tepotinib elicited the largest benefit in patients with MET amplification or high-level MET overexpression<sup>371</sup>; in the

cohort with MET amplification, gefitinib with tepotinib significantly improved ORR (75.0% vs. 42.9%, OR = 4.00) and median PFS (19.8 vs. 5.5 months, HR = 0.25) as compared with pemetrexed and platinum chemotherapy<sup>372</sup>. Gefitinib achieved an ORR of 69.8% and an OS of 19.2 months as first-line treatment for Caucasian patients with non-small cell lung carcinoma (NSCLC) and EGFR sensitizing mutations<sup>373</sup>. In the retrospective analysis of a Phase 3 study for East Asian patients, gefitinib was reported to have a longer PFS for patients with EGFR mutation-positive NSCLC compared with carboplatin/paclitaxel doublet chemotherapy<sup>365,374</sup>. Two Phase 3 trials of gefitinib plus pemetrexed and carboplatin compared with gefitinib alone for patients with advanced NSCLC harboring EGFR activating mutations reported significantly higher ORRs (75.3% and 84% vs. 62.5% and 67%), longer median PFSs (16 and 20.9 months vs. 8 and 11.9 months), and longer median OSs (50.9 months and not reached vs. 17 and 38.8 months) with combination treatment; however, combination treatment was associated with increased Grade 3 or higher adverse events<sup>375-376</sup>. Retrospective analysis of East Asian patients with advanced NSCLC receiving first-line gefitinib therapy reported that patients with EGFR exon 19 mutations experienced a longer median PFS (10.9 months) compared with patients with EGFR mutations in exon 18 (7.9 months), exon 20 (1.2 months), exon 21 (7.7 months), or double mutations (5.7 months); however, no differences in OS were seen between EGFR mutations<sup>377</sup>. In a Phase 1 study for treatment-naïve patients with NSCLC, best ORRs of 78% (7/9) were observed in patients treated with combination gefitinib and the PD-L1 inhibitor durvalumab as first-line treatment and of 80% (8/10) in those treated with the combination after gefitinib monotherapy<sup>378</sup>.

© 2021 Foundation Medicine, Inc. All rights reserved.

ORDERED TEST # ORD-1002521-02

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

## Cabozantinib

*Assay findings association*
**MET**  
amplification - equivocal

### AREAS OF THERAPEUTIC USE

Cabozantinib inhibits multiple tyrosine kinases, including MET, RET, VEGFRs, and ROS1. It is FDA approved to treat patients with advanced renal cell carcinoma (RCC), hepatocellular carcinoma (HCC) after prior treatment with sorafenib, or progressive, metastatic medullary thyroid cancer (MTC). Please see the drug label for full prescribing information.

### GENE ASSOCIATION

Sensitivity of MET alterations to cabozantinib is suggested by clinical responses in patients with non-small cell lung cancer (NSCLC) harboring MET mutations associated with MET exon 14 skipping, with or without concurrent MET amplification<sup>320,379</sup>, as well as by extensive preclinical data<sup>380-386</sup>.

### SUPPORTING DATA

Cabozantinib elicited a CR in a patient with lung adenocarcinoma harboring a MET amplification and a mutation affecting MET exon 14 splicing<sup>320</sup>. A Phase 2 randomized discontinuation trial of cabozantinib reported a 10.0% (6/60) ORR and a 58.3% (35/60) DCR, with median PFS of 4.2 months, for patients with genomically unselected, heavily pretreated NSCLC<sup>387</sup>. Patients with EGFR wild-type non-squamous NSCLC who had progressed after previous treatment experienced longer median PFS with cabozantinib alone or combined with erlotinib (4.3 and 4.7 months, HR=0.39 and 0.37, respectively) compared with single agent erlotinib (1.8 months) in a randomized Phase 2 trial<sup>388</sup>. A Phase 1 study of cabozantinib for advanced solid tumors reported an ORR of 20.0% (4/20; 4 PRs, all in EGFR-mutated tumors) and DCR of 100% (20/20) in the expansion cohort for Japanese patients with NSCLC<sup>389</sup>.

## Cetuximab

*Assay findings association*
**EGFR**  
amplification

### AREAS OF THERAPEUTIC USE

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS-wild-type, EGFR-expressing metastatic colorectal cancer (CRC). Please see the drug label for full prescribing information.

### GENE ASSOCIATION

For patients with metastatic CRC receiving cetuximab or panitumumab as mono- or combination therapy, increased EGFR copy number associated with improved OS (HR=0.62) in a meta-analysis, although increased survival was not seen in populations that received first-line treatment with EGFR antibodies<sup>64</sup>.

### SUPPORTING DATA

In previously untreated patients with non-small cell lung cancer (NSCLC), the FLEX study demonstrated that in NSCLC tumors with high expression of EGFR, treatment with cetuximab plus chemotherapy resulted in longer overall survival compared to chemotherapy alone; there was no clear association between cetuximab response and EGFR mutations in this trial<sup>63</sup>. In a Phase 2 study of 31 patients with Stage 3 NSCLC, the addition of cetuximab to radiotherapy and chemotherapy produced an overall

response rate of 67%; EGFR gene copy number was not predictive of efficacy outcome<sup>390</sup>. A Phase 3 study of 938 patients with progressive non-small cell lung cancer after platinum-based therapy concluded that, in unselected patients, the addition of cetuximab to chemotherapy was not recommended in this second-line setting<sup>391</sup>. Cetuximab is also being studied as part of a therapeutic regimen for patients with EGFR mutations who develop secondary resistance to erlotinib or gefitinib. A Phase 1b study combining afatinib and the anti-EGFR antibody cetuximab in patients with advanced EGFR-mutant lung cancer with acquired resistance to erlotinib/ gefitinib observed an overall objective response rate of 29%, and comparable response rates in both T790M-positive and T790M-negative tumors (32% vs. 25%)<sup>392</sup>. A Phase 1 study of combination erlotinib and cetuximab treatment in patients with NSCLC, including those with squamous tumors, inhibitor-resistant EGFR mutations, and wild-type EGFR, as well as those who had progressed on prior erlotinib treatment, reported partial responses in two of 20 patients and stable disease lasting at least 6 months in three of 20 patients<sup>393</sup>; however, in this study a patient identified with an exon 19 deletion and T790M progressed rapidly on cetuximab and erlotinib<sup>394</sup>.

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Donna Ferguson, M.D. | 01 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1002521-02

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

## Panitumumab

*Assay findings association*
**EGFR**  
amplification

### AREAS OF THERAPEUTIC USE

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved to treat KRAS wild-type and NRAS wild-type metastatic colorectal cancer (CRC) combined with chemotherapy or as monotherapy for patients who have progressed on prior chemotherapy. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

For patients with metastatic CRC receiving cetuximab or panitumumab as mono- or combination therapy, increased EGFR copy number associated with improved OS (HR=0.62) in a meta-analysis, although increased survival was not seen in populations that received first-

line treatment with EGFR antibodies<sup>64</sup>.

### SUPPORTING DATA

In a Phase 2 trial in patients with advanced non-small cell lung cancer (NSCLC), the addition of panitumumab to paclitaxel/carboplatin did not result in improved clinical benefit<sup>395</sup>, and subsequent studies investigating the addition of panitumumab to pemetrexed/cisplatin reported no benefit for patients with wild-type KRAS lung adenocarcinoma<sup>396</sup>. The combination of afatinib and panitumumab has been explored for 2 patients with EGFR T790M NSCLC, with 1 partial response reported<sup>397</sup>.

**NOTE** Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**

**NOTE** Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.foundationmedicine.com/genomic-testing#support-services). Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

## GENE **EGFR**

### ALTERATION amplification

**RATIONALE**  
EGFR activating mutations, rearrangements, or amplification may predict sensitivity to EGFR-targeted therapies. Several strategies to overcome

resistance are under investigation, including next-generation EGFR TKIs and EGFR inhibitor combinations.

#### **NCT03137771**

**PHASE 2**

Maintenance Chemotherapy With or Without Stereotactic Body Radiation Therapy in Treating Patients With Stage IV Non-small Cell Lung Cancer

**TARGETS**  
EGFR, PD-1

**LOCATIONS:** Florida, Georgia, South Carolina, Louisiana

#### **NCT02693535**

**PHASE 2**

TAPUR: Testing the Use of Food and Drug Administration (FDA) Approved Drugs That Target a Specific Abnormality in a Tumor Gene in People With Advanced Stage Cancer

**TARGETS**  
VEGFRs, ABL, SRC, ALK, AXL, MET, ROS1, TRKA, TRKC, CDK4, CDK6, CSF1R, FLT3, KIT, RET, mTOR, EGFR, ERBB2, ERBB3, MEK, BRAF, SMO, DDR2, PARP, PD-1, CTLA-4, ERBB4

**LOCATIONS:** Florida, Texas, Georgia, Alabama, North Carolina, Virginia, Oklahoma, Pennsylvania, Indiana

#### **NCT02795156**

**PHASE 2**

Study to Assess the Activity of Molecularly Matched Targeted Therapies in Select Tumor Types Based on Genomic Alterations

**TARGETS**  
BRAF, KIT, RET, VEGFRs, EGFR, ERBB2, ERBB4, MET, ROS1

**LOCATIONS:** Florida, Tennessee, Missouri, Wisconsin, Colorado

#### **NCT02716116**

**PHASE 1/2**

A Trial of AP32788 in Non-Small Cell Lung Cancer

**TARGETS**  
EGFR, ERBB2

**LOCATIONS:** Florida, Georgia, North Carolina, Virginia, Arizona, California

#### **NCT03829436**

**PHASE 1**

TPST-1120 as Monotherapy and in Combination With (Nivolumab, Docetaxel or Cetuximab) in Subjects With Advanced Cancers

**TARGETS**  
PD-1, PPARalpha, EGFR

**LOCATIONS:** Florida, North Carolina, Tennessee, Oklahoma, Maryland, Pennsylvania, New York, Massachusetts, Michigan



ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**NCT02609776**
**PHASE 1**

A Dose Escalation Study of JNJ-61186372 in Participants With Advanced Non-Small Cell Lung Cancer

**TARGETS**  
MET, EGFR

**LOCATIONS:** Florida, Virginia, Maryland, Pennsylvania, Missouri, New York, Massachusetts, Michigan, Illinois, Toronto (Canada)

**NCT03783403**
**PHASE 1**

A Study of CC-95251, a Monoclonal Antibody Directed Against SIRP $\alpha$ , in Subjects With Advanced Solid and Hematologic Cancers

**TARGETS**  
CD20, EGFR, SIRP-alpha

**LOCATIONS:** Texas, Alabama, North Carolina, Tennessee, Oklahoma, Pennsylvania, Toronto (Canada), Arizona

**NCT02099058**
**PHASE 1**

A Phase 1/1b Study With ABBV-399, an Antibody Drug Conjugate, in Subjects With Advanced Solid Cancer Tumors

**TARGETS**  
MET, EGFR, PD-1

**LOCATIONS:** Texas, Tennessee, Virginia, New Jersey, Massachusetts, Michigan, Illinois, Colorado, California

**NCT01553942**
**PHASE 2**

Afinib With CT and RT for EGFR-Mutant NSCLC

**TARGETS**  
EGFR, ERBB2, ERBB4

**LOCATIONS:** Massachusetts

**NCT02947386**
**PHASE 1/2**

Nimotuzumab and Nivolumab in Treating Patients With Advanced Non-small Cell Lung Cancer

**TARGETS**  
EGFR, PD-1

**LOCATIONS:** New York

ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**GENE**  
**HGF**
**RATIONALE**  
HGF amplification or activating mutations may predict sensitivity to therapeutic agents targeting

its receptor, MET, or to agents directly targeting HGF.

**ALTERATION**  
amplification - equivocal

**NCT03906071**
**PHASE 3**

Phase 3 Study of Sitravatinib Plus Nivolumab vs Docetaxel in Patients With Advanced Non-Squamous NSCLC

**TARGETS**  
PD-1, AXL, DDR2, FLT3, KIT, MET, PDGFRA, RET, TRKA, TRKB, VEGFRs

**LOCATIONS:** Florida, Louisiana

**NCT03175224**
**PHASE 1/2**

CBT-101 Study for Advanced Solid Tumors and c-Met Dysregulation

**TARGETS**  
MET

**LOCATIONS:** Rio Piedras (Puerto Rico), Florida, Louisiana, South Carolina

**NCT04310007**
**PHASE 2**

Testing the Addition of the Pill Chemotherapy, Cabozantinib, to the Standard Immune Therapy Nivolumab Compared to Standard Chemotherapy for Non-small Cell Lung Cancer

**TARGETS**  
MET, RET, ROS1, VEGFRs, PD-1

**LOCATIONS:** Florida, Louisiana, Georgia, South Carolina, Texas, North Carolina

**NCT02795156**
**PHASE 2**

Study to Assess the Activity of Molecularly Matched Targeted Therapies in Select Tumor Types Based on Genomic Alterations

**TARGETS**  
BRAF, KIT, RET, VEGFRs, EGFR, ERBB2, ERBB4, MET, ROS1

**LOCATIONS:** Florida, Tennessee, Missouri, Wisconsin, Colorado

**NCT03170960**
**PHASE 1/2**

Study of Cabozantinib in Combination With Atezolizumab to Subjects With Locally Advanced or Metastatic Solid Tumors

**TARGETS**  
PD-L1, MET, RET, ROS1, VEGFRs

**LOCATIONS:** Florida, Louisiana, South Carolina, Texas, Georgia, Virginia

**NCT02414139**
**PHASE 2**

Clinical Study of Oral cMET Inhibitor INC280 in Adult Patients With EGFR Wild-type Advanced Non-small Cell Lung Cancer

**TARGETS**  
MET

**LOCATIONS:** Florida, Georgia, Texas, South Carolina, North Carolina, Arkansas, Tennessee, Virginia, District of Columbia

ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**NCT04173338**
**PHASE 1**

Cabozantinib With Pemetrexed in Advanced Non-small Cell Lung Cancer, Urothelial Cancer and Malignant Mesothelioma

**TARGETS**  
MET, RET, ROS1, VEGFRs

**LOCATIONS:** Georgia

**NCT02099058**
**PHASE 1**

A Phase 1/1b Study With ABBV-399, an Antibody Drug Conjugate, in Subjects With Advanced Solid Cancer Tumors

**TARGETS**  
MET, EGFR, PD-1

**LOCATIONS:** Texas, Tennessee, Virginia, New Jersey, Massachusetts, Michigan, Illinois, Colorado, California

**NCT04139317**
**PHASE 2**

Safety and Efficacy of Capmatinib (INC280) Plus Pembrolizumab vs Pembrolizumab Alone in NSCLC With PD-L1 ≥ 50%

**TARGETS**  
MET, PD-1

**LOCATIONS:** Tennessee, Madrid (Spain), Valencia (Spain), Barcelona (Spain), Badalona (Spain), Toulouse Cedex 9 (France), LILLE Cédex (France), Bruxelles (Belgium), Yvoir (Belgium), Liege (Belgium)

**NCT01639508**
**PHASE 2**

Cabozantinib in Patients With RET Fusion-Positive Advanced Non-Small Cell Lung Cancer and Those With Other Genotypes: ROS1 or NTRK Fusions or Increased MET or AXL Activity

**TARGETS**  
MET, RET, ROS1, VEGFRs

**LOCATIONS:** New Jersey, New York

ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**GENE**  
**MET**
**RATIONALE**  
Activation of MET may lead to increased MET expression and activation and may therefore

confer sensitivity to MET inhibitors.

**ALTERATION**  
amplification - equivocal

**NCT03906071**
**PHASE 3**

Phase 3 Study of Sitravatinib Plus Nivolumab vs Docetaxel in Patients With Advanced Non-Squamous NSCLC

**TARGETS**  
PD-1, AXL, DDR2, FLT3, KIT, MET, PDGFRA, RET, TRKA, TRKB, VEGFRs

**LOCATIONS:** Florida, Louisiana

**NCT03175224**
**PHASE 1/2**

CBT-101 Study for Advanced Solid Tumors and c-Met Dysregulation

**TARGETS**  
MET

**LOCATIONS:** Rio Piedras (Puerto Rico), Florida, Louisiana, South Carolina

**NCT04310007**
**PHASE 2**

Testing the Addition of the Pill Chemotherapy, Cabozantinib, to the Standard Immune Therapy Nivolumab Compared to Standard Chemotherapy for Non-small Cell Lung Cancer

**TARGETS**  
MET, RET, ROS1, VEGFRs, PD-1

**LOCATIONS:** Florida, Louisiana, Georgia, South Carolina, Texas, North Carolina

**NCT02795156**
**PHASE 2**

Study to Assess the Activity of Molecularly Matched Targeted Therapies in Select Tumor Types Based on Genomic Alterations

**TARGETS**  
BRAF, KIT, RET, VEGFRs, EGFR, ERBB2, ERBB4, MET, ROS1

**LOCATIONS:** Florida, Tennessee, Missouri, Wisconsin, Colorado

**NCT03170960**
**PHASE 1/2**

Study of Cabozantinib in Combination With Atezolizumab to Subjects With Locally Advanced or Metastatic Solid Tumors

**TARGETS**  
PD-L1, MET, RET, ROS1, VEGFRs

**LOCATIONS:** Florida, Louisiana, South Carolina, Texas, Georgia, Virginia

**NCT02414139**
**PHASE 2**

Clinical Study of Oral cMET Inhibitor INC280 in Adult Patients With EGFR Wild-type Advanced Non-small Cell Lung Cancer

**TARGETS**  
MET

**LOCATIONS:** Florida, Georgia, Texas, South Carolina, North Carolina, Arkansas, Tennessee, Virginia, District of Columbia

ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**NCT02609776**
**PHASE 1**

A Dose Escalation Study of JNJ-61186372 in Participants With Advanced Non-Small Cell Lung Cancer

**TARGETS**  
MET, EGFR

**LOCATIONS:** Florida, Virginia, Maryland, Pennsylvania, Missouri, New York, Massachusetts, Michigan, Illinois, Toronto (Canada)

**NCT03539536**
**PHASE 2**

Study of Telisotuzumab Vedotin (ABBV-399) in Subjects With Previously Treated c-Met+ Non-Small Cell Lung Cancer

**TARGETS**  
MET

**LOCATIONS:** Alabama, Texas, Tennessee, Kentucky, Arkansas, Virginia, Missouri, Pennsylvania

**NCT04173338**
**PHASE 1**

Cabozantinib With Pemetrexed in Advanced Non-small Cell Lung Cancer, Urothelial Cancer and Malignant Mesothelioma

**TARGETS**  
MET, RET, ROS1, VEGFRs

**LOCATIONS:** Georgia

**NCT02099058**
**PHASE 1**

A Phase 1/1b Study With ABBV-399, an Antibody Drug Conjugate, in Subjects With Advanced Solid Cancer Tumors

**TARGETS**  
MET, EGFR, PD-1

**LOCATIONS:** Texas, Tennessee, Virginia, New Jersey, Massachusetts, Michigan, Illinois, Colorado, California

ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**GENE**  
**RICTOR**
**ALTERATION**  
amplification

**RATIONALE**  
RICTOR amplification may predict sensitivity to dual mTORC1/mTORC2 inhibitors, as well as dual

PI3K/mTOR inhibitors.

**NCT02159989**
**PHASE 1**

Sapanisertib and Ziv-Aflibercept in Treating Patients With Recurrent Solid Tumors That Are Metastatic or Cannot Be Removed by Surgery

**TARGETS**  
PIGF, VEGFA, VEGFB, mTORC1, mTORC2

**LOCATIONS:** Texas

**NCT03366103**
**PHASE 1/2**

Navitoclax and Vistusertib in Treating Patients With Relapsed Small Cell Lung Cancer and Other Solid Tumors

**TARGETS**  
mTORC1, mTORC2, BCL-W, BCL-XL, BCL2

**LOCATIONS:** Maryland, New Jersey, New York

**NCT03017833**
**PHASE 1**

Sapanisertib and Metformin in Treating Patients With Advanced or Metastatic Relapsed or Refractory Cancers

**TARGETS**  
mTORC1, mTORC2

**LOCATIONS:** Texas

**NCT03430882**
**PHASE 1**

Sapanisertib, Carboplatin, and Paclitaxel in Treating Patients With Recurrent or Refractory Malignant Solid Tumors

**TARGETS**  
mTORC1, mTORC2

**LOCATIONS:** Texas

**NCT04250545**
**PHASE 1**

Testing of the Anti Cancer Drugs CB-839 HCl (Telaglenastat) and MLN0128 (Sapanisertib) in Advanced Stage Non-small Cell Lung Cancer

**TARGETS**  
mTORC1, mTORC2, GLS

**LOCATIONS:** New York, California

**NCT03065062**
**PHASE 1**

Study of the CDK4/6 Inhibitor Palbociclib (PD-0332991) in Combination With the PI3K/mTOR Inhibitor Gedatolisib (PF-05212384) for Patients With Advanced Squamous Cell Lung, Pancreatic, Head &amp; Neck and Other Solid Tumors

**TARGETS**  
PI3K-alpha, PI3K-gamma, mTORC1, mTORC2, CDK4, CDK6

**LOCATIONS:** Massachusetts



ORDERED TEST # ORD-1002521-02

**CLINICAL TRIALS**
**NCT03154294**
**PHASE 1**

Evaluation of the Safety and Tolerability of TAK-228 With TAK-117 and Paclitaxel in Advanced Solid Tumors

**TARGETS**  
PI3K-alpha, mTORC1, mTORC2

**LOCATIONS:** South Dakota

**NCT02664935**
**PHASE 2**

National Lung Matrix Trial: Multi-drug Phase II Trial in Non-Small Cell Lung Cancer

**TARGETS**  
FGFRs, mTORC1, mTORC2, CDK4, CDK6, ALK, AXL, MET, ROS1, TRKA, TRKC, MEK, AKTs, EGFR, PD-L1, DDR2, FLT3, KIT, PDGFRA, RET, TRKB, VEGFRs

**LOCATIONS:** Exeter (United Kingdom), Belfast (United Kingdom), Cardiff (United Kingdom), Bristol (United Kingdom), Wirral (United Kingdom), Southampton (United Kingdom), Glasgow (United Kingdom), Birmingham (United Kingdom), Manchester (United Kingdom), Oxford (United Kingdom)

**NCT04337463**
**PHASE NULL**

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

**TARGETS**  
mTORC1, mTORC2, PD-1

**LOCATIONS:** Chengdu (China)

ORDERED TEST # ORD-1002521-02

**APPENDIX**
**Variants of Unknown Significance**

**NOTE** One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

**ALK**  
M338I

**ARFRP1**  
A114V

**C11ORF30 (EMSY)**  
E917A

**ERF1**  
amplification

**FLT1**  
S356C

**H3F3A**  
amplification

**MAP3K1**  
S939C

**MLL2**  
P2717S

**SDHA**  
amplification

**TNFRSF14**  
amplification

ORDERED TEST # ORD-1002521-02

**APPENDIX**
**Genes Assayed in FoundationOne®CDx**

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

**DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS**

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXJ2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMS5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

**DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS**

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPR352

\*TERC is an NCRNA

\*\*Promoter region of TERT is interrogated

**ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS**

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

ORDERED TEST # ORD-1002521-02

**APPENDIX**

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


**ABOUT FOUNDATIONONE CDx**

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:  
[www.rochefoundationmedicine.com/f1cdxtech](http://www.rochefoundationmedicine.com/f1cdxtech).

**INTENDED USE**

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

**TEST PRINCIPLES**

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes. Using an Illumina® HiSeq platform, hybrid

capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

**THE REPORT**

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

**Diagnostic Significance**

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

**Qualified Alteration Calls (Equivocal and Subclonal)**

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

**Ranking of Alterations and Therapies Biomarker and Genomic Findings**

Therapies are ranked based on the following criteria: Therapies with clinical benefit in patient's tumor type (ranked alphabetically within each NCCN category) followed by therapies with clinical benefit in other tumor type (ranked alphabetically within each NCCN category).

**Clinical Trials**

Pediatric trial qualification → Geographical proximity → Later trial phase.

**NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION**

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) ([www.nccn.org](http://www.nccn.org)). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2020. All rights reserved. To view the most recent and complete version of the guideline, go online to [NCCN.org](http://NCCN.org). NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

**Limitations**

1. The MSI-H/MSS designation by FMI F1CDx test is based on genome wide analysis of 95 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines. The threshold for MSI-H/MSS was determined by analytical concordance to comparator assays (IHC and PCR) using uterine, cecum and colorectal cancer FFPE tissue. The clinical validity of the qualitative MSI designation has not been established. For Microsatellite Instability (MSI) results, confirmatory testing using a validated orthogonal method should be considered.
2. TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Donna Ferguson, M.D. | 01 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1002521-02

APPENDIX

About FoundationOne®CDx

of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/P170019B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf). The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.

3. The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

### VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

#### Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31

INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

\*Interquartile Range = 1<sup>st</sup> Quartile to 3<sup>rd</sup> Quartile

### VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters with no conflicts), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

### LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

### NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

### NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

### TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or

none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

### SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 2.2.0

ORDERED TEST # ORD-1002521-02

APPENDIX

About FoundationOne®CDx

The median exon coverage for this sample is 742x

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Donna Ferguson, M.D. | 01 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # ORD-1002521-02

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Warth A, et al. Virchows Arch. (2016) PMID: 26637197
7. Ninomiya H, et al. Br. J. Cancer (2006) PMID: 16641899
8. Vanderwalde A, et al. Cancer Med (2018) PMID: 29436178
9. Zang YS, et al. Cancer Med (2019) PMID: 31270941
10. Dudley JC, et al. Clin. Cancer Res. (2016) PMID: 26880610
11. Takamochi K, et al. Lung Cancer (2017) PMID: 28676214
12. Pylkkänen L, et al. Environ. Mol. Mutagen. (1997) PMID: 9329646
13. Gonzalez R, et al. Ann. Oncol. (2000) PMID: 11061602
14. Chen XQ, et al. Nat. Med. (1996) PMID: 8782463
15. Merlo A, et al. Cancer Res. (1994) PMID: 8174113
16. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
17. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
18. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
19. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
20. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
21. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
22. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
23. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
24. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
25. Cristescu R, et al. Science (2018) PMID: 30309915
26. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
27. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
28. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
29. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
30. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
31. Rizvi NA, et al. Science (2015) PMID: 25765070
32. Colli LM, et al. Cancer Res. (2016) PMID: 27197178
33. Wang VE, et al. J Immunother Cancer (2017) PMID: 28923100
34. Carbone DP, et al. N. Engl. J. Med. (2017) PMID: 28636851
35. Rizvi H, et al. J. Clin. Oncol. (2018) PMID: 29337640
36. Forde PM, et al. N. Engl. J. Med. (2018) PMID: 29658848
37. Miao D, et al. Nat. Genet. (2018) PMID: 30150660
38. Chae YK, et al. Clin Lung Cancer (2019) PMID: 30425022
39. Paz-Ares et al., 2019; ESMO Abstract LBA80
40. Hellmann MD, et al. N. Engl. J. Med. (2019) PMID: 31562796
41. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
42. Spigel et al., 2016; ASCO Abstract 9017
43. Xiao D, et al. Oncotarget (2016) PMID: 27009843
44. Shim HS, et al. J Thorac Oncol (2015) PMID: 26200269
45. Govindan R, et al. Cell (2012) PMID: 22980976
46. Ding L, et al. Nature (2008) PMID: 18948947
47. Imielinski M, et al. Cell (2012) PMID: 22980975
48. Kim Y, et al. J. Clin. Oncol. (2014) PMID: 24323028
49. Stein et al., 2019; DOI: 10.1200/PO.18.00376
50. Chen Y, et al. J. Exp. Clin. Cancer Res. (2019) PMID: 31088500
51. Yu H, et al. J Thorac Oncol (2019) PMID: 30253973
52. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
53. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
54. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
55. Johnson BE, et al. Science (2014) PMID: 24336570
56. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
57. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
58. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
59. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
60. Nature (2012) PMID: 22810696
61. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
62. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
63. Pirker R, et al. Lancet Oncol. (2012) PMID: 22056021
64. Jiang Z, et al. PLoS ONE (2013) PMID: 23441167
65. Licitra L, et al. Ann. Oncol. (2011) PMID: 21048039
66. Herbst RS, et al. Lancet Oncol. (2018) PMID: 29169877
67. Paz-Ares L, et al. Ann. Oncol. (2016) PMID: 27207107
68. Zhang X, et al. J. Invest. Med. (2017) PMID: 27664271
69. Dahabreh IJ, et al. Ann. Oncol. (2011) PMID: 20826716
70. Cappuzzo F, et al. J. Natl. Cancer Inst. (2005) PMID: 15870435
71. Cappuzzo F, et al. J Thorac Oncol (2015) PMID: 25514804
72. Kwak EL, et al. Cancer (2013) PMID: 23775486
73. Thatcher N, et al. Lancet Oncol. (2015) PMID: 26045340
74. Paz-Ares L, et al. Lancet Oncol. (2015) PMID: 25701171
75. Elez E, et al. Br. J. Cancer (2016) PMID: 26766738
76. Kuenen B, et al. Clin. Cancer Res. (2010) PMID: 20197484
77. Shimamura T, et al. Cancer Res. (2005) PMID: 16024644
78. Shimamura T, et al. Cancer Res. (2008) PMID: 18632637
79. Sawai A, et al. Cancer Res. (2008) PMID: 18199556
80. Bernardes CE, et al. J Phys Condens Matter (2015) PMID: 25923649
81. Xu W, et al. Br. J. Cancer (2007) PMID: 17712310
82. Zeng Q, et al. J. Med. Chem. (2015) PMID: 26313252
83. Yang Z, et al. Sci Transl Med (2016) PMID: 27928026
84. Ahn et al., 2019; ASCO 31587882
85. Strong JE, et al. EMBO J. (1998) PMID: 9628872
86. Coffey MC, et al. Science (1998) PMID: 9812900
87. Gong J, et al. Front Oncol (2014) PMID: 25019061
88. Forsyth P, et al. Mol. Ther. (2008) PMID: 18253152
89. Vidal L, et al. Clin. Cancer Res. (2008) PMID: 18981012
90. Gollamudi R, et al. Invest New Drugs (2010) PMID: 19572105
91. Harrington KJ, et al. Clin. Cancer Res. (2010) PMID: 20484020
92. Comins C, et al. Clin. Cancer Res. (2010) PMID: 20926400
93. Lolkema MP, et al. Clin. Cancer Res. (2011) PMID: 21106728
94. Galanis E, et al. Mol. Ther. (2012) PMID: 22871663
95. Karapanagiotou EM, et al. Clin. Cancer Res. (2012) PMID: 22316603
96. Morris DG, et al. Invest New Drugs (2013) PMID: 22886613
97. Villalona-Calero MA, et al. Cancer (2016) PMID: 26709987
98. Li A, et al. Clin. Cancer Res. (2017) PMID: 28396313
99. Ou SI, et al. Lung Cancer (2016) PMID: 27393507
100. Engelman JA, et al. Science (2007) PMID: 17463250
101. Planchard D, et al. Ann. Oncol. (2015) PMID: 26269204
102. Benedettini E, et al. Am. J. Pathol. (2010) PMID: 20489150
103. Scheffler M, et al. J Thorac Oncol (2015) PMID: 26001148
104. Bahcall M, et al. Cancer Discov (2016) PMID: 27694386
105. Wu YL, et al. J. Clin. Oncol. (2018) PMID: 30156984
106. York ER, et al. J Thorac Oncol (2017) PMID: 28274743
107. Nature (2014) PMID: 25079552
108. Nature (2012) PMID: 22960745
109. Park S, et al. Histol. Histopathol. (2012) PMID: 22207554
110. Liang Z, et al. BMC Cancer (2010) PMID: 20637128
111. Grob TJ, et al. Lung Cancer (2013) PMID: 23238037
112. Watzka SB, et al. Eur J Cardiothorac Surg (2010) PMID: 20353893
113. Dobashi Y, et al. Hum. Pathol. (2011) PMID: 21040950
114. Ludovini V, et al. Cancer Chemother. Pharmacol. (2013) PMID: 23314677
115. Skrzypski M, et al. Clin Lung Cancer (2013) PMID: 23870818
116. Kim SH, et al. Histol. Histopathol. (2012) PMID: 22419022
117. Lee JS, et al. Ann. Surg. Oncol. (2013) PMID: 23525704
118. Oakley GJ, et al. J Thorac Oncol (2011) PMID: 21587084
119. Traynor AM, et al. Lung Cancer (2013) PMID: 23628526
120. Marks JL, et al. J Thorac Oncol (2008) PMID: 18303429
121. Izar B, et al. Ann. Thorac. Surg. (2013) PMID: 23932319
122. Ciardiello F, et al. N. Engl. J. Med. (2008) PMID: 18337605
123. Bhargava R, et al. Mod. Pathol. (2005) PMID: 15920544
124. Yang YL, et al. Chin. Med. J. (2012) PMID: 22490401
125. Yang et al., 2020; AACR Abstract CT127
126. Ou SH, et al. J Thorac Oncol (2011) PMID: 21623265
127. Schwab R, et al. Lung Cancer (2014) PMID: 24192513
128. Le X, et al. Clin Lung Cancer (2015) PMID: 25922291
129. Schrock AB, et al. J Thorac Oncol (2017) PMID: 28315738
130. Lennerz JK, et al. J. Clin. Oncol. (2011) PMID: 22042947
131. Chi AS, et al. J. Clin. Oncol. (2012) PMID: 22162573
132. Palma NA, et al. Case Rep Oncol (2014) PMID: 25232318
133. Schuler et al., 2016; ASCO Abstract 9067
134. Wu et al., 2018; WCLC Abstract P1.01-97
135. Gainor JF, et al. J Thorac Oncol (2020) PMID: 31864558
136. Gautschi O, et al. J Thorac Oncol (2020) PMID: 31864554
137. Decaens et al., 2019; ESMO Abstract 698P
138. Yang et al., 2019; AACR Abstract CT193
139. Park et al., 2019; ESMO Abstract 4770
140. Wu et al., 2019; IASLC Abstract MA09.09
141. Gan HK, et al. Clin. Cancer Res. (2019) PMID: 30952639
142. Lee J, et al. Cancer Discov (2019) PMID: 31315834
143. Kim ST, et al. Transl Oncol (2019) PMID: 30695737
144. Kwak et al., 2015; ASCO GI Abstract 01
145. Spigel DR, et al. J. Clin. Oncol. (2013) PMID: 24101053
146. Catenacci DV, et al. Cancer Discov (2011) PMID: 22389872
147. Harding JJ, et al. Clin. Cancer Res. (2019) PMID: 31142504
148. Strickler JH, et al. J. Clin. Oncol. (2018) PMID: 30285518
149. Dziadziuszko R, et al. J Thorac Oncol (2012) PMID: 22237262
150. Murray S, et al. J. Exp. Clin. Cancer Res. (2012) PMID: 22992338
151. Xia N, et al. Exp. Lung Res. (2013) PMID: 23919423
152. Nakamura Y, et al. Cancer Sci. (2007) PMID: 17459054
153. Cappuzzo F, et al. J. Clin. Oncol. (2009) PMID: 19255323
154. Zucali PA, et al. Ann. Oncol. (2008) PMID: 18467317
155. An SJ, et al. PLoS ONE (2012) PMID: 22768234
156. Le et al., 2020; AACR Abstract 3385
157. Yang JJ, et al. Lung Cancer (2013) PMID: 23079155
158. Chen YT, et al. J Thorac Oncol (2011) PMID: 22052229

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Donna Ferguson, M.D. | 01 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1002521-02

## APPENDIX

## References

159. Kanteti R, et al. J. Environ. Pathol. Toxicol. Oncol. (2009) PMID: 19817696
160. To C, et al. Exp. Cell Res. (2002) PMID: 11795945
161. Tsuta K, et al. J Thorac Oncol (2012) PMID: 22198430
162. Tanaka A, et al. Lung Cancer (2012) PMID: 21733594
163. Tong JH, et al. Clin. Cancer Res. (2016) PMID: 26847053
164. Lee GD, et al. J Thorac Oncol (2017) PMID: 28502721
165. Gow CH, et al. Lung Cancer (2017) PMID: 28024701
166. Awad et al., 2017; ASCO Abstract 8511
167. J. Clin. Oncol. (2011) PMID: 22042966
168. Jung KH, et al. Arch. Pharm. Res. (2012) PMID: 22553051
169. Gao J, et al. Sci Signal (2013) PMID: 23550210
170. Ang CS, et al. Anticancer Res. (2013) PMID: 23898085
171. Abou-Bakr AA, et al. GJF J Oncolog (2013) PMID: 23996864
172. Ho JC, et al. Semin Respir Crit Care Med (2013) PMID: 24258573
173. Madoz-Gürpide J, et al. J Transl Med (2015) PMID: 26319934
174. Ali SM, et al. Oncologist (2015) PMID: 25882375
175. Kwak EL, et al. Cancer Discov (2015) PMID: 26432108
176. Zhang Y, et al. Clin. Cancer Res. (2013) PMID: 23386689
177. Xie Q, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22203985
178. Kentsis A, et al. Nat. Med. (2012) PMID: 22683780
179. Xie Q, et al. Genes Cancer (2013) PMID: 24167653
180. Wanjala J, et al. Mol. Cancer Ther. (2015) PMID: 25381262
181. Cloughesy et al., 2015; ASCO Abstract 2015
182. Koepfen H, et al. Clin. Cancer Res. (2014) PMID: 24687921
183. Patnaik A, et al. Br. J. Cancer (2014) PMID: 24901237
184. Iveson T, et al. Lancet Oncol. (2014) PMID: 24965569
185. Yano S, et al. Cancer Res. (2008) PMID: 19010923
186. Turke AB, et al. Cancer Cell (2010) PMID: 20129249
187. Nakade J, et al. J Thorac Oncol (2014) PMID: 24828661
188. Nanjo S, et al. PLoS ONE (2013) PMID: 24386407
189. Chen X, et al. Anticancer Drugs (2013) PMID: 23962905
190. Straussman R, et al. Nature (2012) PMID: 22763439
191. Onitsuka T, et al. J Thorac Oncol (2010) PMID: 20150826
192. Han JY, et al. Lung Cancer (2011) PMID: 21440951
193. Kasahara K, et al. Clin. Cancer Res. (2010) PMID: 20679350
194. Stamos J, et al. EMBO J. (2004) PMID: 15167892
195. Tolbert WD, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20624990
196. Zack TI, et al. Nat. Genet. (2013) PMID: 24071852
197. Beroukhi R, et al. Nature (2010) PMID: 20164920
198. Cheng H, et al. Cancer Discov (2015) PMID: 26370156
199. Wicki A, et al. Eur. J. Cancer (2018) PMID: 29660598
200. Park et al., 2019; ASCO abstract 8514
201. Ruder et al., 2015; AACR Abstract 3576
202. Sarbassov DD, et al. Curr. Biol. (2004) PMID: 15268862
203. Jacinto E, et al. Nat. Cell Biol. (2004) PMID: 15467718
204. Pearce LR, et al. Biochem. J. (2007) PMID: 17461779
205. Ross JS, et al. J. Clin. Pathol. (2014) PMID: 24978188
206. Cheng et al., 2014; ASCO Abstract 8027
207. Kristeleit et al., 2015; ASCO Abstract 2592
208. Nomura S, et al. Br. J. Cancer (2008) PMID: 18594526
209. Memarzadeh S, et al. Cancer Cell (2007) PMID: 18068633
210. Javle M, et al. Hum. Pathol. (2014) PMID: 24508317
211. Ooi A, et al. Mod. Pathol. (2015) PMID: 25743022
212. Chattopadhyay I, et al. Mutat. Res. (2010) PMID: 20083228
213. Cha JD, et al. Oral Surg Oral Med Oral Pathol Oral Radiol Endod (2011) PMID: 21334929
214. Theodorou V, et al. Oncogene (2004) PMID: 15208658
215. Nakao Y, et al. Int. J. Oncol. (2013) PMID: 24002438
216. Zhang X, et al. J. Biol. Chem. (2006) PMID: 16597617
217. Abolhassani A, et al. J. Cancer (2014) PMID: 25057305
218. Volckaert T, et al. Fibrogenesis Tissue Repair (2014) PMID: 24891877
219. Entesarian M, et al. Nat. Genet. (2005) PMID: 15654336
220. Mirza MR, et al. N. Engl. J. Med. (2016) PMID: 27717299
221. Sandhu SK, et al. Lancet Oncol. (2013) PMID: 23810788
222. Fong PC, et al. N. Engl. J. Med. (2009) PMID: 19553641
223. Audeh MW, et al. Lancet (2010) PMID: 20609468
224. Fong PC, et al. J. Clin. Oncol. (2010) PMID: 20406929
225. Gelmon KA, et al. Lancet Oncol. (2011) PMID: 21862407
226. Kaye SB, et al. J. Clin. Oncol. (2012) PMID: 22203755
227. Swisher EM, et al. Lancet Oncol. (2017) PMID: 27908594
228. Drew Y, et al. Br. J. Cancer (2016) PMID: 27002934
229. Farago et al., 2017; AACR Abstract CT048
230. Pietanza MC, et al. Clin. Cancer Res. (2012) PMID: 22228633
231. Rank L, et al. Nucleic Acids Res. (2016) PMID: 27694308
232. Pettitt SJ, et al. Nat Commun (2018) PMID: 29748565
233. Pettitt SJ, et al. PLoS ONE (2013) PMID: 23634208
234. Zehir A, et al. Nat. Med. (2017) PMID: 28481359
235. Murnyak B, et al. Oncotarget (2017) PMID: 28654422
236. Byers LA, et al. Cancer Discov (2012) PMID: 22961666
237. Amé JC, et al. Bioessays (2004) PMID: 15273990
238. Ray Chaudhuri A, et al. Nat. Rev. Mol. Cell Biol. (2017) PMID: 28676700
239. Qin Q, et al. PLoS ONE (2014) PMID: 24853559
240. Hua RX, et al. PLoS ONE (2014) PMID: 24489833
241. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
242. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
243. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
244. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
245. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
246. Xu L, et al. Mol. Med. (2001) PMID: 11713371
247. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
248. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
249. Pirollo KF, et al. Mol. Ther. (2016) PMID: 27357628
250. Hajdenberg et al., 2012; ASCO Abstract e15010
251. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
252. Moore et al., 2019; ASCO Abstract 5513
253. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
254. Oza et al., 2015; ASCO Abstract 5506
255. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
256. Ma CX, et al. J. Clin. Invest. (2012) PMID: 22446188
257. Kwok M, et al. Blood (2016) PMID: 26563132
258. Boudny M, et al. Haematologica (2019) PMID: 30975914
259. Dillon MT, et al. Mol. Cancer Ther. (2017) PMID: 28062704
260. Middleton FK, et al. Cancers (Basel) (2018) PMID: 30127241
261. Mogi A, et al. J. Biomed. Biotechnol. (2011) PMID: 21331359
262. Tekpli X, et al. Int. J. Cancer (2013) PMID: 23011884
263. Vignot S, et al. J. Clin. Oncol. (2013) PMID: 23630207
264. Maeng CH, et al. Anticancer Res. (2013) PMID: 24222160
265. Cortot AB, et al. Clin Lung Cancer (2014) PMID: 24169260
266. Itakura M, et al. Br. J. Cancer (2013) PMID: 23922113
267. Dong ZY, et al. Clin. Cancer Res. (2017) PMID: 28039262
268. Seo JS, et al. Genome Res. (2012) PMID: 22975805
269. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
270. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
271. Xie M, et al. Nat. Med. (2014) PMID: 25326804
272. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
273. Severson EA, et al. Blood (2018) PMID: 29678827
274. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
275. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
276. Chabon JJ, et al. Nature (2020) PMID: 32269342
277. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
278. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
279. Joergers AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
280. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
281. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
282. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
283. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
284. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
285. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
286. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100
287. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776
288. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316
289. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208
290. Lalloo F, et al. Lancet (2003) PMID: 12672316
291. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713
292. De Grève J, et al. Lung Cancer (2015) PMID: 25682316
293. Januszewski et al., 2018; IASLC WCLC Abstract P1.13-17
294. Chang et al., 2018; IASLC WCLC Abstract P1.01-11
295. Sequist LV, et al. J. Clin. Oncol. (2013) PMID: 23816960
296. Wu YL, et al. Lancet Oncol. (2014) PMID: 24439929
297. Yang JC, et al. Lancet Oncol. (2015) PMID: 25589191
298. Soria JC, et al. Lancet Oncol. (2015) PMID: 26156651
299. Paz-Ares L, et al. Ann. Oncol. (2017) PMID: 28426106
300. Dziadziuszko R, et al. J Thorac Oncol (2019) PMID: 30825613
301. Lai WV, et al. Eur. J. Cancer (2019) PMID: 30685684
302. Greulich H, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22908275
303. Gow CH, et al. J Thorac Oncol (2015) PMID: 26134234
304. Mazières J, et al. Ann. Oncol. (2016) PMID: 26598547
305. Mazières J, et al. J. Clin. Oncol. (2013) PMID: 23610105
306. De Grève J, et al. Lung Cancer (2012) PMID: 22325357
307. Li BT, et al. Lung Cancer (2015) PMID: 26559459
308. Costa DB, et al. J Thorac Oncol (2016) PMID: 26964772
309. Schuler M, et al. Ann. Oncol. (2016) PMID: 26646759
310. Wu et al., 2018; WCLC Abstract P1.01-97
311. Wolf et al., 2020; ASCO Abstract 9509
312. Schuler M, et al. Ann. Oncol. (2020) PMID: 32240796
313. Tan et al., 2017; IASLC Abstract P3.02b-117
314. van den Bent M, et al. J. Neurooncol. (2020) PMID: 31776899
315. Bang YJ, et al. Cancer Sci. (2020) PMID: 31778267
316. Esaki T, et al. Cancer Sci. (2019) PMID: 30724423
317. Vassal et al., 2015; ASCO Abstract 2595
318. Li et al., 2015; ASCO Abstract 8090
319. Frampton GM, et al. Cancer Discov (2015) PMID: 25971938
320. Paik PK, et al. Cancer Discov (2015) PMID: 25971939
321. Benderra MA, et al. J Thorac Oncol (2016) PMID:

© 2021 Foundation Medicine, Inc. All rights reserved.

ORDERED TEST # **ORD-1002521-02**

- 26845121
322. Waqar SN, et al. J Thorac Oncol (2015) PMID: 25898962
323. Mendenhall MA, et al. J Thorac Oncol (2015) PMID: 25898965
324. Jenkins RW, et al. Clin Lung Cancer (2015) PMID: 25769807
325. Stein MN, et al. Eur. Urol. (2015) PMID: 25457019
326. Shaw et al., 2016; ASCO Abstract 9066
327. Lu et al., 2016; ASCO Abstract 9058
328. Yoshida T, et al. J. Clin. Oncol. (2016) PMID: 27354483
329. Solomon BJ, et al. N. Engl. J. Med. (2014) PMID: 25470694
330. Shaw AT, et al. N. Engl. J. Med. (2013) PMID: 23724913
331. Moro-Sibilot D, et al. Ann. Oncol. (2019) PMID: 31584608
332. Goto et al., 2016; ASCO Abstract 9022
333. Shaw AT, et al. N. Engl. J. Med. (2014) PMID: 25264305
334. Mazières J, et al. J. Clin. Oncol. (2015) PMID: 25667280
335. Scheffler M, et al. Oncotarget (2015) PMID: 25868855
336. Vaishnavi A, et al. Nat. Med. (2013) PMID: 24162815
337. Drilon et al., 2016; ASCO Abstract 108
338. Camidge et al., 2014; ASCO Abstract 8001
339. Schrock AB, et al. J Thorac Oncol (2016) PMID: 27343443
340. Jorge SE, et al. Lung Cancer (2015) PMID: 26791794
341. Mahjoubi L, et al. Invest New Drugs (2016) PMID: 26892698
342. Awad MM, et al. J. Clin. Oncol. (2016) PMID: 26729443
343. Zhang Y, et al. J Thorac Oncol (2016) PMID: 26724472
344. Landi L, et al. Clin. Cancer Res. (2019) PMID: 31416808
345. Caparica R, et al. J Thorac Oncol (2017) PMID: 27664533
346. Dagogo-Jack I, et al. Clin Cancer Res (2020) PMID: 32086345
347. Wu YL, et al. Lancet Oncol. (2017) PMID: 28958502
348. Mok TS, et al. J. Clin. Oncol. (2018) PMID: 29864379
349. Necchi A, et al. BJU Int. (2018) PMID: 28921872
350. Zhu Y, et al. Cancer Biol. Ther. (2014) PMID: 24658109
351. Hokenfu Zasshi (1979) PMID: 259761
352. Opsomer RJ, et al. Acta Urol Belg (1985) PMID: 2986437
353. Wu et al., 2018; WCLC abstract MA26.11
354. Ramalingam SS, et al. Ann. Oncol. (2016) PMID: 26768165
355. Yu HA, et al. Lung Cancer (2017) PMID: 29191595
356. Reckamp KL, et al. Cancer (2014) PMID: 24501009
357. Jänne PA, et al. Clin. Cancer Res. (2011) PMID: 21220471
358. van Geel RMJM, et al. Br. J. Cancer (2020) PMID: 32147669
359. Jänne PA, et al. J Thorac Oncol (2016) PMID: 26899759
360. Dahabreh JJ, et al. Clin. Cancer Res. (2010) PMID: 20028749
361. Shepherd FA, et al. N. Engl. J. Med. (2005) PMID: 16014882
362. Han JY, et al. J. Clin. Oncol. (2012) PMID: 22370314
363. Maemondo M, et al. N. Engl. J. Med. (2010) PMID: 20573926
364. Mitsudomi T, et al. Lancet Oncol. (2010) PMID: 20022809
365. Mok TS, et al. N. Engl. J. Med. (2009) PMID: 19692680
366. Petrelli F, et al. Clin Lung Cancer (2012) PMID: 22056888
367. Qi WX, et al. Curr Med Res Opin (2015) PMID: 25329826
368. Zhao H, et al. J Thorac Oncol (2015) PMID: 25546556
369. Petty RD, et al. J. Clin. Oncol. (2017) PMID: 28537764
370. Dutton SJ, et al. Lancet Oncol. (2014) PMID: 24950987
371. Cheng et al., 2018; ESMO Abstract 13770
372. Yang et al., 2019; AACR Abstract CT193/7
373. Douillard JY, et al. Br. J. Cancer (2014) PMID: 24263064
374. Fukuoka M, et al. J. Clin. Oncol. (2011) PMID: 21670455
375. Noronha V, et al. J. Clin. Oncol. (2019) PMID: 31411950
376. Hosomi Y, et al. J. Clin. Oncol. (2020) PMID: 31682542
377. Sutiman N, et al. J Thorac Oncol (2017) PMID: 27908825
378. Gibbons DL, et al. J Thorac Oncol (2016) PMID: 27198414
379. Klempner SJ, et al. J Thorac Oncol (2017) PMID: 27693535
380. Yakes FM, et al. Mol. Cancer Ther. (2011) PMID: 21926191
381. Weber H, et al. J Biomol Screen (2014) PMID: 25260782
382. Navis AC, et al. PLoS ONE (2013) PMID: 23484006
383. Yeh I, et al. Nat Commun (2015) PMID: 26013381
384. Lee YH, et al. Cancers (Basel) (2014) PMID: 25534569
385. Torres KE, et al. Clin. Cancer Res. (2011) PMID: 21540237
386. Sameni M, et al. Clin. Cancer Res. (2016) PMID: 26432786
387. Hellerstedt BA, et al. Clin Lung Cancer (2019) PMID: 30528315
388. Neal JW, et al. Lancet Oncol. (2016) PMID: 27825638
389. Nokihara H, et al. Clin Lung Cancer (2019) PMID: 30718102
390. Ramalingam SS, et al. Lung Cancer (2013) PMID: 23849982
391. Kim ES, et al. Lancet Oncol. (2013) PMID: 24231627
392. Janjigian YY, et al. Cancer Discov (2014) PMID: 25074459
393. Wheler JJ, et al. Mol. Cancer Ther. (2013) PMID: 23963360
394. Tsigelny IF, et al. Oncotarget (2015) PMID: 25760241
395. Crawford J, et al. J Thorac Oncol (2013) PMID: 24389433
396. Schuette W, et al. Clin Lung Cancer (2015) PMID: 26094080
397. Castellanos EH, et al. Clin Lung Cancer (2015) PMID: 25842367