

**ABOUT THE TEST** FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

## PATIENT

**DISEASE** Breast carcinoma (NOS)

**DATE OF BIRTH** 20 March 1974

**SEX** Female

**MEDICAL RECORD #** Not given

## PHYSICIAN

**MEDICAL FACILITY** Hospital Regional Lambayeque

**ADDITIONAL RECIPIENT** None

**MEDICAL FACILITY ID** 319645

**PATHOLOGIST** Not Provided

## SPECIMEN

**SPECIMEN SITE** Breast

**SPECIMEN ID** H21-3840

**SPECIMEN TYPE** Block

**DATE OF COLLECTION** 03 April 2019

**SPECIMEN RECEIVED** 16 February 2021

## Biomarker Findings

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 6 Muts/Mb

## Genomic Findings

*For a complete list of the genes assayed, please refer to the Appendix.*

**FGFR2** amplification

**PTEN** deletion exon 8

**TP53** splice site 375+1G>A

**4 Disease relevant genes with no reportable alterations: *BRCA1, BRCA2, ERBB2, PIK3CA***

**4 Therapies with Clinical Benefit**

**20 Clinical Trials**

**0 Therapies with Lack of Response**

## BIOMARKER FINDINGS

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 6 Muts/Mb

## GENOMIC FINDINGS

**PTEN** - deletion exon 8

**10 Trials** *see p. 10*

**FGFR2** - amplification

**10 Trials** *see p. 8*

## ACTIONABILITY

**No therapies or clinical trials.** *see Biomarker Findings section*

**No therapies or clinical trials.** *see Biomarker Findings section*

### THERAPIES WITH CLINICAL BENEFIT (IN PATIENT'S TUMOR TYPE)

none

none

### THERAPIES WITH CLINICAL BENEFIT (IN OTHER TUMOR TYPE)

Everolimus 2A

Temsirolimus

Erdaftinib

Pazopanib

☐ NCCN category

## GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIALS OPTIONS

*For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.*

**TP53** - splice site 375+1G>A ..... **p. 5**

**NOTE** Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

ORDERED TEST # ORD-1021124-01

## BIOMARKER FINDINGS

## BIOMARKER

## Microsatellite status

## RESULT

MS-Stable

### POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors<sup>1-3</sup>, including approved therapies nivolumab and pembrolizumab<sup>4</sup>. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared

with non-MSI-H cases (70% vs. 12%,  $p=0.001$ )<sup>5</sup>.

### FREQUENCY & PROGNOSIS

MSI is extremely rare in breast cancer, reported in 0-1% of cases across studies<sup>6-11</sup>. The incidence of MSI is increased in triple-negative breast cancer<sup>9-11</sup> and in tumors with homologous recombination defects, such as mutations in BRCA1/2<sup>9,11</sup>. Notably, in Lynch syndrome-related breast cancer, MSI has been reported in 51-85% of cases<sup>12-17</sup>. A prospective study of 123 patients with breast cancer treated with chemotherapy reported an increase in the incidence of MSI-H following chemotherapy treatment (from 0% pre-treatment to 19% post-treatment) and a significant association between MSI and tumor recurrence<sup>18</sup>.

### FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor<sup>19</sup>. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2<sup>19-21</sup>. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers<sup>22-24</sup>. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins<sup>19,21,23-24</sup>.

## BIOMARKER

## Tumor Mutational Burden

## RESULT

6 Muts/Mb

### POTENTIAL TREATMENT STRATEGIES

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1<sup>25-27</sup>, anti-PD-1 therapies<sup>25-28</sup>, and combination nivolumab and ipilimumab<sup>29-33</sup>. In multiple pan-tumor studies, higher TMB has been reported to be associated with increased ORR and OS from treatment with immune checkpoint inhibitors<sup>25-28,34</sup>. Higher TMB was found to be significantly associated with improved OS upon immune checkpoint inhibitor treatment of patients with 9 types of advanced tumors<sup>25</sup>. Analyses across several solid tumor types reported that patients with higher TMB (defined as  $\geq 16-20$  Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy, compared with patients with higher TMB treated with chemotherapy<sup>35</sup> or those with lower TMB treated with PD-1 or PD-L1-targeting agents<sup>26</sup>. However, the KEYNOTE 158 trial of pembrolizumab monotherapy in patients with solid tumors found significant improvement in ORR in patients with TMB  $\geq 10$  Muts/Mb (based on this assay or others)

compared to those with TMB  $< 10$  Muts/Mb, in a large cohort that included multiple tumor types; similar findings were observed in the KEYNOTE 028 and 012 trials<sup>28,34</sup>. Together, these studies suggest that patients with TMB  $\geq 10$  Muts/Mb may derive clinical benefit from PD-1 or PD-L1 inhibitors.

### FREQUENCY & PROGNOSIS

Breast carcinoma harbors a median TMB of 3.8 muts/Mb, and 3.1% of cases have high TMB ( $> 20$  muts/Mb)<sup>36</sup>. A study of 3,969 patients with breast cancer reported a median TMB of 2.63 mutations per megabase (Muts/Mb), with 5% of cases harboring TMB  $\geq 10$  Muts/Mb; median TMB was significantly higher in hormone receptor (HR)-negative and HER2-negative tumors than HR-positive or HER2-positive tumors<sup>37</sup>. The Breast Invasive Carcinoma TCGA analysis reported an average (non-silent) mutation load of 0.84 Muts/Mb for luminal A tumors, 1.38 Muts/Mb for luminal B tumors, 2.05 Muts/Mb for HER2-enriched tumors, and 1.68 Muts/Mb for basal-like tumors<sup>38</sup>. In breast cancer, TMB is significantly higher in recurrent versus primary tumors, metastatic versus localized cancers, triple-negative versus HR-positive tumors, and CDH1-mutated versus CDH1-wildtype tumors<sup>37,39-40</sup>. Among metastatic tumors, TMB-high samples have been reported more frequently in invasive lobular carcinoma (9-17% of cases, depending on the TMB cutoff to designate TMB-high) than in invasive ductal carcinoma (2-8% of cases, depending on the cutoff), and TMB-high (at

either cutoff) has not been observed in papillary carcinoma<sup>37,39-40</sup>. In a large study of patients with breast cancer, hypermutation was more frequently observed in metastatic tumors than in primary tumors<sup>37</sup>. In a study of 14,867 patients with breast cancer, high TMB was associated with older age and metastatic disease but was not significantly associated with PD-L1 positivity using the TMB cutoff of  $\geq 10$  Muts/Mb<sup>40</sup>. In estrogen receptor-positive breast cancer, increased TMB in tissue samples ( $> \text{mean of } 1.25 \text{ Muts/Mb}$ ) associated with shorter OS (HR=2.02) in an analysis of the TCGA data<sup>41</sup>.

### FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma<sup>42-43</sup> and cigarette smoke in lung cancer<sup>44-45</sup>, treatment with temozolomide-based chemotherapy in glioma<sup>46-47</sup>, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes<sup>48-52</sup>, and microsatellite instability (MSI)<sup>48,51-52</sup>. This sample harbors a TMB level associated with lower rates of clinical benefit from treatment with PD-1- or PD-L1-targeting immune checkpoint inhibitors compared with patients with tumors harboring higher TMB levels, based on several studies in multiple solid tumor types<sup>26-27,34</sup>.

ORDERED TEST # ORD-1021124-01

**GENOMIC FINDINGS**
**GENE**
**FGFR2**
**ALTERATION**

amplification

**POTENTIAL TREATMENT STRATEGIES**

FGFR2 activating mutations, amplification, or fusions may confer sensitivity to FGFR-targeting kinase inhibitors such as erdafitinib<sup>53</sup>, pemigatinib<sup>54-55</sup>, E7090<sup>56</sup>, pazopanib<sup>57-58</sup>, ponatinib<sup>59</sup>, AZD4547<sup>60-62</sup>, derazantinib<sup>61-62</sup>, Debio-1347<sup>63</sup>, infigratinib<sup>64</sup>, and futibatinib<sup>65</sup>. In the context of FGFR2 amplification, clinical benefit has been reported for patients with breast cancer treated with erdafitinib<sup>66</sup> and infigratinib<sup>64</sup>. In a Phase 2 study of the FGFR inhibitor AZD4547, responses were reported in

33% (3/9) of patients with FGFR2-amplified gastroesophageal cancer; in this study, higher-level amplification correlated with higher likelihood of response to FGFR inhibitors<sup>60</sup>. However, a randomized Phase 2 study of AZD4547 compared with paclitaxel for the treatment of patients with advanced stomach adenocarcinoma harboring FGFR2 amplification or polysomy reported no significant increase in median PFS, median OS, or ORR<sup>61</sup>.

**FREQUENCY & PROGNOSIS**

FGFR2 amplification has been reported in breast cancer at frequencies ranging from 1-12%<sup>67-70</sup>. FGFR2 protein expression has been reported in 13% of triple-negative breast cancer cases<sup>70</sup> and in ~60% of invasive ductal carcinomas<sup>71</sup>. In a study of 125 patients with invasive ductal carcinoma, high-level FGFR2 protein expression was associated with decreased overall and disease-free

survival<sup>71</sup> and a meta-analysis of over 11,000 patients with breast cancer showed a significant association between FGFR2/3 expression and reduced overall survival<sup>72</sup>. In contrast, a study of triple-negative breast cancer showed no correlation between FGFR2 amplification or overexpression with survival<sup>70</sup>.

**FINDING SUMMARY**

FGFR2 encodes a tyrosine kinase cell surface receptor, which plays an important role in cell differentiation, growth, and angiogenesis<sup>73-74</sup>. FGFR2 amplification has been reported in a variety of cancer types<sup>75</sup> and has been shown to correlate with increased mRNA and protein expression<sup>60,76</sup>. Higher level, clonal FGFR2 amplification has been reported to correlate with higher response rates to FGFR inhibitors<sup>60,77</sup>.

ORDERED TEST # ORD-1021124-01

## GENOMIC FINDINGS

## GENE

## PTEN

## ALTERATION

deletion exon 8

## POTENTIAL TREATMENT STRATEGIES

PTEN loss or mutation leads to activation of the PI3K-AKT-mTOR pathway and may predict sensitivity to inhibitors of this pathway<sup>78-82</sup> such as the mTOR inhibitors temsirolimus and everolimus or the PI3K inhibitor copanlisib. Preclinical studies suggest that PTEN-deficient cancers, in the absence of other oncogenic mutations, depend primarily on the beta isoform of PI3K (PI3K-beta)<sup>83-85</sup>, and PI3K-beta-selective inhibitors are in clinical trials for PTEN-deficient tumors. However, the NCI-MATCH Phase 2 study observed limited activity of the PI3K-beta-selective inhibitor GSK2636771 as monotherapy in PTEN-deficient cancers, with a median PFS of 1.8 months. The best outcomes were 1 PR (1/22, prostate cancer), SD (7/22) for patients with PTEN deletion/mutation, and SD (9/34) for patients with PTEN protein loss<sup>86</sup>. Clinical data in breast<sup>87-88</sup> and prostate cancer<sup>89-90</sup> suggest that PTEN alterations may predict sensitivity to pan-AKT inhibitors such as ipatasertib or capivasertib. Phase 2 studies have reported improved PFS from the addition of either ipatasertib (9.0 vs. 4.9 months, HR=0.44) or capivasertib (9.3 vs. 3.7 months, HR=0.30) to paclitaxel, compared with paclitaxel and placebo, for patients with metastatic triple-negative breast cancer harboring PIK3CA/AKT1/PTEN alterations<sup>91</sup>. However, data from the Phase 2 LOTUS trial reported an improved PFS for patients with PIK3CA/AKT1/PTEN-altered triple negative breast cancer treated with ipatasertib plus paclitaxel compared to paclitaxel

alone (9.0 vs 4.9 months)<sup>92</sup>; however, the Phase 3 IPATunity130 trial did not report a significant PFS improvement for first-line ipatasertib in combination with paclitaxel relative to paclitaxel alone (7.4 vs 6.1 months)<sup>93</sup>. Preclinical data indicate that PTEN loss or inactivation may predict sensitivity to PARP inhibitors<sup>94-98</sup>, and clinical benefit has been observed for patients with PTEN-altered breast cancer<sup>99</sup>, ovarian cancer<sup>100</sup>, endometrial cancer<sup>98</sup>, and other tumor types<sup>101</sup> treated with PARP inhibitors. However, several studies have reported a lack of association between PTEN mutation and PARP inhibitor sensitivity<sup>97,102-103</sup>. Limited clinical evidence in glioblastoma<sup>104</sup>, leiomyosarcoma<sup>105</sup>, NSCLC<sup>106</sup>, and melanoma<sup>107</sup> suggests that PTEN alterations may predict a lack of response to anti-PD-1 therapy. In an analysis of 39 patients with metastatic melanoma treated with pembrolizumab or nivolumab, patients with PTEN-expressing tumors achieved significantly greater reduction of tumor size than those with reduction or loss of PTEN expression<sup>107</sup>. In a retrospective analysis of 66 patients with glioblastoma, tumors from nivolumab or pembrolizumab non-responders were significantly enriched for PTEN mutations<sup>104</sup>. In a patient with uterine leiomyosarcoma treated with pembrolizumab monotherapy, a treatment-resistant tumor arose that harbored PTEN loss<sup>105</sup>. A patient with NSCLC whose tumor harbored a PTEN alteration exhibited a lack of response to nivolumab and pembrolizumab<sup>106</sup>. Clinical and preclinical evidence suggests that PTEN loss or mutation may predict resistance to PI3K inhibitors<sup>108-110</sup>, and to CDK inhibitors such as palbociclib, ribociclib, and abemaciclib<sup>108,111</sup>.

## FREQUENCY &amp; PROGNOSIS

In the TCGA dataset, PTEN mutation has been reported in 4% of breast invasive carcinomas,

while putative homozygous deletion of PTEN has been reported in 2% of cases<sup>38</sup>. PTEN mutation has also been observed in 5.3% (1/19) of metaplastic breast cancers<sup>112</sup> and 2% of invasive lobular carcinoma tumors analyzed<sup>113</sup>. PTEN mutations are associated more frequently with triple-negative breast cancer than with HER2- or hormone-positive breast cancer<sup>114-115</sup>. Loss or reduction of PTEN expression has been observed in 28% of invasive ductal breast carcinomas and has been correlated with metastasis and poor patient prognosis, including decreased 2-year disease-free survival<sup>116-118</sup>.

## FINDING SUMMARY

PTEN encodes an inositol phosphatase that functions as a tumor suppressor by negatively regulating the PI3K-AKT-mTOR pathway; loss of PTEN can lead to uncontrolled cell growth and suppression of apoptosis<sup>80</sup>. Alterations such as seen here may disrupt PTEN function or expression<sup>119-159</sup>.

## POTENTIAL GERMLINE IMPLICATIONS

PTEN mutations underlie several inherited disorders, collectively termed PTEN hamartoma tumor syndrome (PHTS), which include Cowden syndrome (CS) and its variant Lhermitte-Duclos disease (LD), Bannayan-Riley-Ruvalcaba syndrome (BRRS), PTEN-related Proteus syndrome (PS), and Proteus-like syndrome<sup>160-161</sup>. The mutation rate for PTEN in these disorders ranges from 20 to 85% of patients<sup>160,162</sup>. The estimated incidence of Cowden syndrome is 1/200,000, which may be an underestimate due to the high variability of this disorder<sup>160</sup>. Given the association between PTEN and these inherited syndromes, in the appropriate clinical context, germline testing for mutations affecting PTEN is recommended.

ORDERED TEST # ORD-1021124-01

## GENOMIC FINDINGS

## GENE

## TP53

## ALTERATION

splice site 375+1G&gt;A

## TRANSCRIPT ID

NM\_000546

## CODING SEQUENCE EFFECT

375+1G&gt;A

## VARIANT ALLELE FREQUENCY (% VAF)

32.0%

## POTENTIAL TREATMENT STRATEGIES

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib<sup>163-166</sup>, or p53 gene therapy and immunotherapeutics such as SGT-53<sup>167-171</sup> and ALT-801<sup>172</sup>. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% (17/176) and SDs in 53.4% (94/176) of patients with solid tumors; the response rate was 21.1% (4/19) in patients with TP53 mutations versus 12.1% (4/33) in patients who were TP53 wild-type<sup>173</sup>. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 31.9% (30/94, 3 CR) ORR and a 73.4% (69/94) DCR in patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer<sup>174</sup>. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 42.9% (9/21, 1 CR) ORR and a 76.2% (16/21) DCR in patients with platinum-refractory TP53-mutated ovarian cancer<sup>175</sup>. The combination of adavosertib with paclitaxel and carboplatin in patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and

carboplatin alone<sup>176</sup>. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24.0% (6/25) ORR with adavosertib combined with paclitaxel<sup>177</sup>. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71.4% (5/7) response rate for patients with TP53 alterations<sup>178</sup>. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel in patients with solid tumors, 75.0% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage<sup>171</sup>. Additionally, the combination of a CHK1 inhibitor and irinotecan reportedly reduced tumor growth and prolonged survival in a TP53-mutated, but not TP53-wild-type, breast cancer xenotransplant mouse model<sup>179</sup>. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies<sup>180-181</sup>; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies<sup>182-183</sup>. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

## FREQUENCY &amp; PROGNOSIS

TP53 is one of the most commonly mutated genes in breast cancer; mutations in this gene have been identified in 27-37% of breast carcinoma samples<sup>38,184-188</sup>. TP53 mutations that are located within the region encoding the DNA binding domain are associated with poor prognosis in patients with breast cancer<sup>186,189-190</sup>. TP53 mutation is also implicated in breast cancer susceptibility, as TP53 mutation carriers have an 18-60 fold increased risk for early onset breast cancer<sup>191-193</sup>.

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion<sup>194-199</sup>. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy<sup>194-195</sup>. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease<sup>200</sup>. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH<sup>198,201-202</sup>. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

## FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers<sup>203</sup>. Alterations such as seen here may disrupt TP53 function or expression<sup>204-208</sup>.

## POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers<sup>209-211</sup>, including sarcomas<sup>212-213</sup>. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000<sup>214</sup> to 1:20,000<sup>213</sup>. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30<sup>215</sup>. In the appropriate clinical context, germline testing of TP53 is recommended.



ORDERED TEST # ORD-1021124-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

## Erdaftinib

*Assay findings association*
**FGFR2**  
amplification

### AREAS OF THERAPEUTIC USE

Erdaftinib is a pan-fibroblast growth factor receptor (FGFR) inhibitor. It is FDA approved for the treatment of patients with advanced or metastatic urothelial carcinoma who have FGFR2 or FGFR3 alterations and have progressed after prior chemotherapy. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

On the basis of strong clinical evidence for FGFR2 fusions<sup>53,216-217</sup>, limited evidence for FGFR2 mutations<sup>217-218</sup> and limited evidence for FGFR2 amplification<sup>66</sup>, and preclinical data<sup>219-220</sup>, FGFR2 activating alterations may confer sensitivity to erdaftinib.

### SUPPORTING DATA

Clinical data on the efficacy of erdaftinib for the treatment of breast cancer are limited (PubMed, Feb 2021).

Erdaftinib has been primarily studied for the treatment of FGFR-altered urothelial carcinoma. A Phase 2 study evaluating erdaftinib for the treatment of patients with metastatic or unresectable urothelial carcinoma (mUC) previously treated with chemotherapy and harboring FGFR2/3 fusions or FGFR3 activating mutations reported an ORR of 40% (40/99, 3 CR), and a DCR of 80% (79/99)<sup>221</sup>. A Phase 1 trial of erdaftinib reported clinical responses in for patients with various FGFR2- or FGFR3-altered solid tumors<sup>53,218,222-223</sup>, including cholangiocarcinoma (27% ORR, 3/11), NSCLC (5% ORR, 1/21), breast (9% ORR, 3/34), and ovarian (9% ORR, 1/11), while other cancers including endometrial carcinoma and glioblastoma showed a low ORR (2%, 1/58)<sup>66</sup>. Following progression on multiple other lines of therapy, a patient with metastatic FGFR2-fusion-positive NSCLC treated with erdaftinib exhibited an 11-month PR<sup>222</sup>.

## Everolimus

*Assay findings association*
**PTEN**  
deletion exon 8

### AREAS OF THERAPEUTIC USE

Everolimus is an orally available mTOR inhibitor that is FDA approved to treat renal cell carcinoma (RCC) following antiangiogenic therapy; pancreatic neuroendocrine tumors; and well-differentiated non-functional neuroendocrine tumors of the lung or gastrointestinal tract. Everolimus is also approved to treat either renal angiomyolipoma or subependymal giant cell astrocytoma in association with tuberous sclerosis complex (TSC). Please see the drug label for full prescribing information.

### GENE ASSOCIATION

PTEN inactivation may predict benefit from mTOR inhibitors, such as everolimus, based on clinical data in various tumor types. For patients with prostate cancer, PTEN loss correlated with response to single-agent everolimus<sup>224</sup>. Retrospective clinical data suggest that patients with advanced breast cancer and PTEN inactivation, particularly in the context of HER2-positive disease, may benefit from everolimus combined with targeted therapy and/or chemotherapy<sup>225-227</sup>.

### SUPPORTING DATA

In an exploratory cohort of the BOLERO-2 Phase 3 study, the addition of everolimus to exemestane in the first line for hormone receptor-positive (HR+), HER2-negative (HER2-) breast cancer improve the median PFS compared to exemestane alone (11.5 vs. 4.1 months, HR = 0.39)<sup>228</sup>. Everolimus combined with exemestane as second-line therapy in the same setting also improved the median PFS

compared with exemestane in BOLERO-2 (7.8 vs. 3.2 months, HR = 0.45)<sup>229-231</sup>, and modestly improved the median PFS compared with everolimus alone in BOLERO-6 (8.4 vs. 6.8 months, HR = 0.74)<sup>232</sup>. Patients with HR+, HER2- breast cancer also benefited from everolimus combined with other antiestrogen therapies, including letrozole, tamoxifen, and anastrozole<sup>226,233-234</sup>. For patients with HR+, HER- breast cancer who progressed on antiestrogen therapies, addition of everolimus to the most recent endocrine therapy showed efficacy with 8% ORR and median PFS of 6.6 months<sup>235</sup>. For patients with HER2+ breast cancer, everolimus combined with trastuzumab plus paclitaxel did not significantly improve median PFS in the full study population (15.0 months with everolimus vs. 14.5 months with placebo), but increased PFS in the HR-negative subpopulation (20.3 vs. 13.1 months)<sup>236</sup>. For patients with trastuzumab-resistant HER2+ breast cancer, the addition of everolimus to trastuzumab plus vinorelbine prolonged median PFS (7.0 vs. 5.8 months)<sup>237</sup>. Patients with metastatic triple-negative breast cancer treated with everolimus plus carboplatin achieved a clinical benefit rate of 36% (9/25)<sup>238</sup>. Whereas frequent adverse events precluded a recommended Phase 2 dose and schedule for the combination of trametinib and everolimus in a Phase 1b trial for solid tumors<sup>239</sup>, a retrospective study for heavily pretreated patients with solid tumors reported tolerable regimens of the combination for 23/31 patients, with 16 patients treated >3 months and evaluable patients achieving a median PFS of 6.5 months<sup>240</sup>.

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by J. Keith Killian, M.D. | 23 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1021124-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

## Pazopanib

*Assay findings association*
**FGFR2**  
amplification

### AREAS OF THERAPEUTIC USE

Pazopanib is a tyrosine kinase inhibitor that targets VEGFRs, PDGFRs, FGFRs, KIT, ITK, LCK, and c-FMS. It is FDA approved for the treatment of advanced renal cell carcinoma and soft tissue sarcomas that have progressed after prior chemotherapy. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

FGFR2 amplification may predict sensitivity to pazopanib. In a Phase 2 study of pazopanib plus capecitabine and oxaliplatin for the treatment of patients with advanced gastric cancer, 6/7 patients with FGFR2 protein expression exhibited a PR, and PFS was significantly improved in patients with FGFR2 expression than in those without (8.5 vs. 5.6 months,  $p=0.050$ )<sup>241</sup>.

### SUPPORTING DATA

A Phase 2 clinical trial of pazopanib in breast cancer

reported 55% disease stabilization<sup>242</sup>. A Phase 2 study of heavily pretreated post-menopausal hormone receptor positive (HR+) breast cancer treated with a combination of pazopanib and nonsteroidal aromatase inhibitor reported 7% partial responses (PRs; 2/28) and 18% stable diseases (SDs; 5/28), with 7 patients having progression-free survival (PFS) greater than 6 months<sup>243</sup>. Phase 2 clinical trials of pazopanib with lapatinib in patients with HER2-positive breast cancer reported that the combination was associated with higher response rate than lapatinib alone but did not bring about an increase in PFS<sup>244-245</sup>. A multicenter single-arm Phase 2 study evaluating pazopanib combined with paclitaxel as neoadjuvant following doxorubicin/cyclophosphamide reported complete responses in 9% (6/67) and 38% (10/26) of patients with HR+ and triple-negative locally advanced breast cancer cases, respectively; however, a high level of toxicity led to discontinuation of pazopanib in 61% of patients<sup>246</sup>.

## Temsirolimus

*Assay findings association*
**PTEN**  
deletion exon 8

### AREAS OF THERAPEUTIC USE

Temsirolimus is an intravenous mTOR inhibitor that is FDA approved for the treatment of advanced renal cell carcinoma. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

PTEN inactivation may predict benefit from mTOR inhibitors, such as temsirolimus, based on clinical data in various tumor types. Out of 10 patients with metaplastic breast cancer and PTEN alterations, 2 cases responded to temsirolimus or everolimus plus doxorubicin and bevacizumab<sup>247-248</sup>. Temsirolimus achieved SD for 6/7 patients with PTEN-deficient cervical carcinoma<sup>249</sup>. Clinical studies in renal cell carcinoma<sup>250-251</sup>, glioblastoma<sup>252-253</sup>, or endometrial cancer<sup>254-257</sup> did not observe a correlation of PTEN deficiency with response to temsirolimus, although several patients with those tumor types and PTEN loss have benefited from mTOR inhibitors.

### SUPPORTING DATA

A Phase 1 trial examining the combination of

temsirolimus, liposomal doxorubicin, and bevacizumab in 74 patients with breast and gynecological malignancies reported that 37.9% of patients experienced either a CR (1.4%), PR (18.9%), or SD (17.6%); among 25 patients with PIK3CA mutation or PTEN loss, 52% experienced a CR, PR (36%), or SD (16%)<sup>258</sup>. Another Phase 1 trial including patients with several types of cancer reported a 42% incidence of complete or partial responses in patients with metastatic breast cancer<sup>259</sup>. However, a Phase 2 study of temsirolimus in pretreated patients with metastatic breast cancer reported minimal clinical activity and no association with PTEN protein or PIK3CA mutation status<sup>260</sup>. A Phase 3 placebo-controlled trial of letrozole plus oral temsirolimus as first-line endocrine therapy in postmenopausal women with locally advanced or metastatic breast cancer was terminated at the second interim since the addition of temsirolimus to letrozole did not improve PFS as a first-line therapy<sup>261</sup>. A study examining the efficacy of temsirolimus-involving regimens in 24 patients with mesenchymal/metaplastic breast cancer (MpBCs) reported 2 CRs, 4 PRs, 2 instances of SD longer than 6 months, and 4 instances of SD shorter than 6 months<sup>248</sup>.

**NOTE** Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

ORDERED TEST # ORD-1021124-01

**CLINICAL TRIALS**

**NOTE** Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.foundationmedicine.com/genomic-testing#support-services). Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

## GENE **FGFR2**

**RATIONALE**  
FGFR inhibitors may be relevant in tumors with alterations that activate FGFR2.

## ALTERATION amplification

### NCT04083976

**PHASE 2**

A Study of Erdafitinib in Participants With Advanced Solid Tumors and Fibroblast Growth Factor Receptor (FGFR) Gene Alterations

**TARGETS**  
FGFRs

**LOCATIONS:** San Salvador De Jujuy (Argentina), Cordoba (Argentina), Ijuí (Brazil), Ciudad Autónoma de Buenos Aires (Argentina), Buenos Aires (Argentina), Capital Federal (Argentina), Ciudad Autonoma de Buenos Aires (Argentina), Barretos (Brazil), La Plata Lpl Lpl (Argentina), Curitiba (Brazil)

### NCT04042116

**PHASE 1/2**

A Study to Evaluate Lucitanib in Combination With Nivolumab in Patients With a Solid Tumor

**TARGETS**  
FGFRs, VEGFRs, PD-1

**LOCATIONS:** Florida, North Carolina, Tennessee, Oklahoma, Ohio, Pennsylvania, New York, Massachusetts, Colorado, California

### NCT03797326

**PHASE 2**

Efficacy and Safety of Pembrolizumab (MK-3475) Plus Lenvatinib (E7080/MK-7902) in Previously Treated Participants With Select Solid Tumors (MK-7902-005/E7080-G000-224/LEAP-005)

**TARGETS**  
PD-1, FGFRs, KIT, PDGFRA, RET, VEGFRs

**LOCATIONS:** Florida, Tennessee, New Jersey, New York, Toronto (Canada), Wisconsin, South Dakota, Quebec (Canada), Winnipeg (Canada), California

### NCT04024436

**PHASE 2**

A Study of TAS-120 in Patients With Metastatic Breast Cancer

**TARGETS**  
FGFRs, ER

**LOCATIONS:** Florida, Texas, Tennessee, Missouri

### NCT03992131

**PHASE 1/2**

A Study to Evaluate Rucaparib in Combination With Other Anticancer Agents in Patients With a Solid Tumor (SEASTAR)

**TARGETS**  
PARP, FGFRs, VEGFRs, TOP1

**LOCATIONS:** Texas, Tennessee, Massachusetts



ORDERED TEST # ORD-1021124-01

**CLINICAL TRIALS**
**NCT03564691**
**PHASE 1**

Study of MK-4830 as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-4830-001)

**TARGETS**  
ITL4, FGFRs, KIT, PDGFRA, RET, VEGFRs, PD-1

**LOCATIONS:** Texas, Missouri, Ohio, New York, New Jersey, Michigan, Hamilton (Canada), Toronto (Canada), Ottawa (Canada), California

**NCT04233567**
**PHASE 2**

Infigratinib for the Treatment of Advanced or Metastatic Solid Tumors in Patients With FGFR Gene Mutations

**TARGETS**  
FGFR1, FGFR2, FGFR3

**LOCATIONS:** Ohio

**NCT02691767**
**PHASE NULL**

Study to Evaluate the Safety and Efficacy of Pazopanib, in Subject With Refractory Solid Tumors

**TARGETS**  
FGFR1, FGFR2, FGFR3, KIT, VEGFRs

**LOCATIONS:** Seoul (Korea, Republic of)

**NCT02450136**
**PHASE NULL**

Single-arm Study to Evaluate the Safety and Efficacy of Pazopanib, in Subjects With FGFR2 Amplification, FGFR2 Mutation Refractory Solid Tumors

**TARGETS**  
FGFR1, FGFR2, FGFR3, KIT, VEGFRs

**LOCATIONS:** Seoul (Korea, Republic of)

**NCT03547037**
**PHASE 1**

A Study to Evaluate the Safety, Pharmacokinetics, Pharmacodynamics, and Immunogenicity of JNJ-63723283, an Anti-Programmed Cell Death (PD)-1 Monoclonal Antibody, as Monotherapy or in Combination With Erdafitinib in Japanese Participants With Advanced Solid Cancers

**TARGETS**  
PD-1, FGFRs

**LOCATIONS:** Kashiwa (Japan), Chuo-Ku (Japan)

ORDERED TEST # ORD-1021124-01

CLINICAL TRIALS

**GENE**  
**PTEN**
**ALTERATION**  
deletion exon 8

**RATIONALE**  
PTEN loss or inactivating mutations may lead to increased activation of the PI3K-AKT-mTOR pathway and may indicate sensitivity to inhibitors

of this pathway. PTEN loss or inactivation may also predict sensitivity to PARP inhibitors.

**NCT04305496**
**PHASE 3**

Capiasertib+Fulvestrant vs Placebo+Fulvestrant as Treatment for Locally Advanced (Inoperable) or Metastatic HR+/HER2- Breast Cancer

**TARGETS**  
ER, AKTs

**LOCATIONS:** Lima (Peru), Arequipa (Peru), La Rioja (Argentina), Rosario (Argentina), Ciudad Autonoma De Buenos Aire (Argentina), Viedma (Argentina), Florida, Georgia, Tennessee

**NCT04191135**
**PHASE 2/3**

Study of Olaparib Plus Pembrolizumab Versus Chemotherapy Plus Pembrolizumab After Induction With First-Line Chemotherapy Plus Pembrolizumab in Triple Negative Breast Cancer (TNBC) (MK-7339-009/KEYLYNK-009)

**TARGETS**  
PD-1, PARP

**LOCATIONS:** Cali (Colombia), Medellin (Colombia), La Serena (Chile), Monteria (Colombia), Vina del Mar (Chile), Santiago (Chile), Barranquilla (Colombia), Temuco (Chile), Florida, Texas

**NCT03997123**
**PHASE 3**

Capiasertib+Paclitaxel as First Line Treatment for Patients With Locally Advanced or Metastatic TNBC

**TARGETS**  
AKTs

**LOCATIONS:** Rosario (Argentina), Londrina (Brazil), Goiania (Brazil), São José do Rio Preto (Brazil), Mar del Plata (Argentina), Caba (Argentina), Ciudad Autonoma De Buenos Aire (Argentina), Ciudad Autónoma de Bs. As. (Argentina), Barretos (Brazil), La Plata (Argentina)

**NCT04060862**
**PHASE 3**

A Study of Ipatasertib Plus Palbociclib and Fulvestrant Versus Placebo Plus Palbociclib and Fulvestrant in Hormone Receptor Positive and HER2 Negative Locally Advanced Unresectable or Metastatic Breast Cancer

**TARGETS**  
AKTs, CDK4, CDK6, ER

**LOCATIONS:** Porto Alegre (Brazil), Georgia, New Jersey, Hamilton (Canada), Calgary (Canada), Barcelona (Spain), Manchester (United Kingdom), Sutton (United Kingdom), London (United Kingdom), Malvern (Australia)

**NCT03598257**
**PHASE 2**

Radiation Therapy With or Without Olaparib in Treating Patients With Inflammatory Breast Cancer

**TARGETS**  
PARP

**LOCATIONS:** San Juan (Puerto Rico), Florida, Louisiana, Georgia, Texas, South Carolina

**NCT03994796**
**PHASE 2**

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

**TARGETS**  
ALK, ROS1, TRKA, TRKB, TRKC, CDK4, CDK6, PI3K, mTOR

**LOCATIONS:** Florida, Louisiana, Texas

ORDERED TEST # ORD-1021124-01

**CLINICAL TRIALS**
**NCT02498613**
**PHASE 2**

A Phase 2 Study of Cediranib in Combination With Olaparib in Advanced Solid Tumors

**TARGETS**  
PARP, VEGFRs

**LOCATIONS:** Florida, Texas, Tennessee, Virginia, Connecticut, Massachusetts, Toronto (Canada), California

**NCT03907969**
**PHASE 1/2**

A Clinical Trial to Evaluate AZD7648 Alone and in Combination With Other Anti-cancer Agents in Patients With Advanced Cancers

**TARGETS**  
PARP, DNA-PK

**LOCATIONS:** Texas, Connecticut, Newcastle upon Tyne (United Kingdom)

**NCT03939897**
**PHASE 1/2**

Testing the Addition of Copanlisib to Usual Treatment (Fulvestrant and Abemaciclib) in Metastatic Breast Cancer - Dose-Finding Study

**TARGETS**  
PI3K, CDK4, CDK6, ER

**LOCATIONS:** Alabama, Kentucky, Missouri, Ohio, New York, California

**NCT03280563**
**PHASE 1/2**

A Study of Multiple Immunotherapy-Based Treatment Combinations in Hormone Receptor (HR)-Positive Human Epidermal Growth Factor Receptor 2 (HER2)-Negative Breast Cancer

**TARGETS**  
PD-L1, ER, HDAC, AKTs, CDK4, CDK6

**LOCATIONS:** North Carolina, Tennessee, Maryland, Pennsylvania, New York, Illinois, California

ORDERED TEST # ORD-1021124-01

**APPENDIX**
**Variants of Unknown Significance**

**NOTE** One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

**AXIN1**  
R298Q

**CIC**  
P1551A

**CTNNB1**  
E477Q

**EPHB1**  
E65K

**FANCC**  
I290V

**FH**  
F225S

**KMT2A (MLL)**  
A53V

**LTK**  
I662T

**MAF**  
L380M

**MAP3K1**  
L78P

**MERTK**  
E712D

**NSD3 (WHSC1L1)**  
A1293V

**RET**  
Q70R

ORDERED TEST # ORD-1021124-01

**APPENDIX**
**Genes Assayed in FoundationOne®CDx**

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

**DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS**

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTB	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXO2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMS5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

**DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS**

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPR352

\*TERC is an NCRNA

\*\*Promoter region of TERT is interrogated

**ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS**

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)



ORDERED TEST # ORD-1021124-01

**APPENDIX**

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a., Cipalstraat 3, 2440 Geel, Belgium.


**ABOUT FOUNDATIONONE CDx**

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:  
[www.rochefoundationmedicine.com/f1cdxtech](http://www.rochefoundationmedicine.com/f1cdxtech).

**INTENDED USE**

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

**TEST PRINCIPLES**

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes. Using an Illumina® HiSeq platform, hybrid

capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

**THE REPORT**

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

**Diagnostic Significance**

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

**Qualified Alteration Calls (Equivocal and Subclonal)**

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

**Ranking of Alterations and Therapies Biomarker and Genomic Findings**

Therapies are ranked based on the following criteria: Therapies with clinical benefit in patient's tumor type (ranked alphabetically within each NCCN category) followed by therapies with clinical benefit in other tumor type (ranked alphabetically within each NCCN category).

**Clinical Trials**

Pediatric trial qualification → Geographical proximity → Later trial phase.

**NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION**

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) ([www.nccn.org](http://www.nccn.org)). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2020. All rights reserved. To view the most recent and complete version of the guideline, go online to [NCCN.org](http://NCCN.org). NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

**Limitations**

1. The MSI-H/MSS designation by FMI F1CDx test is based on genome wide analysis of 95 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines. The threshold for MSI-H/MSS was determined by analytical concordance to comparator assays (IHC and PCR) using uterine, cecum and colorectal cancer FFPE tissue. The clinical validity of the qualitative MSI designation has not been established. For Microsatellite Instability (MSI) results, confirmatory testing using a validated orthogonal method should be considered.
2. TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by J. Keith Killian, M.D. | 23 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1021124-01

APPENDIX

About FoundationOne®CDx

of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/P170019B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf). The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.

3. The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

### VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

#### Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31

INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

\*Interquartile Range = 1<sup>st</sup> Quartile to 3<sup>rd</sup> Quartile

### VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters with no conflicts), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

### LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

### NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

### NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

### TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or

none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

### SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 2.2.0

ORDERED TEST # ORD-1021124-01

APPENDIX

About FoundationOne®CDx

The median exon coverage for this sample is 937x

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by J. Keith Killian, M.D. | 23 February 2021  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1021124-01

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Anbazhagan R, et al. Clin. Cancer Res. (1999) PMID: 10213220
7. Adem C, et al. Int. J. Cancer (2003) PMID: 14520695
8. Horimoto Y, et al. Cancer Sci (2020) PMID: 32449246
9. Heeke AL, et al. Breast Cancer Res Treat (2020) PMID: 32776290
10. Kurata K, et al. Breast Cancer (2020) PMID: 31907878
11. Sivapiragasam A, et al. Cancer Med (2020) PMID: 33314633
12. Walsh MD, et al. Clin. Cancer Res. (2010) PMID: 20215533
13. Risinger JI, et al. Cancer (1996) PMID: 8646682
14. de Leeuw WJ, et al. Cancer Res. (2003) PMID: 12615735
15. Shanley S, et al. Fam. Cancer (2009) PMID: 19123071
16. Buerki N, et al. Genes Chromosomes Cancer (2012) PMID: 22034109
17. Yee CJ, et al. Cancer Res. (1994) PMID: 8137273
18. Kamat N, et al. BMC Cancer (2012) PMID: 22928966
19. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
20. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
21. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
22. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
23. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
24. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
25. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
26. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
27. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
28. Cristescu R, et al. Science (2018) PMID: 30309915
29. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
30. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
31. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
32. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
33. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
34. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
35. Legrand et al., 2018; ASCO Abstract 12000
36. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
37. Barroso-Sousa R, et al. Ann. Oncol. (2020) PMID: 32067680
38. Nature (2012) PMID: 23000897
39. Sokol ES, et al. Ann. Oncol. (2019) PMID: 30423024
40. Chumsri S, et al. J Natl Compr Canc Netw (2020) PMID: 32380464
41. Haricharan S, et al. Breast Cancer Res. Treat. (2014) PMID: 24839032
42. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
43. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
44. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
45. Rizvi NA, et al. Science (2015) PMID: 25765070
46. Johnson BE, et al. Science (2014) PMID: 24336570
47. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
48. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
49. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
50. Heitz E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
51. Nature (2012) PMID: 22810696
52. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
53. Tabernero J, et al. J. Clin. Oncol. (2015) PMID: 26324363
54. Krook MA, et al. Cold Spring Harb Mol Case Stud (2019) PMID: 31371345
55. Abou-Alfa GK, et al. Lancet Oncol. (2020) PMID: 32203698
56. Morizane et al., 2020; ASCO GI Abstract 538
57. Borad MJ, et al. PLoS Genet. (2014) PMID: 24550739
58. Liao RG, et al. Cancer Res. (2013) PMID: 23786770
59. Gozgit JM, et al. Mol. Cancer Ther. (2012) PMID: 22238366
60. Pearson A, et al. Cancer Discov (2016) PMID: 27179038
61. Van Cutsem E, et al. Ann. Oncol. (2017) PMID: 29177434
62. Aggarwal C, et al. J Thorac Oncol (2019) PMID: 31195180
63. Voss MH, et al. Clin. Cancer Res. (2019) PMID: 30745300
64. Nogova L, et al. J. Clin. Oncol. (2017) PMID: 27870574
65. Goyal L, et al. Cancer Discov (2019) PMID: 31109923
66. Bahlada R, et al. Clin. Cancer Res. (2019) PMID: 31088831
67. Heiskanen M, et al. Anal Cell Pathol (2001) PMID: 11564899
68. Adnane J, et al. Oncogene (1991) PMID: 1851551
69. Turner N, et al. Oncogene (2010) PMID: 20101236
70. Lee HJ, et al. Ann. Surg. Oncol. (2014) PMID: 24385208
71. Sun S, et al. J Surg Oncol (2012) PMID: 22006548
72. Templeton AJ, et al. Cancer Treat. Rev. (2014) PMID: 25217796
73. Powers CJ, et al. Endocr. Relat. Cancer (2000) PMID: 11021964
74. Turner N, et al. Nat. Rev. Cancer (2010) PMID: 20094046
75. Gao J, et al. Sci Signal (2013) PMID: 23550210
76. Tokunaga R, et al. Oncotarget (2016) PMID: 26933914
77. André F, et al. Clin. Cancer Res. (2013) PMID: 23658459
78. Courtney KD, et al. J. Clin. Oncol. (2010) PMID: 20085938
79. Wu R, et al. Clin. Cancer Res. (2011) PMID: 21903772
80. Simpson L, et al. Exp. Cell Res. (2001) PMID: 11237521
81. Dreyling M, et al. Ann. Oncol. (2017) PMID: 28633365
82. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
83. Wee S, et al. Proc. Natl. Acad. Sci. U.S.A. (2008) PMID: 18755892
84. Jia S, et al. Nature (2008) PMID: 18594509
85. Schmit F, et al. Proc. Natl. Acad. Sci. U.S.A. (2014) PMID: 24737887
86. Janku et al., 2018; ESMO Abstract 418PD
87. Dent et al., 2018; ASCO Abstract 1008
88. Schmid et al., 2018; ASCO Abstract 1007
89. de Bono JS, et al. Clin. Cancer Res. (2019) PMID: 30037818
90. Saura C, et al. Cancer Discov (2017) PMID: 27872130
91. Schmid P, et al. J. Clin. Oncol. (2019) PMID: 31841354
92. Kim SB, et al. Lancet Oncol (2017) PMID: 28800861
93. Kim et al., 2020; SABCS Abstract GS3-04
94. Mendes-Pereira AM, et al. EMBO Mol Med (2009) PMID: 20049735
95. Shen Y, et al. Clin. Cancer Res. (2013) PMID: 23881923
96. Chatterjee P, et al. PLoS ONE (2013) PMID: 23565244
97. McCormick A, et al. Int. J. Gynecol. Cancer (2016) PMID: 26905328
98. Forster MD, et al. Nat Rev Clin Oncol (2011) PMID: 21468130
99. Gruber et al., 2019; ASCO Abstract 3006
100. Dougherty et al., 2014; ASCO Abstract 5536
101. Piha-Paul et al., 2018; AACR Abstract A096
102. Ihnen M, et al. Mol. Cancer Ther. (2013) PMID: 23729402
103. Sandhu SK, et al. Lancet Oncol. (2013) PMID: 23810788
104. Zhao J, et al. Nat. Med. (2019) PMID: 30742119
105. George S, et al. Immunity (2017) PMID: 28228279
106. Parikh AR, et al. Lung Cancer (Auckl) (2018) PMID: 29844707
107. Peng W, et al. Cancer Discov (2016) PMID: 26645196
108. Costa C, et al. Cancer Discov (2019) PMID: 31594766
109. Le X, et al. Cancer Discov (2016) PMID: 27604488
110. Juric D, et al. Nature (2015) PMID: 25409150
111. Chen SH, et al. Oncogene (2018) PMID: 29059158
112. Hennessy BT, et al. Cancer Res. (2009) PMID: 19435916
113. Mercapide J, et al. Mol. Carcinog. (2002) PMID: 12203362
114. Hohensee I, et al. Am. J. Pathol. (2013) PMID: 23665199
115. Perez EA, et al. J. Clin. Oncol. (2013) PMID: 23650412
116. Tsutsui S, et al. Oncology (2005) PMID: 16020969
117. Zhang HY, et al. Oncol Lett (2013) PMID: 23946797
118. Capodanno A, et al. Hum. Pathol. (2009) PMID: 19428048
119. Campbell RB, et al. J. Biol. Chem. (2003) PMID: 12857747
120. Rodríguez-Escudero I, et al. Hum. Mol. Genet. (2011) PMID: 21828076
121. He X, et al. Cancer Res. (2013) PMID: 23475934
122. Han SY, et al. Cancer Res. (2000) PMID: 10866302
123. Myers MP, et al. Proc. Natl. Acad. Sci. U.S.A. (1998) PMID: 9811831
124. Pradella LM, et al. BMC Cancer (2014) PMID: 24498881
125. Kim JS, et al. Mol. Cell. Biol. (2011) PMID: 21536651
126. Denning G, et al. Oncogene (2007) PMID: 17213812
127. Hlobilkova A, et al. Anticancer Res. ( ) PMID: 16619501
128. Redfern RE, et al. Protein Sci. (2010) PMID: 20718038
129. Shenoy S, et al. PLoS ONE (2012) PMID: 22505997
130. Wang Y, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19329485
131. Okumura K, et al. J. Biol. Chem. (2006) PMID: 16829519
132. Lee JO, et al. Cell (1999) PMID: 10555148
133. Maxwell GL, et al. Cancer Res. (1998) PMID: 9635567
134. Risinger JI, et al. Clin. Cancer Res. (1998) PMID: 9865913
135. Kato H, et al. Clin. Cancer Res. (2000) PMID: 11051241
136. Fenton TR, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22891331
137. Ngeow J, et al. J. Clin. Endocrinol. Metab. (2012) PMID: 23066114
138. Lobo GP, et al. Hum. Mol. Genet. (2009) PMID: 19457929
139. Liu J, et al. Oncogene (2014) PMID: 23995781
140. Maehama T, et al. Annu. Rev. Biochem. (2001) PMID: 11395408
141. De Vivo I, et al. J. Med. Genet. (2000) PMID: 10807691
142. Ramaswamy S, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10051603
143. Liu JL, et al. Mol. Cell. Biol. (2005) PMID: 15988030
144. Karoui M, et al. Br. J. Cancer (2004) PMID: 15026806
145. Gil A, et al. PLoS ONE (2015) PMID: 25875300
146. Furnari FB, et al. Cancer Res. (1998) PMID: 9823298
147. Spinelli L, et al. J. Med. Genet. (2015) PMID: 25527629
148. Mingo J, et al. Eur. J. Hum. Genet. (2018) PMID: 29706633
149. Wang Q, et al. J. Mol. Graph. Model. (2010) PMID: 20538496
150. Andrés-Pons A, et al. Cancer Res. (2007) PMID: 17942903
151. Butler MG, et al. J. Med. Genet. (2005) PMID: 15805158
152. Georgescu MM, et al. Proc. Natl. Acad. Sci. U.S.A. (1999)

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by J. Keith Killian, M.D. | 23 February 2021  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # **ORD-1021124-01**

- pmid: 10468583
153. Staal FJ, et al. Br. J. Cancer (2002) pmid: 12085208
  154. Nguyen HN, et al. Oncogene (2014) pmid: 24292679
  155. Rahdar M, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) pmid: 19114656
  156. Das S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) pmid: 12808147
  157. Wang X, et al. Biochem. J. (2008) pmid: 18498243
  158. Valiente M, et al. J. Biol. Chem. (2005) pmid: 15951562
  159. Nguyen HN, et al. Oncogene (2015) pmid: 25263454
  160. Blumenthal GM, et al. Eur. J. Hum. Genet. (2008) pmid: 18781191
  161. Orloff MS, et al. Oncogene (2008) pmid: 18794875
  162. Zbuk KM, et al. Nat. Rev. Cancer (2007) pmid: 17167516
  163. Hirai H, et al. Cancer Biol. Ther. (2010) pmid: 20107315
  164. Bridges KA, et al. Clin. Cancer Res. (2011) pmid: 21799033
  165. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) pmid: 21389100
  166. Osman AA, et al. Mol. Cancer Ther. (2015) pmid: 25504633
  167. Xu L, et al. Mol. Cancer Ther. (2002) pmid: 12489850
  168. Xu L, et al. Mol. Med. (2001) pmid: 11713371
  169. Camp ER, et al. Cancer Gene Ther. (2013) pmid: 23470564
  170. Kim SS, et al. Nanomedicine (2015) pmid: 25240597
  171. Pirollo KF, et al. Mol. Ther. (2016) pmid: 27357628
  172. Hajdenberg et al., 2012; ASCO Abstract e15010
  173. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27601554
  174. Moore et al., 2019; ASCO Abstract 5513
  175. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27998224
  176. Oza et al., 2015; ASCO Abstract 5506
  177. Lee J, et al. Cancer Discov (2019) pmid: 31315834
  178. Méndez E, et al. Clin. Cancer Res. (2018) pmid: 29535125
  179. Ma CX, et al. J. Clin. Invest. (2012) pmid: 22446188
  180. Kwok M, et al. Blood (2016) pmid: 26563132
  181. Boudny M, et al. Haematologica (2019) pmid: 30975914
  182. Dillon MT, et al. Mol. Cancer Ther. (2017) pmid: 28062704
  183. Middleton FK, et al. Cancers (Basel) (2018) pmid: 30127241
  184. Banerji S, et al. Nature (2012) pmid: 22722202
  185. Stephens PJ, et al. Nature (2012) pmid: 22722201
  186. Alsner J, et al. Acta Oncol (2008) pmid: 18465328
  187. Alkam Y, et al. Histopathology (2013) pmid: 24004112
  188. Uji K, et al. Cancer Lett. (2014) pmid: 23973262
  189. Olivier M, et al. Clin. Cancer Res. (2006) pmid: 16489069
  190. Végran F, et al. PLoS ONE (2013) pmid: 23359294
  191. Walsh T, et al. JAMA (2006) pmid: 16551709
  192. Garber JE, et al. J. Clin. Oncol. (2005) pmid: 15637391
  193. Apostolou P, et al. Biomed Res Int (2013) pmid: 23586058
  194. Jaiswal S, et al. N. Engl. J. Med. (2014) pmid: 25426837
  195. Genovese G, et al. N. Engl. J. Med. (2014) pmid: 25426838
  196. Xie M, et al. Nat. Med. (2014) pmid: 25326804
  197. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) pmid: 28669404
  198. Severson EA, et al. Blood (2018) pmid: 29678827
  199. Fuster JJ, et al. Circ. Res. (2018) pmid: 29420212
  200. Hematology Am Soc Hematol Educ Program (2018) pmid: 30504320
  201. Chabon JJ, et al. Nature (2020) pmid: 32269342
  202. Razavi P, et al. Nat. Med. (2019) pmid: 31768066
  203. Brown CJ, et al. Nat. Rev. Cancer (2009) pmid: 19935675
  204. Joerger AC, et al. Annu. Rev. Biochem. (2008) pmid: 18410249
  205. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) pmid: 12826609
  206. Kamada R, et al. J. Biol. Chem. (2011) pmid: 20978130
  207. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) pmid: 28472496
  208. Yamada H, et al. Carcinogenesis (2007) pmid: 17690113
  209. Bougeard G, et al. J. Clin. Oncol. (2015) pmid: 26014290
  210. Sorrell AD, et al. Mol Diagn Ther (2013) pmid: 23355100
  211. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) pmid: 11219776
  212. Kleihues P, et al. Am. J. Pathol. (1997) pmid: 9006316
  213. Gonzalez KD, et al. J. Clin. Oncol. (2009) pmid: 19204208
  214. Lalloo F, et al. Lancet (2003) pmid: 12672316
  215. Mandelker D, et al. Ann. Oncol. (2019) pmid: 31050713
  216. Siefker-Radtke et al., 2018; ASCO Abstract 4503
  217. Park et al., 2019; ASCO Abstract 4117
  218. Soria et al., 2017; ASCO Abstract 4074
  219. Perera TPS, et al. Mol. Cancer Ther. (2017) pmid: 28341788
  220. Karkera JD, et al. Mol. Cancer Ther. (2017) pmid: 28416604
  221. Llorca Y, et al. N. Engl. J. Med. (2019) pmid: 31340094
  222. Qin A, et al. J Thorac Oncol (2019) pmid: 30267839
  223. Di Stefano AL, et al. Clin. Cancer Res. (2015) pmid: 25609060
  224. Templeton AJ, et al. Eur. Urol. (2013) pmid: 23582881
  225. André F, et al. J. Clin. Oncol. (2016) pmid: 27091708
  226. Wheeler JJ, et al. Oncotarget (2014) pmid: 24912489
  227. Janku F, et al. Cell Rep (2014) pmid: 24440717
  228. Beck JT, et al. Breast Cancer Res. Treat. (2014) pmid: 24362951
  229. Yardley DA, et al. Adv Ther (2013) pmid: 24158787
  230. Baselga J, et al. N. Engl. J. Med. (2012) pmid: 22149876
  231. Piccart M, et al. Ann. Oncol. (2014) pmid: 25231953
  232. Jerusalem G, et al. JAMA Oncol (2018) pmid: 29862411
  233. Baselga J, et al. J. Clin. Oncol. (2009) pmid: 19380449
  234. Bachelot T, et al. J. Clin. Oncol. (2012) pmid: 22565002
  235. Yardley DA, et al. Clin. Breast Cancer (2019) pmid: 31932237
  236. Hurvitz SA, et al. Lancet Oncol. (2015) pmid: 26092818
  237. André F, et al. Lancet Oncol. (2014) pmid: 24742739
  238. Singh J, et al. Breast Cancer Res. (2014) pmid: 24684785
  239. Tolcher AW, et al. Ann. Oncol. (2015) pmid: 25344362
  240. Patterson et al., 2018; AACR Abstract 3891
  241. Kim ST, et al. J. Cancer Res. Clin. Oncol. (2016) pmid: 26983912
  242. Taylor SK, et al. Oncologist (2010) pmid: 20682606
  243. Majure et al., 2016; ASCO Abstract 560
  244. Johnston SR, et al. Breast Cancer Res. Treat. (2013) pmid: 23283526
  245. Cristofanilli M, et al. Breast Cancer Res. Treat. (2013) pmid: 23239151
  246. Tan AR, et al. Breast Cancer Res. Treat. (2015) pmid: 25542269
  247. Basho et al., 2015; SABCS Abstract P3-14-02
  248. Moulder S, et al. Ann. Oncol. (2015) pmid: 25878190
  249. Tinker AV, et al. Gynecol. Oncol. (2013) pmid: 23672928
  250. Figlin RA, et al. Cancer (2009) pmid: 19526589
  251. Cho D, et al. Clin Genitourin Cancer (2007) pmid: 17956710
  252. Galanis E, et al. J. Clin. Oncol. (2005) pmid: 15998902
  253. Cloughesy TF, et al. PLoS Med. (2008) pmid: 18215105
  254. Oza AM, et al. J. Clin. Oncol. (2011) pmid: 21788564
  255. Mackay HJ, et al. Cancer (2014) pmid: 24166148
  256. Fleming GF, et al. Gynecol. Oncol. (2014) pmid: 24456823
  257. Tsoref D, et al. Gynecol. Oncol. (2014) pmid: 25173583
  258. Moroney JW, et al. Clin. Cancer Res. (2011) pmid: 21890452
  259. Moroney J, et al. Clin. Cancer Res. (2012) pmid: 22927482
  260. Fleming GF, et al. Breast Cancer Res. Treat. (2012) pmid: 22245973
  261. Wolff AC, et al. J. Clin. Oncol. (2013) pmid: 23233719