

Lab 01 Conversion from Matlab

Aaron Swoboda

2025-04-15

This file aims to convert a series of Matlab scripts and data files into a R Markdown file that replicates the Tol (2023) Lab 1 for the economics of climate change. <https://github.com/rtol/ClimateEconomics>

The Matlab files this replicates are:

- MLIAM01.m
- MLIAM.mat
- init01.m
- MRHparam.m
- MRH.m
- RFparam.m
- RadForc.m
- STparam.m
- ST.m

Overview - the MLIAM01.m script

```
# clear all
# load MLIAM (Step 1 below)
# init01 (Step 2 below)
#
# (Loop is Step 3)
# for t=StartYear+1:EndYear,
#   i = t - StartYear + 1;
#   MRHbox(i,:) = MRH(MRHbox(i-1,:),CO2emit(i-1),MRHlife,MRHshare,CO2convert);
#   CO2conc(i) = sum(MRHbox(i,:));
#   RF(i) = RadForc(CO2conc(i),RF0,CO20);
#   [atmtemp(i) oceantemp(i)] = ST(atmtemp(i-1),oceantemp(i-1),RF(i),STpar);
# end
```

Step 1 - load the MLIAM.mat file

```
library(R.matlab)
```

```
## R.matlab v3.7.0 (2022-08-25 21:52:34 UTC) successfully loaded. See ?R.matlab for help.
##
## Attaching package: 'R.matlab'
## The following objects are masked from 'package:base':
##
##      getOption, isOpen
```

```

MLIAM <- readMat("../TolMatlabFiles/MLIAM.mat")
ls.str()

## MLIAM : List of 11
## $ CO2global : num [1:551, 1] 0 3 3 3 3 3 3 3 3 3 ...
## $ CO2emit : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ CO2concobs : num [1:689, 1:2] 2006 2005 2004 2003 2002 ...
## $ population : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ output : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ atmtempobs : num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## $ oceantempobs: num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## $ CO2emitobs : num [1:551, 1] 0 3 3 3 3 3 3 3 3 3 ...
## $ outputobs : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ energy : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ NReg : num [1, 1] 3

# CO2global <- MLIAM$CO2global # should we use this?
CO2emit <- MLIAM$CO2emit
# CO2concobs <- MLIAM$CO2concobs
# population <- MLIAM$population
# output <- MLIAM$output
# atmtempobs <- MLIAM$atmtempobs
# oceantempobs <- MLIAM$oceantempobs
# CO2emitobs <- MLIAM$CO2emitobs # should we use this?
# outputobs <- MLIAM$outputobs
# energy <- MLIAM$energy
# NReg <- MLIAM$NReg

```

Step 2 - run the init01.m script

```

# %initialize
#
# climsens = 4.260547;
# EndYear = 2300;
# StartYear = 1750;
# NYear = EndYear-StartYear+1;
# year = zeros(NYear,1);
# year(1) = StartYear;
# for i=2:NYear
# year(i) = year(i-1)+1;
# end
#
# MRHparam; #Step 2a
# RFparam; # Step 2b
# STparam; # Step 2c

```

and now in R

```

climsens = 4.260547;
EndYear = 2300;
StartYear = 1750;
year = StartYear:EndYear
NYear = length(year)

```

Step 2a - MRHparam

```
# %MRHparam
#
# MRHlife(1) = 0;
# MRHlife(2) = 1-exp(-1/363);
# MRHlife(3) = 1-exp(-1/74);
# MRHlife(4) = 1-exp(-1/17);
# MRHlife(5) = 1-exp(-1/2);
#
# MRH1750(1) = 275;
# MRH1750(2) = 0;
# MRH1750(3) = 0;
# MRH1750(4) = 0;
# MRH1750(5) = 0;
#
# MRHshare(1) = 0.13;
# MRHshare(2) = 0.20;
# MRHshare(3) = 0.32;
# MRHshare(4) = 0.25;
# MRHshare(5) = 0.10;
#
# CO2convert = 1/2.13/1000;
#
# MRHbox = zeros(NYear,5); %five boxes
# MRHbox(1,:) = MRH1750;
# CO2conc = zeros(NYear,1);
# CO20 = sum(MRH1750);
# CO2conc(1) = CO20;
```

in R

```
# %MRHparam
MRHlife <- c(0, 1-exp(-1/363), 1-exp(-1/74), 1-exp(-1/17), 1-exp(-1/2))
#
MRH1750 <- c(275, 0, 0, 0, 0);
#
MRHshare <- c(0.13, 0.20, 0.32, 0.25, 0.10);
#
CO2convert = 1/2.13/1000;
#
MRHbox = matrix(0, NYear, 5); # "five boxes" for each year
MRHbox[1, ] = MRH1750;
CO2conc = matrix(0, NYear, 1);
CO20 = sum(MRH1750);
CO2conc[1] = CO20;
```

Step 2b - RFparam

```
# %RFparam
#
# RFO = 5.35;
# RF = zeros(NYear,1);
# RF(1) = RadForc(CO20,RFO,CO20); #Need the RadForc function
```

First, create the RadForc function

```
### RadForc
RadForc <-function(CO2,RF0,CO20) RF0*log(CO2/CO20);
```

Then run the RFparam

```
# %RFparam
#
RF0 = 5.35;
RF = matrix(0, NYear, 1);
RF[1] = RadForc(CO20, RF0, CO20) #Needs the RadForc function
```

Step 2c - ST param

```
# %STparam
#
# atmtmp0 = 0;
# oceantemp0 = 0;
#
# atmtmp = zeros(NYear,1);
# oceantemp = zeros(NYear,1);
#
# atmtmp(1) = atmtmp0;
# oceantemp(1) = oceantemp0;
#
# STpar(1) = climsens/RF0/log(2);
# STpar(2) = 0.0256;
# STpar(3) = 0.00738;
# STpar(4) = 0.00568;

# %STparam

atmtmp0 = 0;
oceantemp0 = 0;

atmtmp = matrix(0, NYear, 1);
oceantemp = matrix(0, NYear, 1);

atmtmp[(1)] = atmtmp0;
oceantemp[(1)] = oceantemp0;

STpar <- c(climsens/RF0/log(2), 0.0256, 0.00738, 0.00568)
```

Loop is Step 3

But we need some functions - MRH and ST

```
### MRH - Maier-Reimer Hasselmann model
# function CO2connew = MRH(CO2concold,CO2emit,CO2life,CO2share,CO2convert)
# %function CO2connew = MRH(CO2concold,CO2emit,CO2life,CO2share,CO2convert)
# %
# %updates the atmospheric concentration of carbon dioxide using the
# %Maier-Reimer Hasselmann model
#
# CO2connew = (1-CO2life).*CO2concold + CO2convert*CO2share*CO2emit;
```

MRH in R

```
### MRH - Maier-Reimer Hasselmann model
```

```
MRH <- function(CO2concold, CO2emit, CO2life, CO2share, CO2convert) {  
  CO2concnew = (1-CO2life)*CO2concold + CO2convert*CO2share*CO2emit  
  CO2concnew  
}
```

```
### Schneider-Thompson model of Ocean and Atm Temps
```

```
# function [atmtempnew oceantempnew] = ST(atmtempold, oceantempold, radforc, STpar)
```

```
# %function [atmtempnew oceantempnew] = ST(atmtempold, oceantempold, radforc, STpar)
```

```
# %
```

```
# %updates the temperature of the atmosphere and the ocean using the Schneider-Thompson model
```

```
#
```

```
# atmtempnew = atmtempold + STpar(2)*(STpar(1)*radforc-atmtempold) + STpar(3)*(oceantempold-atmtempold)
```

```
# oceantempnew = oceantempold + STpar(4)*(atmtempold-oceantempold)
```

```
### Schneider-Thompson model of Ocean and Atm Temps
```

```
# updates the temperature of the atmosphere and the ocean using the Schneider-Thompson model
```

```
ST <- function(atmtempold, oceantempold, radforc, STpar) {
```

```
  atmtempnew = atmtempold +
```

```
    STpar[2]*(STpar[1]*radforc-atmtempold) +
```

```
    STpar[3]*(oceantempold-atmtempold)
```

```
  oceantempnew = oceantempold +
```

```
    STpar[4]*(atmtempold-oceantempold)
```

```
  temps <- c(atmtempnew, oceantempnew)
```

```
  names(temps) <- c("atm", "ocean")
```

```
  temps
```

```
}
```

Step 3 - the main loop!

```
# for t=StartYear+1:EndYear,
```

```
# i = t - StartYear + 1;
```

```
# MRHbox(i,:) = MRH(MRHbox(i-1,:), CO2emit(i-1), MRHlife, MRHshare, CO2convert); # need the MRH function
```

```
# CO2conc(i) = sum(MRHbox(i,:));
```

```
# RF(i) = RadForc(CO2conc(i), RF0, CO20);
```

```
# [atmtemp(i) oceantemp(i)] = ST(atmtemp(i-1), oceantemp(i-1), RF(i), STpar); # need the ST function
```

```
# end
```

in R

```
# for t=StartYear+1:EndYear,
```

```
# i = t - StartYear + 1;
```

```
for (i in 2:(EndYear-StartYear+1)) {
```

```
  MRHbox[i, ] = MRH(MRHbox[i-1, ], CO2emit[i-1], MRHlife, MRHshare, CO2convert); # need the MRH function
```

```
  CO2conc[i] = sum(MRHbox[i, ]);
```

```
  RF[i] = RadForc(CO2conc[i], RF0, CO20);
```

```
  temps <- ST(atmtemp[i-1], oceantemp[i-1], RF[i], STpar); # need the ST function
```

```
  atmtemp[i] <- temps[["atm"]]
```

```
  oceantemp[i] <- temps[["ocean"]]
```

```
}
```

End of Matlab Scripts

```
ls.str()

## atmtemp : num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## atmtemp0 : num 0
## climsens : num 4.26
## CO20 : num 275
## CO2conc : num [1:551, 1] 275 275 275 275 275 275 275 275 275 275 ...
## CO2convert : num 0.000469
## CO2emit : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## EndYear : num 2300
## i : int 551
## MLIAM : List of 11
## $ CO2global : num [1:551, 1] 0 3 3 3 3 3 3 3 3 3 ...
## $ CO2emit : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ CO2concoobs : num [1:689, 1:2] 2006 2005 2004 2003 2002 ...
## $ population : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ output : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ atmtempobs : num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## $ oceantempobs : num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## $ CO2emitobs : num [1:551, 1] 0 3 3 3 3 3 3 3 3 3 ...
## $ outputobs : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ energy : num [1:551, 1:3] 0 0 0 0 0 0 0 0 0 0 ...
## $ NReg : num [1, 1] 3
## MRH : function (CO2concoold, CO2emit, CO2life, CO2share, CO2convert)
## MRH1750 : num [1:5] 275 0 0 0 0
## MRHbox : num [1:551, 1:5] 275 275 275 275 275 275 275 275 275 275 ...
## MRHlife : num [1:5] 0 0.00275 0.01342 0.05713 0.39347
## MRHshare : num [1:5] 0.13 0.2 0.32 0.25 0.1
## NYear : int 551
## oceantemp : num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## oceantemp0 : num 0
## RadForc : function (CO2, RF0, CO20)
## RF : num [1:551, 1] 0 0 0 0 0 0 0 0 0 0 ...
## RF0 : num 5.35
## ST : function (atmtempold, oceantempold, radforc, STpar)
## StartYear : num 1750
## STpar : num [1:4] 1.14891 0.0256 0.00738 0.00568
## temps : Named num [1:2] 1.09 1.05
## year : int [1:551] 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 ...
```

How should we display the output?

Plot the output data for comparison?

Are these outouts consistent with the Excel file results?

Make sure we are using the correct inputs?

- CO2emit vs. CO2emitobs vs. CO2global