

QUESTION PAPER CODE 65/1/3
EXPECTED ANSWER/VALUE POINTS

SECTION – A

Question Numbers 1 to 20 carry 1 mark each.

Question Numbers 1 to 10 are multiple choice type questions.

Select the correct option.

Q.No.	Marks
<p>1. The matrix $\begin{bmatrix} 2 & -1 & 3 \\ \lambda & 0 & 7 \\ -1 & 1 & 4 \end{bmatrix}$ is not invertible for</p> <p>(A) $\lambda = -1$ (B) $\lambda = 0$ (C) $\lambda = 1$ (D) $\lambda \in \mathbb{R} - \{1\}$</p> <div>Ans: (C) $\lambda = 1$</div>	1
<p>2. The number of arbitrary constants in the particular solution of a differential equation of second order is (are)</p> <p>(A) 0 (B) 1 (C) 2 (D) 3</p> <div>Ans: (A) 0</div>	1
<p>3. The value of $\tan^{-1} \left(\tan \frac{7\pi}{6} \right)$ is</p> <p>(A) $\frac{\pi}{6}$ (B) $\frac{\pi}{2}$ (C) $\frac{\pi}{3}$ (D) $\frac{7\pi}{6}$</p> <div>Ans: (A) $\frac{\pi}{6}$</div>	1
<p>4. The corner points of the feasible region determined by the system of linear inequalities are (0, 0), (4, 0), (2, 4) and (0, 5). If the maximum value of $z = ax + by$, where $a, b > 0$ occurs at both (2, 4) and (4, 0), then</p> <p>(A) $a = 2b$ (B) $2a = b$ (C) $a = b$ (D) $3a = b$</p> <div>Ans: (A) $a = 2b$</div>	1
<p>5. If A and B are two independent events with $P(A) = \frac{1}{3}$ and $P(B) = \frac{1}{4}$, then $P(B' A)$ is equal to</p> <p>(A) $\frac{1}{4}$ (B) $\frac{1}{3}$ (C) $\frac{3}{4}$ (D) 1</p> <div>Ans: (C) $\frac{3}{4}$</div>	1
<p>6. If A is a square matrix such that $A^2 = A$, then $(I - A)^3 + A$ is equal to</p> <p>(A) I (B) 0 (C) I - A (D) I + A</p> <div>Ans: (A) I</div>	1

7. $\int_1^e \frac{\log x}{x} dx$, is equal to

- (A) $\frac{e^2}{2}$ (B) 1 (C) $\frac{1}{2}$ (D) $-\infty$

Ans: (C) $\frac{1}{2}$

1

8. A point P lies on the line segment joining the points $(-1, 3, 2)$ and $(5, 0, 6)$. If x-coordinate of P is 2, then its z-coordinate is

- (A) -1 (B) 4 (C) $\frac{3}{2}$ (D) 8

Ans: (B) 4

1

9. If the projection of $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}$ on $\vec{b} = 2\hat{i} + \lambda\hat{k}$ is zero, then the value of λ is

- (A) 0 (B) 1 (C) $-\frac{2}{3}$ (D) $-\frac{3}{2}$

Ans: (C) $-\frac{2}{3}$

1

10. The vector equation of the line passing through the point $(-1, 5, 4)$ and perpendicular to the plane $z = 0$ is

- (A) $\vec{r} = -\hat{i} + 5\hat{j} + 4\hat{k} + \lambda(\hat{i} + \hat{j})$ (B) $\vec{r} = -\hat{i} + 5\hat{j} + (4 + \lambda)\hat{k}$
(C) $\vec{r} = \hat{i} - 5\hat{j} - 4\hat{k} + \lambda\hat{k}$ (D) $\vec{r} = \lambda\hat{k}$

Ans: (B) $\vec{r} = -\hat{i} + 5\hat{j} + (4 + \lambda)\hat{k}$

1

Fill in the blanks in question numbers 11 to 15

11. The position vectors of two points A and B are $\vec{OA} = 2\hat{i} - \hat{j} - \hat{k}$ and

$\vec{OB} = 2\hat{i} - \hat{j} + 2\hat{k}$, respectively. The position vector of a point P which divides the line segment joining A and B in the ratio 2 : 1 is _____.

Ans: $2\hat{i} - \hat{j} + \hat{k}$

1

12. The equation of the normal to the curve $y^2 = 8x$ at the origin is _____.

Ans: $y = 0$

1

OR

The radius of a circle is increasing at the uniform rate of 3 cm/sec. At the instant when the radius of the circle is 2 cm, its area increases at the rate of _____ cm^2/s .

Ans: 12π

1

13. If A is a square matrix of order 3 and A_{ij} is the cofactor of the element a_{ij} , then value of $a_{21} A_{11} + a_{22} A_{12} + a_{23} A_{13}$ is _____.

Ans: 0

1

OR

If the matrix A is both symmetric and skew symmetric, then A is a _____.

Ans: Zero matrix

1

14. A relation R in a set A is called _____, if $(a_1, a_2) \in R$ implies $(a_2, a_1) \in R$, for all $a_1, a_2 \in A$.

Ans: Symmetric

1

15. The greatest integer function defined by $f(x) = [x]$, $0 < x < 2$ is not differentiable at $x = \underline{\hspace{1cm}}$.

Ans: 1

1

Question numbers 16 to 20 are very short answer type questions

16. If A is a square matrix of order 3 and $|A| = 2$, then find the value of $|-AA'|$.

$$\begin{aligned} \text{Ans: } |-AA'| &= -|A|^2 \\ &= -4 \end{aligned}$$

1/2

1/2

17. Two cards are drawn at random and one-by-one without replacement from a well-shuffled pack of 52 playing cards. Find the probability that one card is red and the other is black.

$$\text{Ans: } \frac{{}^{26}C_1 \times {}^{26}C_1}{{}^{52}C_2} = \frac{26}{51}$$

1/2+1/2

18. Evaluate: $\int_1^3 |2x - 1| dx$.

$$\begin{aligned} \text{Ans: } \int_1^3 |2x - 1| dx &= \int_1^3 (2x - 1) dx = \left[\frac{1}{4} (2x - 1)^2 \right]_1^3 \\ &= 6 \end{aligned}$$

1/2

1/2

19. Find : $\int \frac{dx}{\sqrt{9 - 4x^2}}$

$$\begin{aligned} \text{Ans: } \int \frac{dx}{\sqrt{9 - 4x^2}} &= \int \frac{dx}{\sqrt{3^2 - (2x)^2}} \\ &= \frac{1}{2} \sin^{-1} \left(\frac{2x}{3} \right) + C \end{aligned}$$

1/2

1/2

20. Find: $\int x^4 \log x \, dx$.

Ans: $\int x^4 \cdot \log x \, dx = \log x \cdot \frac{x^5}{5} - \int \frac{1}{x} \cdot \frac{x^5}{5} \, dx$ 1/2

$$= \frac{x^5 \cdot \log x}{5} - \frac{x^5}{25} + c$$
 1/2

OR

Find: $\int \frac{2x}{\sqrt[3]{x^2+1}} \, dx$.

Ans: Let, $x^2 + 1 = t$

$\therefore 2x \, dx = dt$ 1/2

$$\int \frac{2x}{\sqrt[3]{x^2+1}} \, dx = \int \frac{1}{\sqrt[3]{t}} \, dt = \int t^{-1/3} \, dt = \frac{3}{2} t^{2/3} + c$$

$$= \frac{3}{2} (x^2 + 1)^{2/3} + c$$
 1/2

SECTION-B

Question numbers 21 to 26 carry 2 marks each.

21. Find a unit vector perpendicular to each of the vectors \vec{a} and \vec{b}

where $\vec{a} = 5\hat{i} + 6\hat{j} - 2\hat{k}$ and $\vec{b} = 7\hat{i} + 6\hat{j} + 2\hat{k}$.

Ans: $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 6 & -2 \\ 7 & 6 & 2 \end{vmatrix} = 24\hat{i} - 24\hat{j} - 12\hat{k}$ 1

Unit vector perpendicular to both \vec{a} and \vec{b} is $\frac{2}{3}\hat{i} - \frac{2}{3}\hat{j} - \frac{1}{3}\hat{k}$ 1

OR

Find the volume of the parallelepiped whose adjacent edges are represented by

$2\vec{a}$, $-\vec{b}$ and $3\vec{c}$, where $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{b} = 3\hat{i} + 4\hat{j} - 5\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 3\hat{k}$

Ans: Volume of the parallelepiped = $\begin{vmatrix} 2 & -2 & 4 \\ -3 & -4 & 5 \\ 6 & -3 & 9 \end{vmatrix}$ 1

$$= |-24| = 24$$
 1

22. If $f(x) = \sqrt{\tan \sqrt{x}}$, then find $f' \left(\frac{\pi^2}{16} \right)$.

Ans: $f'(x) = \frac{\sec^2 \sqrt{x}}{4\sqrt{x}\sqrt{\tan(\sqrt{x})}}$ 1

$f' \left(\frac{\pi^2}{16} \right) = \frac{2}{4 \cdot \frac{\pi}{4}} = \frac{2}{\pi}$ 1

23. Using differentials, find the approximate value of $\sqrt{25 \cdot 3}$ up to two places of decimals.

Ans: Let $y = f(x) = \sqrt{x}$, Let $x = 25$, $x + \Delta x = 25.3$, $\Delta x = 0.3$ 1

$\Delta y \simeq \left. \frac{dy}{dx} \right|_{x=25} \cdot \Delta x = \frac{1}{2\sqrt{25}} (0.3) = 0.03$ 1/2

$\sqrt{25.3} = f(25) + \Delta y = 5 + 0.03 = 5.03$ (approx.) 1/2

24. The probability of finding a green signal on a busy crossing X is 30%. What is the probability of finding a green signal on X on two consecutive days out of three?

Ans: Probability of green signal on crossing X = $\frac{30}{100} = \frac{3}{10}$ 1
 Probability of not a green signal on crossing X = $1 - \frac{3}{10} = \frac{7}{10}$

Probability of a green signal on X on two consecutive days out of three

$= \frac{3}{10} \times \frac{3}{10} \times \frac{7}{10} + \frac{7}{10} \times \frac{3}{10} \times \frac{3}{10} = \frac{63}{500}$ 1

25. Prove that $\sin^{-1} \left(2x\sqrt{1-x^2} \right) = 2 \cos^{-1} x$, $\frac{1}{\sqrt{2}} \leq x \leq 1$

Ans: Put $x = \cos \theta \Leftrightarrow \theta = \cos^{-1} x$ 1/2

L.H.S. = $\sin^{-1} \left(2x\sqrt{1-x^2} \right)$

$= \sin^{-1} (2 \cos \theta \sin \theta) = \sin^{-1} (\sin 2\theta) = 2\theta = 2 \cos^{-1} x = \text{R.H.S.}$ 1 $\frac{1}{2}$

OR

Consider a bijective function $f: \mathbb{R}_+ \rightarrow (7, \infty)$ given by $f(x) = 16x^2 + 24x + 7$, where \mathbb{R}_+ is the set of all positive real numbers. Find the inverse function of f .

Ans: Let $y = f(x) = 16x^2 + 24x + 7 = (4x + 3)^2 - 2$

$$\Rightarrow f^{-1}(y) = x = \frac{\sqrt{y+2}-3}{4}$$

26. Find the value of k so that the lines $x = -y = kz$ and $x - 2 = 2y + 1 = -z + 1$ are perpendicular to each other.

Ans: The lines, $\frac{x}{1} = \frac{y}{-1} = \frac{z}{\frac{1}{k}}$ and $\frac{x-2}{1} = \frac{y+\frac{1}{2}}{\frac{1}{2}} = \frac{z-1}{-1}$

are perpendicular $\therefore 1 - \frac{1}{2} - \frac{1}{k} = 0 \Rightarrow k = 2$

SECTION-C

Question numbers 27 to 32 carry 4 marks each.

27. A furniture trader deals in only two items – chairs and tables. He has ₹ 50,000 to invest and a space to store at most 35 items. A chair costs him ₹ 1000 and a table costs him ₹ 2000. The trader earns a profit of ₹ 150 and ₹ 250 on a chair and table, respectively. Formulate the above problem as an LPP to maximise the profit and solve it graphically.

Ans.

Let No. of chairs = x , No. of tables = y

Then L.P. P. is:


Maximize (Profit) : $Z = 150x + 250y$

Subject to : $x + y \leq 35$

$1000x + 2000y \leq 50000 \Rightarrow x + 2y \leq 50$

$x, y \geq 0$

Correct graph



Corner: Value of Z

$A(0, 25)$ ₹ 6250

$B(20, 15)$ ₹ 6750 (Max)

$C(35, 0)$ ₹ 5250

$\therefore \text{Max}(Z) = ₹ 6750$

Number of chairs = 20, Tables = 15

28. If $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$, $a > 0$, then find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{3}$.

Ans. $\frac{dy}{d\theta} = a \sin \theta$, $\frac{dx}{d\theta} = a(1 - \cos \theta)$

1/2+1/2

$$\frac{dy}{dx} = \frac{\sin \theta}{1 - \cos \theta} = \frac{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \sin^2 \frac{\theta}{2}} = \cot \frac{\theta}{2}$$

1

$$\frac{d^2y}{dx^2} = -\frac{1}{2} \cdot \operatorname{cosec}^2 \frac{\theta}{2} \cdot \frac{d\theta}{dx} = -\frac{\operatorname{cosec}^2 \frac{\theta}{2}}{2a(1 - \cos \theta)}$$

1

$$\left. \frac{d^2y}{dx^2} \right]_{\theta=\frac{\pi}{3}} = -\frac{1}{2} \times \frac{4}{a \left(1 - \frac{1}{2}\right)} = -\frac{4}{a}$$

1

29. Evaluate $\int_2^3 e^x dx$ as limit of the sums.

Ans. Let $f(x) = e^x$, $a = 1$, $b = 3$, $nh = 2$,

1

$$f(1) + f(1+h) + f(1+2h) + \dots + f(1+(n-1)h)$$

$$= e + e^{1+h} + e^{1+2h} + \dots + e^{1+(n-1)h} = \frac{e(e^{nh} - 1)}{e^h - 1}$$

2

$$\int_1^3 e^x dx = \lim_{h \rightarrow 0} h \cdot \frac{e(e^{nh} - 1)}{e^h - 1} = e(e^2 - 1) \text{ or } e^3 - e$$

1

30. There are two bags, I and II. Bag I contains 3 red and 5 black balls and Bag II contains 4 red and 3 black balls. One ball is transferred randomly from Bag I to Bag II and then a ball is drawn randomly from Bag II. If the ball so drawn is found to be black in colour, then find the probability that the transferred ball is also black.

Ans. $E_1 = \text{Event that the ball transferred from Bag I is Black}$
 $E_2 = \text{Event that the ball transferred from Bag I is Red}$
 $A = \text{Event that the ball drawn from Bag II is Black}$

1/2

$$P(E_1) = \frac{5}{8}; P(E_2) = \frac{3}{8}; P\left(\frac{A}{E_1}\right) = \frac{4}{8} = \frac{1}{2}; P\left(\frac{A}{E_2}\right) = \frac{3}{8}$$

2

Required Probability:

$$P\left(\frac{E_1}{A}\right) = \frac{P(E_1) \cdot P\left(\frac{A}{E_1}\right)}{P(E_1) \cdot P\left(\frac{A}{E_1}\right) + P(E_2) \cdot P\left(\frac{A}{E_2}\right)} = \frac{\frac{5}{8} \cdot \frac{1}{2}}{\frac{5}{8} \cdot \frac{1}{2} + \frac{3}{8} \cdot \frac{3}{8}} = \frac{20}{29} \quad 1\frac{1}{2}$$

OR

An urn contains 5 red, 2 white and 3 black balls. Three balls are drawn, one-by-one, at random without replacement. Find the probability distribution of the number of white balls. Also, find the mean and the variance of the number of white balls drawn.

Ans. Let X = No. of white balls = 0, 1, 2

$$X : \quad \quad \quad 0 \quad \quad \quad 1 \quad \quad \quad 2 \quad \quad \quad 1/2$$

$$P(X) : \quad \frac{8}{10} \times \frac{7}{9} \times \frac{6}{8} = \frac{7}{15} \quad 3 \times \frac{8}{10} \times \frac{7}{9} \times \frac{2}{8} = \frac{7}{15} \quad 3 \times \frac{2}{10} \times \frac{1}{9} \times \frac{8}{8} = \frac{1}{15} \quad 1\frac{1}{2}$$

$$X \cdot P(X) : \quad \quad \quad 0 \quad \quad \quad \frac{7}{15} \quad \quad \quad \frac{2}{15} \quad \quad \quad 1/2$$

$$X^2 P(X) : \quad \quad \quad 0 \quad \quad \quad \frac{7}{15} \quad \quad \quad \frac{4}{15}$$

$$\text{Mean} = \sum X P(X) = \frac{9}{15} = \frac{3}{5} \quad 1/2$$

$$\text{Variance} = \sum X^2 P(x) - \left[\sum X P(X)^2 \right] = \frac{11}{15} - \left[\frac{3}{5} \right]^2 = \frac{28}{75} \quad 1$$

31. Find the general solution of the differential equation $ye^y dx = (y^3 + 2xe^y) dy$.

$$\text{Ans.} \quad y \cdot e^y dx = (y^3 + 2xe^y) dy \Rightarrow y \cdot e^y \frac{dy}{dx} = y^3 + 2xe^y$$

$$\therefore \frac{dx}{dy} - \frac{2}{y}x = y^2 \cdot e^{-y} \quad 1$$

$$\text{I.F. (Integrating factor)} = e^{-2 \int \frac{1}{y} dy} = e^{-2 \log y} = e^{\log \frac{1}{y^2}} = \frac{1}{y^2} \quad 1$$

\therefore Solution is

$$x \cdot \frac{1}{y^2} = \int y^2 \cdot e^{-y} \cdot \frac{1}{y^2} dy + c = \int e^{-y} dy + c \quad 1$$

$$\Rightarrow \frac{x}{y^2} = -e^{-y} + c \quad \text{or} \quad x = -y^2 e^{-y} + cy^2 \quad 1$$

OR

Find the particular solution of the differential equation

$$x \frac{dy}{dx} = y - x \tan\left(\frac{y}{x}\right), \text{ given that } y = \frac{\pi}{4} \text{ at } x = 1.$$

Ans. The differential equation can be written as:

$$\frac{dy}{dx} = \frac{y}{x} - \tan\frac{y}{x}, \text{ let } y = vx \therefore \frac{dy}{dx} = v + x \frac{dv}{dx} \quad 1$$

$$\Rightarrow v + x \frac{dv}{dx} = v - \tan v \Rightarrow \cot v \, dv = -\frac{1}{x} dx$$

Integrate both sides

$$\log \sin v = -\log |x| + \log c \Rightarrow \log \sin \frac{y}{x} = \log \frac{c}{x} \quad 2$$

$$\Rightarrow x \cdot \sin \frac{y}{x} = c, \text{ Put } y = \frac{\pi}{4} \text{ and } x = 1$$

$$\Rightarrow \sin \frac{\pi}{4} = c \text{ or } c = \frac{1}{\sqrt{2}} \quad 1/2$$

$$\therefore \text{ Particular solution is } x \cdot \sin\left(\frac{y}{x}\right) = \frac{1}{\sqrt{2}} \quad 1/2$$

32. Let N be the set of natural numbers and R be the relation on $N \times N$ defined by $(a, b) R (c, d)$ iff $ad = bc$ for all $a, b, c, d \in N$. Show that R is an equivalence relation.

Ans: Reflexive: For any $(a, b) \in N \times N$

$$a \cdot b = b \cdot a$$

$$\therefore (a, b) R (a, b) \text{ thus } R \text{ is reflexive} \quad 1$$

Symmetric: For $(a, b), (c, d) \in N \times N$

$$(a, b) R (c, d) \Rightarrow a \cdot d = b \cdot c$$

$$\Rightarrow c \cdot b = d \cdot a$$

$$\Rightarrow (c, d) R (a, b) \therefore R \text{ is symmetric} \quad 1\frac{1}{2}$$

Transitive : For any $(a, b), (c, d), (e, f), \in N \times N$

$$(a, b) R (c, d) \text{ and } (c, d) R (e, f)$$

$$\Rightarrow a \cdot d = b \cdot c \text{ and } c \cdot f = d \cdot e$$

$$\Rightarrow a \cdot d \cdot c \cdot f = b \cdot c \cdot d \cdot e \Rightarrow a \cdot f = b \cdot e$$

$$\therefore (a, b) R (e, f), \therefore R \text{ is transitive} \quad 1\frac{1}{2}$$


$\therefore R$ is an equivalence Relation

SECTION-D

Question numbers 33 to 36 carry 6 marks each.

33. Show that the height of the right circular cylinder of greatest volume which can be inscribed in a right circular cone of height h and radius r is one-third of the height of the cone, and the greatest volume of the cylinder is $\frac{4}{9}$ times the volume of the cone.

Ans.



Let H = Height of cylinder

R = Radius of cylinder

$$\text{Volume of cone} = \frac{\pi}{3} r^2 h \quad 1/2$$

$$V = \text{Volume of cylinder} = \pi R^2 H \quad 1/2$$

$$\Delta ADF \sim \Delta AEC \Rightarrow \frac{h-H}{h} = \frac{R}{r} \Rightarrow R = \frac{r}{h}(h-H) \quad 1$$

$$\therefore V = \pi \cdot H \cdot \frac{r^2}{h^2} (h-H)^2 = \frac{\pi r^2}{h^2} (H^3 - 2hH^2 + Hh^2) \quad 1$$

$$V'(H) = \frac{\pi r^2}{h^2} (3H^2 - 4hH + h^2), \quad V'(h) = 0 \Rightarrow H = \frac{h}{3} \quad 1+1$$


$$V''(H) = \frac{\pi r^2}{h^2} (6H - 4h), \quad V''\left(H = \frac{h}{3}\right) = \frac{\pi r^2}{h^2} (-2h) < 0 \quad 1/2$$

$$\therefore V \text{ is max iff } H = \frac{h}{3} \text{ and } R = \frac{2r}{3}$$

$$\frac{\text{Volume of cylinder}}{\text{Volume of cone}} = \frac{3\pi R^2 H}{\pi r^2 h} = 3 \cdot \frac{4r^2}{9} \cdot \frac{h}{3} \cdot \frac{1}{r^2 h} = \frac{4}{9} \quad 1/2$$

34. Using integration, find the area of the region enclosed by the parabola $y = 3x^2$ and the line $3x - y + 6 = 0$.

Ans.



$$\text{Points of intersection } x = -1, 2 \quad 1$$

$$\text{Correct Graph} \quad 1$$

Required area

$$= \int_{-1}^2 3(x+2) dx - 3 \int_{-1}^2 x^2 dx \quad 2$$

$$= \frac{3}{2} \left[(x+2)^2 \right]_{-1}^2 - \left[x^3 \right]_{-1}^2 \quad 1 \frac{1}{2}$$

$$= \frac{3}{2} \times 15 - 9 = \frac{27}{2} \quad 1/2$$

35. Find the equation of the plane that contains the point A(2,1,-1) and is perpendicular to the line of intersection of the planes $2x + y - z = 3$ and $x + 2y + z = 2$. Also find the angle between the plane thus obtained and the y-axis.

Ans. Let equation of the required plane be:

$$a(x - 2) + b(y - 1) + c(z + 1) = 0 \quad 1\frac{1}{2}$$

$$\text{Also : } 2a + b - c = 0$$

$$a + 2b + c = 0$$

$$\text{Solving: } \frac{a}{3} = \frac{b}{-3} = \frac{c}{3} = k \Rightarrow a = 3k, b = -3k, c = 3k \quad 1\frac{1}{2}$$

$$\therefore \text{Equation of plane is : } 3k(x - 2) - 3k(y - 1) + 3k(z + 1) = 0$$

$$\Rightarrow x - y + z = 0 \quad 1\frac{1}{2}$$

Let angle between y-axis and plane = θ

$$\text{then, } \sin \theta = \left| \frac{0 - 1 + 0}{\sqrt{1 + 1 + 1}} \right| = \left| \frac{-1}{\sqrt{3}} \right| \Rightarrow \theta = \sin^{-1} \left(\frac{1}{\sqrt{3}} \right) \quad 1\frac{1}{2}$$

OR

Find the distance of the point P(-2, -4, 7) from the point of intersection Q of the line $\vec{r} = (3\hat{i} - 2\hat{j} + 6\hat{k}) + \lambda(2\hat{i} - \hat{j} + 2\hat{k})$ and the plane $\vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 6$. Also write the vector equation of the line PQ.

$$\text{Ans. General point on line is: } \vec{r} = (3 + 2\lambda)\hat{i} + (-2 - \lambda)\hat{j} + (6 + 2\lambda)\hat{k} \quad 1$$

For the point of intersection:

$$\left[(3 + 2\lambda)\hat{i} + (-2 - \lambda)\hat{j} + (6 + 2\lambda)\hat{k} \right] \cdot (\hat{i} - \hat{j} + \hat{k}) = 6 \quad 1$$

$$\Rightarrow 3 + 2\lambda + 2 + \lambda + 6 + 2\lambda = 6 \Rightarrow \lambda = -1 \quad 1$$

$$\therefore Q(\hat{i} - \hat{j} + 4\hat{k}) = Q(1, -1, 4) \quad 1$$

$$PQ = 3\sqrt{3}, \text{ equation of the line PQ : } \vec{r} = -2\hat{i} - 4\hat{j} + 7\hat{k} + \mu(3\hat{i} + 3\hat{j} - 3\hat{k}) \quad 1+1$$

36. If $A = \begin{bmatrix} 1 & 2 & -3 \\ 3 & 2 & -2 \\ 2 & -1 & 1 \end{bmatrix}$, then find A^{-1} and use it to solve the

following system of the equations:

$$x + 2y - 3z = 6$$

$$3x + 2y - 2z = 3$$

$$2x - y + z = 2$$

$$\text{Ans. } |A| = 7; \text{adj}(A) = \begin{bmatrix} 0 & 1 & 2 \\ -7 & 7 & -7 \\ -7 & 5 & -4 \end{bmatrix}; A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{7} \begin{bmatrix} 0 & 1 & 2 \\ -7 & 7 & -7 \\ -7 & 5 & -4 \end{bmatrix} \quad 1 + 1 \frac{1}{2} + \frac{1}{2}$$

The system of equations in Matrix form can be written as :

$$A \cdot X = B, \text{ where } X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}; B = \begin{bmatrix} 6 \\ 3 \\ 2 \end{bmatrix} \quad 1$$

$$X = A^{-1}B \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 0 & 1 & 2 \\ -7 & 7 & -7 \\ -7 & 5 & -4 \end{bmatrix} \begin{bmatrix} 6 \\ 3 \\ 2 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 7 \\ -35 \\ -35 \end{bmatrix} = \begin{bmatrix} 1 \\ -5 \\ -5 \end{bmatrix} \quad 1$$

$$\therefore x = 1, y = -5, z = -5 \quad 1$$

OR

Using properties of determinants, prove that

$$\begin{vmatrix} (b+c)^2 & a^2 & bc \\ (c+a)^2 & b^2 & ca \\ (a+b)^2 & c^2 & ab \end{vmatrix} = (a-b)(b-c)(c-a)(a+b+c)(a^2+b^2+c^2)$$

$$\text{Ans. } \begin{vmatrix} (b+c)^2 & a^2 & bc \\ (c+a)^2 & b^2 & ca \\ (a+b)^2 & c^2 & ab \end{vmatrix}$$

$$= \begin{vmatrix} b^2+c^2 & a^2 & bc \\ c^2+a^2 & b^2 & ca \\ a^2+b^2 & c^2 & ab \end{vmatrix} \quad (C_1 \rightarrow C_1 - 2C_3) \quad 1$$

$$= \begin{vmatrix} a^2+b^2+c^2 & a^2 & bc \\ a^2+b^2+c^2 & b^2 & ca \\ a^2+b^2+c^2 & c^2 & ab \end{vmatrix} \quad (C_1 \rightarrow C_1 + C_2) \quad 1$$

$$= \begin{vmatrix} a^2+b^2+c^2 & a^2 & bc \\ 0 & b^2-a^2 & ca-bc \\ 0 & c^2-a^2 & ab-bc \end{vmatrix} \quad (R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1) \quad 2$$

$$= (b-a)(c-a) \begin{vmatrix} a^2+b^2+c^2 & a^2 & bc \\ 0 & b+a & -c \\ 0 & c+a & -b \end{vmatrix} \quad 1$$

Expand along C_1

$$= (a^2 + b^2 + c^2)(b-a)(c-a)(-b^2 - ab + c^2 + ac)$$

$$= (a-b)(b-c)(c-a)(a+b+c)(a^2 + b^2 + c^2)$$

1
