1

EE5609 Assignment 1

Vimal K B Roll No - AI20MTECH14002

Abstract—This assignment involves finding a vector which is perpendicular to given two vectors and nonperpendicular to a third vector.

The python solution code for this problem can be downloaded from

> https://github.com/vimalkb007/EE5609/ blob/master/Assignment 1/codes/ assignment1 solution.py

The python verification code for this problem can be downloaded from

> https://github.com/vimalkb007/EE5609/ blob/master/Assignment 1/codes/ assignment1 solution verify.py

1 PROBLEM STATEMENT

Let $\mathbf{a} = \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 3 \\ -2 \\ 7 \end{pmatrix}$ and $\mathbf{c} = \begin{pmatrix} 2 \\ -1 \\ 4 \end{pmatrix}$. Find a

vector **d** such that $\mathbf{d} \perp \mathbf{a}, \mathbf{d} \perp \mathbf{b}$ and $\mathbf{d}^T \mathbf{c} = 15$.

2 Theory

If two vectors are perpendicular, then their dot product is 0. If we have two vectors x, y is given by $\mathbf{x} \cdot \mathbf{v} = |\mathbf{x}| |\mathbf{v}| \cos(\theta)$.

When $\theta = \pi/2$ (90°), then $\cos \theta = 0 \implies \mathbf{x} \cdot \mathbf{y}$

If we have 3 equations and 3 unknowns, we can use Guassian Elimination method in order to find the unknowns.

3 Solution

Lets consider vector **d** as $\begin{bmatrix} d_2 \\ d_3 \end{bmatrix}$

It is given that $\mathbf{d} \perp \mathbf{a}$, then their correponding dot product will be 0.

$$\mathbf{d}^T \mathbf{a} = 0 \implies \begin{pmatrix} d_1 \\ d_2 \\ d_3 \end{pmatrix}^T \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix} = 0$$

$$d_1 + 4d_2 + 2d_3 = 0 (3.0.1)$$

Similarly, as $\mathbf{d} \perp \mathbf{b}$,

$$\mathbf{d}^{T}\mathbf{b} = 0 \implies \begin{pmatrix} d_{1} \\ d_{2} \\ d_{3} \end{pmatrix}^{T} \begin{pmatrix} 3 \\ -2 \\ 7 \end{pmatrix} = 0$$

$$3d_{1} - 2d_{2} + 7d_{3} = 0 \tag{3.0.2}$$

Since, it is given that $\mathbf{d}^T \mathbf{c} = 15$, we can write it as $(d_1 \ d_2 \ d_3) \begin{pmatrix} 2 \\ -1 \\ A \end{pmatrix} = 15.$

$$2d_1 - d_2 + 4d_3 = 15 \tag{3.0.3}$$

Using equations 3.0.1, 3.0.2, 3.0.3, we can use Guassian Elimination Method in order to find the values of d_1 , d_2 , d_3 .

$$\begin{pmatrix}
1 & 4 & 2 & 0 \\
3 & -2 & 7 & 0 \\
2 & -1 & 4 & 15
\end{pmatrix}$$
(3.0.4)

$$\begin{pmatrix}
1 & 4 & 2 & 0 \\
3 & -2 & 7 & 0 \\
2 & -1 & 4 & 15
\end{pmatrix}$$

$$\xrightarrow{R_3 \leftarrow R_3 - 2R_1} \begin{pmatrix}
1 & 4 & 2 & 0 \\
0 & -14 & 1 & 0 \\
0 & -9 & 0 & 15
\end{pmatrix}$$
(3.0.4)

$$\stackrel{R_3 \leftarrow R_3 - \frac{9}{14}R_2}{\longleftrightarrow} \begin{pmatrix} 1 & 4 & 2 & 0 \\ 0 & -14 & 1 & 0 \\ 0 & 0 & \frac{-9}{14} & 15 \end{pmatrix}$$
(3.0.6)

$$\begin{array}{c|cccc}
(0 & 0 & \overline{14} & | & 13) \\
\hline
R_3 \leftarrow \frac{-14}{9} R_2 \\
\hline
R_2 \leftarrow \frac{-1}{14} R_2
\end{array}$$

$$\begin{array}{c|cccc}
(1 & 4 & 2 & | & 0 \\
0 & 1 & \frac{-1}{14} & | & 0 \\
0 & 0 & 1 & | & \frac{-210}{9}
\end{array}$$

$$(3.0.7)$$

$$\stackrel{R_1 \leftarrow R_1 + \frac{1}{14}R_3}{\longleftrightarrow} \begin{pmatrix} 1 & 4 & 2 & 0 \\ 0 & 1 & 0 & \frac{-210}{126} \\ 0 & 0 & 1 & \frac{-210}{9} \end{pmatrix}$$
(3.0.8)

$$\stackrel{R_1 \leftarrow R_1 - 4R_3}{\longleftrightarrow} \begin{pmatrix}
1 & 0 & 2 & | & \frac{840}{126} \\
0 & 1 & 0 & | & \frac{-210}{126} \\
0 & 0 & 1 & | & \frac{-210}{9}
\end{pmatrix}$$

$$\stackrel{R_1 \leftarrow R_1 - 2R_3}{\longleftrightarrow} \begin{pmatrix}
1 & 0 & 0 & | & \frac{6720}{126} \\
0 & 1 & 0 & | & \frac{-210}{126} \\
0 & 0 & 1 & | & \frac{-210}{9}
\end{pmatrix}$$
(3.0.9)

By using Guassian Elimination Method, we were able to get the vector
$$\mathbf{d}$$
 as $\begin{pmatrix} \frac{6720}{126} \\ \frac{-210}{126} \\ \frac{-210}{9} \end{pmatrix}$

The resultant vector
$$\mathbf{d} = \begin{pmatrix} 53.333 \\ -1.667 \\ -23.333 \end{pmatrix}$$