

EE5609 Assignment 1

Vimal K B

Roll No - AI20MTECH14002

Abstract—This assignment involves finding a vector which is perpendicular to given two vectors and non-perpendicular to a third vector.

The python solution code for this problem can be downloaded from

https://github.com/vimalkb007/EE5609/blob/master/Assignment_1/codes/assignment1_solution.py

The python verification code for this problem can be downloaded from

https://github.com/vimalkb007/EE5609/blob/master/Assignment_1/codes/assignment1_solution_verify.py

$$\mathbf{d}^T \mathbf{b} = 0 \quad (3.0.2)$$

It is given that

$$\mathbf{d}^T \mathbf{c} = 15 \quad (3.0.3)$$

Using equations 3.0.1, 3.0.2, 3.0.3, we can represent them in a Matrix Representation of Linear Equations $Ax=B$ form as:

$$\begin{pmatrix} \mathbf{a}^T \\ \mathbf{b}^T \\ \mathbf{c}^T \end{pmatrix} \mathbf{d} = \begin{pmatrix} 0 \\ 0 \\ 15 \end{pmatrix} \quad (3.0.4)$$

Numerically, using \mathbf{a} , \mathbf{b} , \mathbf{c} the above equation 3.0.4 can be written as,

$$\begin{pmatrix} 1 & 4 & 2 \\ 3 & -2 & 7 \\ 2 & -1 & 4 \end{pmatrix} \mathbf{d} = \begin{pmatrix} 0 \\ 0 \\ 15 \end{pmatrix} \quad (3.0.5)$$

we can use Guassian Elimination Method in order to find the coordinate values of \mathbf{d} .

1 PROBLEM STATEMENT

Let $\mathbf{a} = \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 3 \\ -2 \\ 7 \end{pmatrix}$ and $\mathbf{c} = \begin{pmatrix} 2 \\ -1 \\ 4 \end{pmatrix}$. Find a vector \mathbf{d} such that $\mathbf{d} \perp \mathbf{a}$, $\mathbf{d} \perp \mathbf{b}$ and $\mathbf{d}^T \mathbf{c} = 15$.

2 THEORY

If two vectors are perpendicular, then their dot product is 0. If we have two vectors \mathbf{x} , \mathbf{y} is given by $\mathbf{x} \cdot \mathbf{y} = |\mathbf{x}||\mathbf{y}|\cos(\theta)$.

When $\theta = \pi/2$ (90°), then $\cos \theta = 0 \implies \mathbf{x} \cdot \mathbf{y} = 0$.

If we have 3 equations and 3 unknowns, we can use Guassian Elimination method in order to find the unknowns.

3 SOLUTION

It is given that $\mathbf{d} \perp \mathbf{a}$, then their corresponding dot product will be 0.

$$\mathbf{d}^T \mathbf{a} = 0 \quad (3.0.1)$$

Similarly, as $\mathbf{d} \perp \mathbf{b}$

$$\begin{pmatrix} 1 & 4 & 2 \\ 3 & -2 & 7 \\ 2 & -1 & 4 \end{pmatrix} \begin{array}{l} 0 \\ 0 \\ 15 \end{array} \quad (3.0.6)$$

$$\begin{array}{l} \xleftarrow{R_3 \leftarrow R_3 - 2R_1} \\ \xleftarrow{R_2 \leftarrow R_2 - 3R_1} \end{array} \begin{pmatrix} 1 & 4 & 2 \\ 0 & -14 & 1 \\ 0 & -9 & 0 \end{pmatrix} \begin{array}{l} 0 \\ 0 \\ 15 \end{array} \quad (3.0.7)$$

$$\xleftarrow{R_3 \leftarrow R_3 - \frac{9}{14}R_2} \begin{pmatrix} 1 & 4 & 2 \\ 0 & -14 & 1 \\ 0 & 0 & \frac{-9}{14} \end{pmatrix} \begin{array}{l} 0 \\ 0 \\ 15 \end{array} \quad (3.0.8)$$

$$\begin{array}{l} \xleftarrow{R_3 \leftarrow \frac{-14}{9}R_3} \\ \xleftarrow{R_2 \leftarrow \frac{-1}{14}R_2} \end{array} \begin{pmatrix} 1 & 4 & 2 \\ 0 & 1 & \frac{-1}{14} \\ 0 & 0 & 1 \end{pmatrix} \begin{array}{l} 0 \\ 0 \\ \frac{-210}{9} \end{array} \quad (3.0.9)$$

$$\xleftarrow{R_1 \leftarrow R_1 + \frac{1}{14}R_2} \begin{pmatrix} 1 & 4 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{array}{l} 0 \\ \frac{-210}{126} \\ \frac{-210}{9} \end{array} \quad (3.0.10)$$

$$\xleftrightarrow{R_1 \leftarrow R_1 - 4R_3} \left(\begin{array}{ccc|c} 1 & 0 & 2 & \frac{840}{126} \\ 0 & 1 & 0 & \frac{-210}{126} \\ 0 & 0 & 1 & \frac{-210}{9} \end{array} \right) \quad (3.0.11)$$

$$\xleftrightarrow{R_1 \leftarrow R_1 - 2R_3} \left(\begin{array}{ccc|c} 1 & 0 & 0 & \frac{6720}{126} \\ 0 & 1 & 0 & \frac{-210}{126} \\ 0 & 0 & 1 & \frac{-210}{9} \end{array} \right) \quad (3.0.12)$$

By using Gaussian Elimination Method, we were

able to get the vector \mathbf{d} as $\begin{pmatrix} \frac{6720}{126} \\ \frac{-210}{126} \\ \frac{-210}{9} \end{pmatrix}$

The resultant vector $\mathbf{d} = \begin{pmatrix} 53.333 \\ -1.667 \\ -23.333 \end{pmatrix}$