



Clinical Decision Support Report

Patient ID	P-001	Age	25
Gender	Male	Weight	60 kg

Diagnosis

typhoid (Confidence: 61%)

Clinical Summary

1. Explanation of Primary Diagnosis:

The primary diagnosis of typhoid fever (confidence: 61%) fits best due to the presence of key symptoms such as fever, headache, abdominal pain, and the patient's age and gender. Typhoid fever is an acute illness characterized by fever caused by infection with the bacterium *Salmonella Typhi*, which matches the patient's symptoms. The insidious onset of fever, headache, and abdominal pain, along with the possibility of malaise, fatigue, and loss of appetite, are all consistent with typhoid fever. The patient's symptoms, such as fever and abdominal pain, are also consistent with the characteristic step-ladder pattern of typhoid fever.

2. Comparison with Other Possible Diagnoses:

In comparison to other possible diagnoses, dengue (52%) and malaria (48%) are also potential causes of fever and abdominal pain. However, typhoid fever is more likely due to the combination of symptoms, such as headache, abdominal pain, and the absence of typical dengue symptoms like severe joint pain or malaria symptoms like cyclical fevers. Dengue and malaria also have distinct diagnostic features, such as thrombocytopenia in dengue or parasitemia in malaria, which are not mentioned in the patient's symptoms.

3. Urgency Level:

The urgency level for this patient is ****High****. Typhoid fever can lead to severe complications, such as intestinal perforation, hemorrhage, or sepsis, if left untreated or untreated promptly. Prompt antibiotic therapy and supportive care are essential to prevent these complications and reduce the risk of mortality.

4. Immediate Actions:

Immediate actions include:

- * Consulting a doctor for further evaluation and guidance
- * Starting antibiotic therapy as prescribed by the doctor
- * Eating high-calorie vegetables to help manage symptoms and support recovery
- * Ensuring adequate hydration and rest
- * Monitoring the patient's condition closely for any signs of complications or deterioration

5. Further Diagnostics:

Further diagnostics may include:

- * Blood cultures to detect Salmonella Typhi
- * Stool or urine tests to detect the presence of Salmonella Typhi
- * Bone marrow biopsy in some cases
- * Imaging studies, such as abdominal ultrasound or CT scan, to evaluate for complications like intestinal perforation
- * Complete blood count (CBC) to monitor for signs of infection or inflammation
- * Liver function tests to evaluate for any hepatic involvement

These diagnostic tests will help confirm the diagnosis of typhoid fever and guide further management and treatment.

Suggested Clinical Management

- 1. Ciprofloxacin - 500 mg twice daily
- 2. Ceftriaxone - 50 mg/kg/day (for 60 kg, 3000 mg/day, given in 2 divided doses: 1500 mg twice daily)
- 3. Azithromycin - 500 mg once daily

■■■ This report is generated by EXiCare AI for clinical decision support only.