

Cubic Splines

$n+1$ piece wise approx



x

y

$$1. a_i = f(x_i)$$

y

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \dots & 0 \\ h_0 & 2(h_0 + h_1) & h_1 & 0 & 0 & \dots & 0 \\ 0 & h_1 & 2(h_1 + h_2) & h_2 & 0 & \dots & 0 \\ 0 & 0 & h_2 & 2(h_2 + h_3) & h_3 & \dots & 0 \\ 0 & 0 & 0 & h_3 & 2(h_3 + h_4) & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \\ c_2 \\ c_3 \\ \vdots \\ \vdots \\ c_n \end{pmatrix} = \begin{pmatrix} 0 \\ \frac{3}{h_1}(a_2 - a_1) - \frac{3}{h_0}(a_1 - a_0) \\ \frac{3}{h_2}(a_3 - a_2) - \frac{3}{h_1}(a_2 - a_1) \\ \frac{3}{h_3}(a_4 - a_3) - \frac{3}{h_2}(a_3 - a_2) \\ \vdots \\ \vdots \\ 0 \end{pmatrix}$$

solving for d_i in (7)

$$d_i = \frac{1}{3h_i}(c_{i+1} - c_i) \quad (11)$$

now substitute d_i in (5) and solve for b_i

$$b_i = \frac{a_{i+1} - a_i}{h_i} - \frac{2c_i + c_{i+1}}{3}h_i \quad (10)$$

Simple example

Example 1 Construct a natural cubic spline that passes through the points (1, 2), (2, 3), and (3, 5).

Solution This spline consists of two cubics. The first for the interval [1, 2], denoted

$$S_0(x) = a_0 + b_0(x - 1) + c_0(x - 1)^2 + d_0(x - 1)^3,$$

Copyright 2010 Cengage Learning. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has determined that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

and the other for [2, 3], denoted

$$S_1(x) = a_1 + b_1(x - 2) + c_1(x - 2)^2 + d_1(x - 2)^3.$$

There are 8 constants to be determined, which requires 8 conditions. Four conditions come from the fact that the splines must agree with the data at the nodes. Hence

$$\begin{aligned} 2 &= f(1) = a_0, & 3 &= f(2) = a_0 + b_0 + c_0 + d_0, & 3 &= f(2) = a_1, & \text{and} \\ 5 &= f(3) = a_1 + b_1 + c_1 + d_1. \end{aligned}$$

Two more come from the fact that $S'_0(2) = S'_1(2)$ and $S''_0(2) = S''_1(2)$. These are

$$S'_0(2) = S'_1(2): \quad b_0 + 2c_0 + 3d_0 = b_1 \quad \text{and} \quad S''_0(2) = S''_1(2): \quad 2c_0 + 6d_0 = 2c_1$$

The final two come from the natural boundary conditions:

$$S''_0(1) = 0: \quad 2c_0 = 0 \quad \text{and} \quad S''_1(3) = 0: \quad 2c_1 + 6d_1 = 0.$$

Solving this system of equations gives the spline

$$S(x) = \begin{cases} 2 + \frac{3}{2}(x - 1) + \frac{1}{4}(x - 1)^3, & \text{for } x \in [1, 2] \\ 3 + \frac{3}{2}(x - 2) + \frac{1}{4}(x - 2)^2 - \frac{1}{4}(x - 2)^3, & \text{for } x \in [2, 3] \end{cases}$$

Example 2 At the beginning of Chapter 3 we gave some Taylor polynomials to approximate the exponential $f(x) = e^x$. Use the data points $(0, 1)$, $(1, e)$, $(2, e^2)$, and $(3, e^3)$ to form a natural spline $S(x)$ that approximates $f(x) = e^x$.

Solution We have $n = 3$, $h_0 = h_1 = h_2 = 1$, $a_0 = 1$, $a_1 = e$, $a_2 = e^2$, and $a_3 = e^3$. So the matrix A and the vectors \mathbf{b} and \mathbf{x} given in Theorem 3.11 have the forms

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 4 & 1 & 0 \\ 0 & 1 & 4 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 0 \\ 3(e^2 - 2e + 1) \\ 3(e^3 - 2e^2 + e) \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} c_0 \\ c_1 \\ c_2 \\ c_3 \end{bmatrix}.$$

The vector-matrix equation $A\mathbf{x} = \mathbf{b}$ is equivalent to the system of equations

$$\begin{aligned} c_0 &= 0, \\ c_0 + 4c_1 + c_2 &= 3(e^2 - 2e + 1), \\ c_1 + 4c_2 + c_3 &= 3(e^3 - 2e^2 + e), \\ c_3 &= 0. \end{aligned}$$

This system has the solution $c_0 = c_3 = 0$, and to 5 decimal places,

$$c_1 = \frac{1}{5}(-e^3 + 6e^2 - 9e + 4) \approx 0.75685, \quad \text{and} \quad c_2 = \frac{1}{5}(4e^3 - 9e^2 + 6e - 1) \approx 5.83007.$$

Solving for the remaining constants gives

$$\begin{aligned} b_0 &= \frac{1}{h_0}(a_1 - a_0) - \frac{h_0}{3}(c_1 + 2c_0) \\ &= (e - 1) - \frac{1}{15}(-e^3 + 6e^2 - 9e + 4) \approx 1.46600, \\ b_1 &= \frac{1}{h_1}(a_2 - a_1) - \frac{h_1}{3}(c_2 + 2c_1) \\ &= (e^2 - e) - \frac{1}{15}(2e^3 + 3e^2 - 12e + 7) \approx 2.22285, \\ b_2 &= \frac{1}{h_2}(a_3 - a_2) - \frac{h_2}{3}(c_3 + 2c_2) \\ &= (e^3 - e^2) - \frac{1}{15}(8e^3 - 18e^2 + 12e - 2) \approx 8.80977, \\ d_0 &= \frac{1}{3h_0}(c_1 - c_0) = \frac{1}{15}(-e^3 + 6e^2 - 9e + 4) \approx 0.25228, \\ d_1 &= \frac{1}{3h_1}(c_2 - c_1) = \frac{1}{3}(e^3 - 3e^2 + 3e - 1) \approx 1.69107, \end{aligned}$$

and

$$d_2 = \frac{1}{3h_2}(c_3 - c_1) = \frac{1}{15}(-4e^3 + 9e^2 - 6e + 1) \approx -1.94336.$$

The natural cubic spline is described piecewise by

$$S(x) = \begin{cases} 1 + 1.46600x + 0.25228x^3, & \text{for } x \in [0, 1], \\ 2.71828 + 2.22285(x - 1) + 0.75685(x - 1)^2 + 1.69107(x - 1)^3, & \text{for } x \in [1, 2], \\ 7.38906 + 8.80977(x - 2) + 5.83007(x - 2)^2 - 1.94336(x - 2)^3, & \text{for } x \in [2, 3]. \end{cases}$$

Derivation.

Graph of a natural cubic spline $S(x)$ passing through points $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$. The spline is composed of n polynomials S_0, S_1, \dots, S_n . The points are marked as nodes, and the intervals between them are marked as $h_i = x_{i+1} - x_i$.

Conditions:

- $S_i(x_i) = y_i \quad \forall i = 0, \dots, n$
- $S_i(x_{i+1}) = S_{i+1}(x_{i+1}) \quad \forall i = 0, \dots, n-1$
- $S_i'(x_{i+1}) = S_{i+1}'(x_{i+1}) \quad \forall i = 0, \dots, n-1$
- $S_i''(x_{i+1}) = S_{i+1}''(x_{i+1}) \quad \forall i = 0, \dots, n-1$
- $S_0''(x_0) = 0 \quad ; \quad S_n''(x_n) = 0$

Polynomial form:

$$S_i(x) = a_i + b_i(x - x_i) + c_i(x - x_i)^2 + d_i(x - x_i)^3$$

Unknowns: $4n$

Number of conditions: $4n + 2$

$$\begin{aligned} S_i(x) &= a_i + b_i(x - x_i) + c_i(x - x_i)^2 + d_i(x - x_i)^3 \\ S_{i+1}(x) &= a_{i+1} + b_{i+1}(x - x_{i+1}) + c_{i+1}(x - x_{i+1})^2 + d_{i+1}(x - x_{i+1})^3 \\ S_i'(x) &= b_i + 2c_i(x - x_i) + 3d_i(x - x_i)^2 \\ S_{i+1}'(x) &= b_{i+1} + 2c_{i+1}(x - x_{i+1}) + 3d_{i+1}(x - x_{i+1})^2 \\ S_i''(x) &= 2c_i + 6d_i(x - x_i) \\ S_{i+1}''(x) &= 2c_{i+1} + 6d_{i+1}(x - x_{i+1}) \end{aligned}$$

Continuity conditions at x_{i+1} :

$$\begin{aligned} S_i(x_{i+1}) &= S_{i+1}(x_{i+1}) \\ S_i'(x_{i+1}) &= S_{i+1}'(x_{i+1}) \\ S_i''(x_{i+1}) &= S_{i+1}''(x_{i+1}) \end{aligned}$$

$$\begin{aligned} ① \quad S_i(x_i) &= y_i \\ a_i &= y_i \\ ② \quad S_{i+1}(x_{i+1}) &= S_i(x_{i+1}) \\ a_{i+1} &= a_i + b_i h_i + c_i h_i^2 + d_i h_i^3 \quad [5] \\ ③ \quad S_{i+1}'(x_{i+1}) &= S_i'(x_{i+1}) \\ b_{i+1} &= b_i + 2c_i h_i + 3d_i h_i^2 \quad [6] \\ ④ \quad S_{i+1}''(x_{i+1}) &= S_i''(x_{i+1}) \\ 2c_{i+1} &= 2c_i + 6d_i h_i \\ c_{i+1} &= c_i + 3d_i h_i \quad [7] \end{aligned}$$

$$\begin{aligned} ⑤ \quad S_0''(x_0) &= 0 & S_n''(x_n) &= 0 \\ c_0 &= 0 & c_n &= 0 \end{aligned}$$

7

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

It is helpful to write the equations as follows:

$$\begin{aligned} a_0 \sum_{i=1}^m x_i^0 + a_1 \sum_{i=1}^m x_i^1 + a_2 \sum_{i=1}^m x_i^2 + \cdots + a_n \sum_{i=1}^m x_i^n &= \sum_{i=1}^m y_i x_i^0, \\ a_0 \sum_{i=1}^m x_i^1 + a_1 \sum_{i=1}^m x_i^2 + a_2 \sum_{i=1}^m x_i^3 + \cdots + a_n \sum_{i=1}^m x_i^{n+1} &= \sum_{i=1}^m y_i x_i^1, \\ &\vdots \\ a_0 \sum_{i=1}^m x_i^n + a_1 \sum_{i=1}^m x_i^{n+1} + a_2 \sum_{i=1}^m x_i^{n+2} + \cdots + a_n \sum_{i=1}^m x_i^{2n} &= \sum_{i=1}^m y_i x_i^n. \end{aligned}$$

Example 2 Fit the data in Table 8.3 with the discrete least squares polynomial of degree at most 2.

Solution For this problem, $n = 2$, $m = 5$, and the three normal equations are

$$\begin{aligned} 5a_0 + 2.5a_1 + 1.875a_2 &= 8.7680, \\ 2.5a_0 + 1.875a_1 + 1.5625a_2 &= 5.4514, \\ 1.875a_0 + 1.5625a_1 + 1.3828a_2 &= 4.4015. \end{aligned}$$

To solve this system using Maple, we first define the equations

$$\begin{aligned} eq1 &:= 5a_0 + 2.5a_1 + 1.875a_2 = 8.7680; \\ eq2 &:= 2.5a_0 + 1.875a_1 + 1.5625a_2 = 5.4514; \\ eq3 &:= 1.875a_0 + 1.5625a_1 + 1.3828a_2 = 4.4015 \end{aligned}$$

Table 8.3

i	x_i	y_i
1	0	1.0000
2	0.25	1.2840
3	0.50	1.6487
4	0.75	2.1170
5	1.00	2.7183

1.5

i	x_i	y_i	$\ln y_i$	x_i^2	$x_i \ln y_i$
1	1.00	5.10	1.629	1.0000	1.629
2	1.25	5.79	1.756	1.5625	2.195
3	1.50	6.53	1.876	2.2500	2.814
4	1.75	7.45	2.008	3.0625	3.514
5	2.00	8.46	2.135	4.0000	4.270
	7.50		9.404	11.875	14.422

If x_i is graphed with $\ln y_i$, the data appear to have a linear relation, so it is reasonable to assume an approximation of the form

$$y = be^{ax}, \quad \text{which implies that} \quad \ln y = \ln b + ax.$$

Extending the table and summing the appropriate columns gives the remaining data in Table 8.5.

Using the normal equations (8.1) and (8.2),

$$a = \frac{(5)(14.422) - (7.5)(9.404)}{(5)(11.875) - (7.5)^2} = 0.5056$$

and

$$\ln b = \frac{(11.875)(9.404) - (14.422)(7.5)}{(5)(11.875) - (7.5)^2} = 1.122.$$

With $\ln b = 1.122$ we have $b = e^{1.122} = 3.071$, and the approximation assumes the form

$$y = 3.071e^{0.5056x}.$$

$$a_0 = \frac{\sum_{i=1}^m x_i^2 \sum_{i=1}^m y_i - \sum_{i=1}^m x_i y_i \sum_{i=1}^m x_i}{m \left(\sum_{i=1}^m x_i^2 \right) - \left(\sum_{i=1}^m x_i \right)^2}$$

$$a_1 = \frac{m \sum_{i=1}^m x_i y_i - \sum_{i=1}^m x_i \sum_{i=1}^m y_i}{m \left(\sum_{i=1}^m x_i^2 \right) - \left(\sum_{i=1}^m x_i \right)^2}.$$

$$p_n(x) = \sum_{j=0}^n a_j x^j \quad \left. \vphantom{\sum_{j=0}^n} \right\} \rightarrow$$

$$E(a_0, a_1, \dots, a_n) = \sum_{i=1}^m (p_n(x_i) - y_i)^2$$

$$= \sum_{i=1}^m \left(\sum_{j=0}^n a_j x_i^j - y_i \right)^2$$

$$\frac{\partial E}{\partial a_0} = 0, \quad \frac{\partial E}{\partial a_1} = 0 \quad \dots \quad \frac{\partial E}{\partial a_n} = 0 \quad \left. \vphantom{\frac{\partial E}{\partial a_n}} \right\} \frac{\partial a_j}{\partial a_n} = \begin{cases} 1 \\ 0 \end{cases}$$

$$\frac{\partial E}{\partial a_n} = \sum_{i=1}^m 2 \left(\sum_{j=0}^n a_j x_i^j - y_i \right) \left(\sum_{j=0}^n x_i^j \underbrace{\frac{\partial a_j}{\partial a_n}} \right)$$

$$p_n(x) = \sum_{j=0}^n a_j x^j \quad \left. \vphantom{\sum_{j=0}^n} \right\} \rightarrow$$

$$E(a_0, a_1, \dots, a_n) = \sum_{i=1}^m (p_n(x_i) - y_i)^2$$

$$\frac{\partial E}{\partial a_n} = \sum_{i=1}^m 2 \left(\sum_{j=0}^n a_j x_i^j - y_i \right) \left(\sum_{j=0}^n x_i^j \frac{\partial a_j}{\partial a_n} \right)$$

$$\frac{\partial E}{\partial a_n} = 2 \sum_{i=1}^m \left(\sum_{j=0}^n a_j x_i^j - y_i \right) x_i^k = 0 \quad \dots \quad k=0 \dots n$$

$$\sum_{i=1}^m \sum_{j=0}^n a_j x_i^j x_i^k - \sum_{i=1}^m x_i^k y_i = 0 \quad \forall k=0 \dots n$$

$$\begin{aligned} n a_0 + \left(\sum_{i=1}^m x_i \right) a_1 + \left(\sum_{i=1}^m x_i^2 \right) a_2 + \dots - \left(\sum_{i=1}^m x_i^n \right) a_n &= \sum_{i=1}^m y_i \\ \left(\sum_{i=1}^m x_i \right) a_0 + \left(\sum_{i=1}^m x_i^2 \right) a_1 + \left(\sum_{i=1}^m x_i^3 \right) a_2 + \dots - \left(\sum_{i=1}^m x_i^{n+1} \right) a_n &= \sum_{i=1}^m x_i y_i \\ \vdots & \vdots \\ \left(\sum_{i=1}^m x_i^n \right) a_0 + \left(\sum_{i=1}^m x_i^{n+1} \right) a_1 + \left(\sum_{i=1}^m x_i^{n+2} \right) a_2 + \dots - \left(\sum_{i=1}^m x_i^{2n} \right) a_n &= \sum_{i=1}^m x_i^n y_i \end{aligned}$$

$$\begin{pmatrix} m & \sum x_i & \sum x_i^2 & \sum x_i^3 & \dots & \sum x_i^n \\ \sum x_i & \sum x_i^2 & \sum x_i^3 & \sum x_i^4 & \dots & \sum x_i^{n+1} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \sum x_i^n & \sum x_i^{n+1} & \sum x_i^{n+2} & \dots & \dots & \sum x_i^{2n} \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{pmatrix} = \begin{pmatrix} \sum y_i \\ \sum x_i y_i \\ \vdots \\ \sum x_i^n y_i \end{pmatrix}$$