

Columbia MA Math Camp

Optimization

Vinayak Iyer ^a

July 22, 2020

^aMaterial adapted from notes by David Thompson and Xingye Wu

Unconstrained Optimization

$$\max f(x)$$

Equality Constrained Optimization

$$\max f(x) \quad \text{s.t.} \cdot g(x) = c$$

$$\max x^{\frac{1}{2}} y^{\frac{1}{2}} \quad \text{s.t.} \cdot x + y = 1$$


Inequality Constrained Opt

$$\max f(x) \quad \text{s.t.} \cdot g(x) \leq c$$

May not cover. (Kuhn Tucker Theorem)

Some Definitions


x_1 & x_2
are local
maxima



Let $\mathcal{D} \subseteq \mathbb{R}^n$ and $f : \mathcal{D} \rightarrow \mathbb{R}$.

- A point c is a **maximum point** or **global maximum** of f if $f(c) \geq f(x)$ for all $x \in \mathcal{D}$
- A point c is a **local maximum** of f if there exists an $\epsilon > 0$ such that $f(c) \geq f(x)$ for all $x \in B_\epsilon(c)$
- If f is differentiable, a point such that $f'(c) = 0$ is a **critical point** of f

Maximum and minimum points are also called **extreme points**, **extremum** and **optimal points**



First-Order Conditions


Suppose f is concave & D is a closed set & bounded & differentiable
Is it necessary that at the maximum, $x^* \in D$
 $f'(x^*) = 0$?

Proposition 1.1

Let $D \subseteq \mathbb{R}^n$ be an open set, and $f : D \rightarrow \mathbb{R}$ a differentiable function. If f has a local extreme point at x , then $f'(x) = 0$

\iff \longrightarrow This condition isolates extreme points!

Notes : This could mean that x is either a maximum or minimum or neither



Second-Order Conditions

Twice continuously differentiable function!

Proposition 1.2

Let \mathcal{D} be an open set of \mathbb{R}^n and $f : \mathcal{D} \rightarrow \mathbb{R}$ a C^2 function.

- If f has a local maximum (minimum) at x , the Hessian of f at x is negative (positive) semi-definite
- If $f'(x) = 0$ and $H(x)$ is negative (positive) definite, then x is a strict local maximum (minimum)


(Proof of second result): Since $H(x)$ is negative definite, there exists $\epsilon > 0$ such that $H(\zeta)$ is negative definite for all $\zeta \in B_\epsilon(x)$. Using the exact form of Taylor's theorem, for any $y \in B_\epsilon(x)$:

$$\begin{aligned} f(y) &= f(x) + f'(x)(y-x) + \frac{1}{2}(y-x)^T H(\zeta)(y-x) \\ &< f(x) \end{aligned}$$

Maximum:- $f'(x) = 0$ & $f''(x) \leq 0$

Hessian negative semi-definite.

Global maxime
 $f(x_5) > f(x_1)$
 $\approx f(x_3)$




$$f'(x) = 0$$

$\{x_1, x_2, x_3, x_4, x_5\}$
 Critical points.

$$f''(x) < 0 \quad \leftarrow \{x_1, x_3, x_5\}$$

✓ $f(x_1), f(x_3), f(x_5)$

To find which point is
 the global maxime.



Summarizing

maxima/minima

- Critical points are a **necessary (not sufficient)** condition for extrema
- The second derivative can give us **local sufficient conditions**

To this point we've only discussed maximization over an open set

- Maxima need not exist on an open set (e.g. $f(x) = x$ on $(0, 1)$)
- If you're maximizing over a closed set S , you can decompose it as $S = \text{int}(S) \cup \text{Boundary}(S)$. Need to check the boundary

max $f(x)$
 $x \in (0, 1)$
 $f(x) = x^{\frac{1}{2}}$
maximizer
does not
exist!

General recipe to maximize a function :

- Find all critical points. If there are many, **SOC** can help filter
- Find the critical point with the largest value
- Check if the function takes on a higher value along the boundary

$f'(x) = 0$
filler out
the minima!

If $f'(x) = 0$ & $f''(x) < 0 \Rightarrow x$ is local max

$$f: [1, 2] \rightarrow \mathbb{R}.$$

Maximize



(1) find all critical points.

$$\rightarrow \langle x_1, x_2, x_3, x_4 \rangle$$

(b) filter out based on SOC

$$\rightarrow \langle x_1, x_4 \rangle$$

$$(2) f(x_1) \text{ \& } f(x_3) \rightarrow f(\overset{\text{critical pt}}{x_3}) > f(x_1)$$

(3) Check $f(1), f(2)$ &
compare to $f(x_3)$

→ 2 is the maxime
because

$$f(2) \geq f(x) \quad \forall x \in [1, 2]$$


$$\max x^{\frac{1}{2}}$$

$x \in (0,1)$ — Not compact

\Rightarrow max
not exists
necessarily.

$$f'(x) = 0 = \left[\frac{1}{2} x^{-\frac{1}{2}} \right]$$

$$x=0 \notin (0,1)$$



Why is it not guaranteed in this
simple problem that a max
exists?

$$x^* \in D \quad x^* \in (0,1)$$

Weierstrass: If f is cts & D is
compact \Rightarrow max exists.

$$\nRightarrow \neg P \Rightarrow \neg Q.$$


Sufficient Conditions for Global Extrema

For convex functions, optimization is dramatically simpler, as evidenced by the following proposition :

Proposition 1.3

Let \mathcal{D} be a convex open set of \mathbb{R}^n and $f : \mathcal{D} \rightarrow \mathbb{R}$ be a convex function. Then :

- ✓ The set of minimizers of f is convex
- ✓ If f is strictly convex, it has at most one minimizer
- Any local minimum of f is a global minimum
- ✓ If f is differentiable, then x is a global minimum of f iff $f'(x) = 0$



The same results hold for concave functions, replacing “minimizers” with “maximizers”

Note :

- Note that we already proved the second result for quasiconvex functions.

Convex fn $:- f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y)$

is a
(1) set of minimizers convex set

$$\forall x_1^*, x_2^* \in S_{\text{minimizers}}$$

$$\lambda x_1^* + (1-\lambda)x_2^* \in S_{\text{minimizers}}$$

$$\Rightarrow f(\lambda x_1^* + (1-\lambda)x_2^*) \leq f(x) \quad \forall x \in D$$

$$f(x_1^*) = f(x_2^*)$$

$$\begin{aligned} \Rightarrow f(\lambda x_1^* + (1-\lambda)x_2^*) &\leq f(x_1^*) = f(x_2^*) \\ &\leq f(x) \\ &\quad \forall x \in D. \end{aligned}$$

$$\Rightarrow \lambda x_1^* + (1-\lambda)x_2^* \in S_{\text{minimizers}}.$$



$$f(\lambda x + (1-\lambda)y) < \lambda f(x) + (1-\lambda)f(y)$$

f is strictly \downarrow convex \Rightarrow At most
1 minimizer.

Pf: Suppose there are 2 minimizers
 $x_1^* \neq x_2^*$. $f(x_1^*) = f(x_2^*)$

$$\begin{aligned} f(\underbrace{\lambda x_1^* + (1-\lambda)x_2^*}) &< \lambda f(x_1^*) \\ &\quad + (1-\lambda)f(x_2^*) \\ &= f(x_1^*) \end{aligned}$$

which contradicts $x_1^* \neq x_2^*$
being minimizers!

hence minimizer (if exists)
is unique.

Table of Contents

Unconstrained Optimization

Equality Constrained Optimization

Constrained Optimization Problems

- In economics, it's more common to maximize an objective function subject to some constraints
- For example, in consumer theory, you will see problems of this form:

$$\max_{c_1, c_2} \log c_1 + \alpha \log c_2 \quad \text{s.t.} \quad p_1 c_1 + p_2 c_2 = M$$

- One approach to these problems is to put the constraints in the objective. For instance, in the above example

$$c_1 = \frac{M - p_2 c_2}{p_1}$$

So we could do the unconstrained maximization problem:

$$\max_{c_2} \log \left(\frac{M - p_2 c_2}{p_1} \right) + \alpha \log c_2$$

[Do More Examples]

A more formal statement

Theorem 2.1

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}^k$ be C^1 functions, and consider the program:

$$\max_{x \in \mathbb{R}^n} f(x) \text{ s.t. } g(x) = 0$$

If x^* is a local maximum and x^* satisfies the constraint qualification ;
 $\text{rank}(g'(x^*)) = k$, then there exist k Lagrange multipliers $\lambda = (\lambda_1, \dots, \lambda_k)^T \in \mathbb{R}^k$ such
that the first-order condition holds:

$$f'(x^*) + \lambda^T g'(x^*) = 0$$

It is common to talk about the **Lagrangian** of a system:

$$\mathcal{L}(x, \lambda) = f(x) + \lambda^T g(x)$$

The first-order conditions wrt x and λ give us the critical point of the Lagrangian.

A sketch of Lagrange's Theorem in two variables

- Write $x = (x_1, x_2)$. Let $x^* = (x_1^*, x_2^*)$ be a local maximum of f subject to g
- By the IFT, we can write $x_2 = h(x_1)$, with $h'(x_1) = -\frac{g_1(x_1, x_2)}{g_2(x_1, x_2)}$
- We now do unconstrained optimization of $f(x_1, h(x_1))$. The FOC is

$$f_1(x^*) + f_2(x^*)h'(x_1^*) = 0$$

- Define $\lambda = -\frac{f_2(x^*)}{g_2(x^*)}$. Then

$$f_1 + \lambda g_1(x^*) = 0$$

$$f_2 + \lambda g_2(x^*) = 0$$

The general case is similar, just with more cumbersome matrix notation.

- The Lagrange condition is a necessary condition.
- As with unconstrained optimization, there are second order conditions that let you check whether a critical point of the Lagrangian is a local maximum or minimum (see FMEA Section 3.4 for details)
- In order to get sufficient conditions for global maxima along the constraint, we need additional structure
- If the constraint qualification fails, the theorem says nothing. So you need to check points where the CQ fails separately

An example of a sufficient condition

Proposition 2.1

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be strictly quasiconcave and consider the program

$$\max_x f(x) \text{ s.t. } Ax = b \quad (1)$$

where A is an $m \times n$ matrix with $m < n$. If (x^*, λ^*) is a critical point of the Lagrangian and $f'(x^*) \neq 0$, then x^* solves (1).

Proof.

The FOC of the Lagrangian implies $f'(x^*) + \lambda^T A = 0$. Suppose there were an \hat{x} such that $A\hat{x} = b$ and $f(\hat{x}) > f(x^*)$. Since f is strictly quasiconcave:

$$\begin{aligned} 0 &< f'(x^*)(\hat{x} - x^*) \\ &= -\lambda^T A(\hat{x} - x^*) \\ &= 0 \end{aligned}$$

a contradiction. □

Note : $f(x)$ being strictly quasiconcave and the constraint being linear ensures that the Lagrangian is strictly quasiconcave as well (which we have shown before implies a unique maximizer)

Define the "**value function**" V as follows

$$V(b) = \max_x f(x) \text{ s.t. } g(x) = b$$

Form the Lagrangian :

$$\mathcal{L}(x, \lambda, b) = f(x) + \lambda^T (b - g(x))$$

Write the solution of this problem as $x^*(b), \lambda^*(b)$. Then

$$V(b) = \mathcal{L}(x^*(b), \lambda^*(b), b)$$

Interpretation of the multipliers (cont.)

Using the chain rule, we have :

$$V'(b) = \frac{\partial \mathcal{L}}{\partial x} \frac{dx^*}{db} + \frac{\partial \mathcal{L}}{\partial \lambda} \frac{d\lambda^*}{db} + \frac{\partial \mathcal{L}}{\partial b}$$

However, we know $\frac{\partial \mathcal{L}}{\partial x}$ and $\frac{\partial \mathcal{L}}{\partial \lambda}$ are 0 at x^*, λ^* , so we have

$$V'(b) = \lambda^T$$

Interpretation: Lagrange multipliers measure the marginal value of loosening a constraint by 1 unit