


Complex Numbers

Consider the complex plane, with the imaginary unit j defined by $j^2 = -1$



Rectangular **Form**

$$z_0 = x_0 + jy_0$$

$$x_0 = Re\{z_0\}$$

$$y_0 = Im\{z_0\}$$

Polar Form


$$z_0 = r_0 e^{j\theta_0}$$

$$r_0 = |z_0|$$
$$\theta_0 = \angle z_0$$

$$\theta_0 = \angle z_0$$



Complex Conjugate




$$z_0^* = x_0 - jy_0$$

$$z_0^* = r_0 e^{-j\theta_0}$$




Euler's Identity



$$e^{j\theta_0} = \cos\theta_0 + j\sin\theta_0$$



Euler's Inverse Identities




$$\cos \theta_0 = \frac{e^{j\theta_0} + e^{-j\theta_0}}{2}$$

$$\sin \theta_0 = \frac{e^{j\theta_0} - e^{-j\theta_0}}{2j}$$



Conversion From Polar to Rectangular Form




$$z_0 = r_0 e^{j\theta_0} = r_0 \cos \theta_0 + jr_0 \sin \theta_0$$

$$x_0 = Re\{z_0\} = r_0 \cos \theta_0$$
$$y_0 = Im\{z_0\} = r_0 \sin \theta_0$$

$$y_0 = Im\{z_0\} = r_0 \sin \theta_0$$



Conversion From Rectangular to Polar Form



$$z_0 = x_0 + jy_0$$

$$r_0 = \sqrt{x_0^2 + y_0^2}$$

$$\theta_0 = \tan^{-1} \frac{y_0}{x_0}$$

$$z_0 = \sqrt{x_0^2 + y_0^2} e^{j \tan^{-1} \left(\frac{y_0}{x_0}\right)}$$



An Example

- Consider the complex number $z_0 = 3 + j4$
 - Express the complex number in polar form
 - Find its complex conjugate

