

Computational Physics - Project 2

Johannes Scheller, Vincent Noculak, Lukas Powalla

September 22, 2015

Contents

1	Introduction And Motivation	3
2	Theory	3
2.1	Rewriting Schrödinger's equation as eigenvalue problem	3
2.1.1	One electron in a harmonic oscillator potential	3
2.1.2	Two interacting electrons in a harmonic oscialltor potential	3
2.2	Jacobi's method	3
3	Execution	3
3.1	Implementing the algorithm	3
3.2	Results	3
4	Comparison and discussion of the results	3

1 Introduction And Motivation

In many fields of both mathematics and physics, we often come to the point that we have to solve so-called eigenvalue problems, which are equations of the form $\hat{A} \cdot \hat{v} = \lambda \hat{v}$, where \hat{A} is a matrix of dimension $n \times n$ and v is a vector of dimension n . Equations of this kind occur not only in linear algebra, but also in mechanics and quantum mechanics and will also be a major part of this report. In this project, we are going to rewrite the Schrödinger's equation of one and two electrons in a harmonic oscillator potential in the form of an eigenvalue problem and solve it numerically by implementing Jacobi's method, an algorithm that can be used to solve any eigenvalue problem.

2 Theory

2.1 Rewriting Schrödinger's equation as eigenvalue problem

2.1.1 One electron in a harmonic oscillator potential

2.1.2 Two interacting electrons in a harmonic oscillator potential

2.2 Jacobi's method

3 Execution

3.1 Implementing the algorithm

3.2 Results

4 Comparison and discussion of the results