

The Quench-Action method in integrable spin chains: A Monte Carlo approach

Vincenzo Alba¹, Pasquale Calabrese¹

¹ International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy, INFN, Sezione di Trieste

Abstract.

fdasfa

1. Introduction

We show that it is possible to numerically simulate the Quench Action approach combining Monte Carlo methods and Bethe ansatz techniques.

We focus on the situation in which the pre-quench initial state is the Neel state or the Majumdar-Ghosh state.

We investigate the importance of the zero-momentum strings in the Quench Action.

Without zero-momentum strings the overlap saturation rules are not valid, i.e., in finite size systems the vast majority of the eigenstates contain zero momentum strings.

The details on the eigenstates counting depend on the pre-quench initial state.

However, we show that one can restrict to the set of non-zero momentum strings. The fact that one neglects zero-momentum strings gives rise only to scaling corrections.

We also investigate the validity of the Bethe-Takahashi approximation for the calculation of the overlap.

2. The Heisenberg spin chain

Here we consider the spin- $\frac{1}{2}$ isotropic Heisenberg chain (XXX chain). The XXX chain with L sites is defined by the Hamiltonian

$$\mathcal{H} \equiv J \sum_{i=1}^L \left[\frac{1}{2} (S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+) + S_i^z S_{i+1}^z \right], \quad (1)$$

where $S_i^\pm \equiv (\sigma_i^x \pm i\sigma_i^y)/2$ are spin operators acting on the site i , $S_i^z \equiv \sigma_i^z/2$, and $\sigma_i^{x,y,z}$ the Pauli matrices. We fix $J = 1$ and use periodic boundary conditions, identifying sites $L+1$ and 1 . The total magnetization $S_T^z \equiv \sum_i S_i^z = L/2 - M$, with M number of down spins (particles), commutes with (1), and it is here used to label its eigenstates.

2.1. Bethe equations and wavefunctions

The generic eigenstate of (1) in the sector with M particles can be written as

$$|\Psi_M\rangle = \sum_{1 \leq x_1 < x_2 < \dots < x_M \leq L} A_M(x_1, x_2, \dots, x_M) |x_1, x_2, \dots, x_M\rangle, \quad (2)$$

where the sum is over the positions $\{x_i\}$ of the particles, and $A_M(x_1, x_2, \dots, x_M)$ is the eigenstate amplitude corresponding to particles at positions x_1, x_2, \dots, x_M . $A_M(x_1, x_2, \dots, x_M)$ is given as

$$A_M(x_1, x_2, \dots, x_M) \equiv \sum_{\mathcal{P} \in S_M} \exp \left[i \sum_{j=1}^M k_{\mathcal{P}_j} x_j + i \sum_{i < j} \theta_{\mathcal{P}_i \mathcal{P}_j} \right]. \quad (3)$$

Here the outermost sum is over the permutations S_M of the so-called quasi-momenta $\{k_1, k_2, \dots, k_M\}$. The two-particle scattering phases $\theta_{m,n}$ are defined as

$$\theta_{m,n} \equiv \frac{1}{2i} \log \left[- \frac{e^{ik_m + ik_n} - 2e^{ik_m} + 1}{e^{ik_m + ik_n} - 2e^{ik_n} + 1} \right]. \quad (4)$$

The energy associated to the eigenstate (2) is

$$E = \sum_{\alpha=1}^M (\cos(k_\alpha) - 1). \quad (5)$$

The quasi-momenta $\{k_\alpha\}$ are obtained by solving the so-called Bethe equations

$$e^{ik_\alpha L} = \prod_{\beta \neq \alpha}^M \left[-\frac{1 - 2e^{ik_\alpha} - e^{ik_\alpha + ik_\beta}}{1 - 2e^{ik_\beta} - e^{ik_\alpha + ik_\beta}} \right]. \quad (6)$$

It is useful to introduce the rapidities $\{\lambda_\alpha\}$ as

$$k_\alpha = \pi - 2 \arctan(\lambda_\alpha) \pmod{2\pi}. \quad (7)$$

Taking the logarithm on both sides in (6), and using (7), one obtains the Bethe equations in logarithmic form as

$$\arctan(\lambda_\alpha) = \frac{\pi}{L} J_\alpha + \frac{1}{L} \sum_{\beta \neq \alpha} \arctan\left(\frac{\lambda_\alpha - \lambda_\beta}{2}\right), \quad (8)$$

where $-L/2 < J_\alpha \leq L/2$ are the so-called Bethe quantum numbers. The J_α are half-integers and integers for $L - M$ even and odd.

These solutions of the Bethe equations (6) form particular “string” patterns in the complex plane, in the limit of large chains $L \rightarrow \infty$ (string hypothesis) [?, ?]. Specifically, rapidities forming a “string” of length $1 \leq n \leq M$ (that we defined here as n -string) are parametrized as

$$\lambda_{n;\gamma}^j = \lambda_{n;\gamma} - i(n - 1 - 2j) + i\delta_{n;\gamma}^j, \quad j = 0, 1, \dots, n - 1, \quad (9)$$

where $\lambda_{n;\gamma}$ is the real part of the string (string center), and γ labels strings with different centers, while j labels the different components of the string. In (9) $\delta_{n;\gamma}^j$ are the string deviations, which typically vanish exponentially in the thermodynamic limit.

Notice that pure real rapidities are strings of unit length (1-strings).

2.2. Bethe-Takahashi equations

The string centers $\lambda_{n;\gamma}$ in (9) are obtained by solving the so-called Bethe-Takahashi equations

$$2L\theta_n(\lambda_{n;\gamma}) = 2\pi I_{n;\gamma} + \sum_{(m,\beta) \neq (n,\gamma)} \Theta_{m,n}(\lambda_{n;\gamma} - \lambda_{m;\beta}), \quad (10)$$

where the generalized scattering phases $\Theta_{m,n}$ read

$$\Theta_{m,n}(x) \equiv \begin{cases} \theta_{|n-m|}(x) + \sum_{r=1}^{(n+m-|n-m|-1)/2} 2\theta_{|n-m|+2r}(x) + \theta_{n+m}(x) & \text{if } n \neq m \\ \sum_{r=1}^{n-1} 2\theta_{2r}(x) + \theta_{2n}(x) & \text{if } n = m \end{cases}$$

and $\theta_\alpha(x) \equiv 2 \arctan(x/\alpha)$. Here $I_{n;\gamma}$ are the Bethe-Takahashi quantum numbers associated with $\lambda_{n;\gamma}$.

Each M -particle eigenstate can be characterized by its “string content” $\mathcal{S} \equiv \{s_1, \dots, s_M\}$, with s_n the number of n -strings.

It can be shown that $I_{n;\gamma}$ are integers or half-integers for $L - s_n$ odd and even, respectively.

Clearly, the constraint $\sum_{\alpha=1}^M \alpha s_\alpha = M$ has to be satisfied. The upper bound for the Bethe-Takahashi quantum numbers can be derived as

$$|I_{n;\gamma}| \leq I_n^{(MAX)} \equiv \frac{1}{2}(L - 1 - \sum_{m=1}^M t_{m,n} s_m), \quad (11)$$

where $t_{m,n} \equiv 2\min(n, m) - \delta_{m,n}$.

3. Overlap with the Neel state

Here we restrict to the parity-invariant eigenstates. These are the only eigenstates with non-zero overlap with the Neel state. Parity-invariant eigenstates contain only pairs of rapidities with opposite sign.

We denote the generic parity invariant eigenstate as $|\{\pm\lambda_j\}_{j=1}^m, n_\infty\rangle$, where m is the number of rapidity pairs, N_∞ is the number of infinite rapidities, with $M = L/2 = N_\infty + 2m$, and $n_\infty \equiv N_\infty/L$ is the density of infinite rapidities.

The overlap with the Neel state $|N\rangle$ reads

$$\frac{\langle N | \{\pm\lambda_j\}_{j=1}^m, n_\infty \rangle}{||\{\lambda_j\}_{j=1}^m, n_\infty\rangle||} = \frac{\sqrt{2}N_\infty!}{\sqrt{(2N_\infty)!}} \left[\prod_{j=1}^m \frac{\sqrt{\lambda_j^2 + 1}}{4\lambda_j} \right] \sqrt{\frac{\det_m(G^+)}{\det_m(G^-)}} \quad (12)$$

where

$$G_{jk}^\pm = \delta_{jk} \left(N K_{1/2}(\lambda_j) - \sum_{l=1}^m K_1^+(\lambda_j, \lambda_l) \right) + K_1^\pm(\lambda_j, \lambda_k), \quad j, k = 1, \dots, m \quad (13)$$

and

$$K_1^\pm(\lambda, \mu) = K_1(\lambda - \mu) \pm K_1(\lambda + \mu) \quad (14)$$

and

$$K_\alpha(\lambda) \equiv \frac{8\alpha}{\lambda^2 + 4\alpha^2} \quad (15)$$

3.1. Reduced Neel overlap

Here we consider the overlap formula for the Neel state (12) in the limit $L \rightarrow \infty$, assuming that the rapidities form perfect strings.

In the case of perfect strings the matrices G_{jk}^\pm become ill-defined. Precisely, $K_1^\pm(\lambda, \mu)$ diverges if λ and μ are successive members of the same string, i.e., $|\lambda - \mu| = 2i$.

It is possible to rewrite (12) in terms of the string centers $\lambda_{n;\alpha}$ only. Here we restrict ourselves to rapidity configurations with no zero-momentum strings. Our results are not valid for zero-rapidity strings. These would require the knowledge of the precise form of

the string deviations, i.e., the dependence of the string deviations on L , as it has been pointed out in Ref. [1].

It is convenient to split the indices i, j of G_{ij}^\pm as $i = (n, \alpha)$ $j = (m, \beta)$, with n, m being the length of the strings and α, β labelling the string centers.

The result reads

$$\frac{1}{2}G_{(n,\alpha)(m,\beta)}^+ = \begin{cases} L\theta'_n(\lambda_{n;\alpha}) - \sum_{(\ell,\gamma) \neq (n,\alpha)} \left[\Theta'_{n,\ell}(\lambda_{n;\alpha} - \lambda_{\ell;\gamma}) + \Theta'_{n,\ell}(\lambda_{n;\alpha} + \lambda_{\ell;\gamma}) \right] & \text{if } (n, \alpha) = (m, \beta) \\ \Theta'_{n,m}(\lambda_{n;\alpha} - \lambda_{m;\beta}) + \Theta'_{n,m}(\lambda_{n;\alpha} + \lambda_{m;\beta}) & \text{if } (n, \alpha) \neq (m, \beta) \end{cases} \quad (16)$$

Here $\theta'_n(x) \equiv d\theta_n(x)/dx = 2n/(n^2 + x^2)$ and $\Theta'(x) \equiv d\Theta(x)/dx$.

For G_{ij}^- one obtains

$$\frac{1}{2}G_{(n,\alpha)(m,\beta)}^- = \begin{cases} (L-1)\theta'_n(\lambda_{n;\alpha}) - 2 \sum_{k=1}^{n-1} \theta'_k(\lambda_{n;\alpha}) & \text{if } (n, \alpha) = (m, \beta) \\ - \sum_{(\ell,\gamma) \neq (n,\alpha)} \left[\Theta'_{n,\ell}(\lambda_{n;\alpha} - \lambda_{\ell;\gamma}) + \Theta'_{n,\ell}(\lambda_{n;\alpha} + \lambda_{\ell;\gamma}) \right] & \text{if } (n, \alpha) = (m, \beta) \\ \Theta'_{n,m}(\lambda_{n;\alpha} - \lambda_{m;\beta}) - \Theta'_{n,m}(\lambda_{n;\alpha} + \lambda_{m;\beta}) & \text{if } (n, \alpha) \neq (m, \beta) \end{cases} \quad (17)$$

Finally, the multiplicative prefactor in (12) for the generic n -string can be rewritten as

$$\prod_{a=1}^n \frac{\sqrt{(\lambda_{n;\alpha}^a)^2 + 1}}{4\lambda_{n;\alpha}^a} = \frac{1}{4^n} \left(\frac{\sqrt{n^2 + \lambda_{n;\alpha}^2}}{\lambda_{n;\alpha}} \prod_{k=0}^{\lceil n/2 \rceil - 1} \frac{(2k)^2 + \lambda_{n;\alpha}^2}{(2k+1)^2 + \lambda_{n;\alpha}^2} \right)^{\mathcal{P}}, \quad (18)$$

with $\mathcal{P} = +$ and $\mathcal{P} = -$ for even and odd strings, respectively.

4. Overlap with the Majumdar-Ghosh state

The overlap between the generic eigenstate of the XXX chain and the Majumdar-Ghosh state is obtained from the overlap with the Neel state as

$$\langle MG | \{\lambda_j\}_{j=1}^M \rangle = \prod_{j=1}^M \frac{1}{\sqrt{2}} \left(1 - \frac{\lambda_j - i}{\lambda_j + i} \right) \langle N | \{\lambda_j\}_{j=1}^M \rangle \quad (19)$$

The mutliplicative factor in (??), using the string hypothesis for the generic n -string is rewritten as

$$\prod_{j=1}^M \frac{1}{\sqrt{2}} \left(1 - \frac{\lambda_j - i}{\lambda_j + i} \right) = 2^n \prod_{k=0}^{\lfloor n/2 \rfloor} \frac{1}{[\lambda_{n;\gamma}^2 + (2k + (1 - (-1)^n)/2)^2]^2} \quad (20)$$




Figure 1. Overlap sum rules for the Neel state: The role of the zero-momentum strings. (a) The overlap sum rule $\sum_{\lambda} |\langle \lambda | N \rangle|^2 = 1$, with $|N\rangle$ the Neel state and $|\lambda\rangle$ the eigenstates of the XXX spin chain. The x -axis shows $1/L$, with L the chain length. The circles are Bethe ansatz results for chains up to $L = 36$. The results are obtained via a full scanning of the chain Hilbert space. Only the eigenstates with no zero-momentum strings are considered. The dash-dotted line is a fit to $A/L^{1/2} + B/L$, with A, B fitting parameters. Inset: The same data as in the main Figure plotted versus $1/L^{1/2}$. (b) The same as in (a) for the sum rule $\sum_{\lambda} |\langle \lambda | N \rangle|^2 e(\lambda) = 1/2$, with $e(\lambda)$ the eigenstates energy density.

5. Overlap sum rules: the role of the zero-momentum strings

In this section we investigate the role of the zero-momentum strings. We focus on conserved quantities sum rules. Specifically, given a generic conserved quantity (charge) \hat{Q} we consider the trivial identity

$$Q_0 \equiv \langle \Psi_0 | \hat{Q} | \Psi_0 \rangle = \sum_{\lambda} |\langle \lambda | \Psi_0 \rangle|^2 Q_{\lambda}, \quad (21)$$

where Q_0 is the charge expectation values over the initial state $|\Psi_0\rangle$, $|\lambda\rangle$ denotes the generic eigenstate of the XXX chain, and Q_{λ} is the charge eigenvalue over the eigenstate.

Here we restrict ourselves to the cases with $\hat{Q} = \mathbb{I}$ and $\hat{Q} = \mathcal{H}$, considering both the case with $|\Psi\rangle = |N\rangle$ and $|\Psi_0\rangle = |MG\rangle$.

For the Néel state, for any finite L Eq. (21) gives $Q_0 = 1, -1/2$ for $\hat{Q} = \mathbb{I}$ and $\hat{Q} = \mathcal{H}$, respectively.

Figure 1 plots the numerical results obtained via a full scanning of the Hilbert space of the Heisenberg chain. Panel (a) and (b) in the Figure plots the sum rule (21) for $\hat{Q} = \mathbb{I}$ and $\hat{Q} = \mathcal{H}$, respectively. The circles are obtained considering all the eigenstates of the Heisenberg spin chain with L sites, excluding eigenstates containing zero-momentum strings.

The total number of eigenstates \tilde{Z}_{Neel} containing no zero-momentum strings is given as

$$\tilde{Z}_{Neel} = B\left(\frac{L}{2}, \frac{L}{4}\right). \quad (22)$$




Figure 2. Overlap sum rules for the Majumdar-Ghosh (MG) state. (a) The overlap sum rule $\sum_{\lambda} |\langle \lambda | MG \rangle|^2 = 1$, with $|MG\rangle$ the Majumdar-Ghosh state and $|\lambda\rangle$ the eigenstates of the XXX spin chain. The x -axis shows $1/L$, with L the chain length. The circles are Bethe ansatz results for chains up to $L = 36$. The results are obtained via a full scanning of the chain Hilbert space. Only the eigenstates with no zero-momentum strings are considered. The dash-dotted line is a fit to $A/L + B/L^2$, with A, B fitting parameters. (b) The same as in (a) for the sum rule $\sum_{\lambda} |\langle \lambda | MG \rangle|^2 e(\lambda) = 2/3$, with $e(\lambda) \equiv E/L$ the eigenstates energy density.

The proof of (22) is reported in [Appendix B](#).

The largest chain size considered is $L = 36$. The data are plotted versus $1/L$.

Interestingly, both the sum rules (panels (a)(b)) are not saturated, as expected since the eigenstates corresponding to zero-momentum strings are excluded in the sum in the right-hand-side in (21).

Interestingly, both sum rules exhibit vanishing behavior upon increasing the chain size. The dash-dotted lines in the Figures are fits to the behavior $A/L^{1/2} + B/L$, with A, B fitting parameters, and perfectly describe the behavior of the data.

The vanishing behavior as $\propto 1/L^{1/2}$ of the sum rules reflects the vanishing of the fraction of non-zero momentum string eigenstates in the thermodynamic limit ad

$$\frac{\tilde{Z}_{Neel}}{Z_{Neel}} \propto \frac{4}{\sqrt{\pi L}}, \quad (23)$$

with Z_{Neel} being the total number of eigenstates corresponding to parity invariant BGT quantum numbers configurations.

One should observe that the asymptotic, i.e., at large L , behavior (23) is not generic, meaning that different initial states $|\Psi_0\rangle$ might give different behaviors.

This is illustrated in Figure 2, focusing on the Majumdar-Ghosh (MG) state. Panels (a) and (b) plot the sum rules with $\hat{Q} = \mathbb{I}$ and $\hat{Q} = \mathcal{H}$ (same as in Figure 1). As for the Néel state only eigenstates containing no zero-momentum strings are considered. Note that their number \tilde{Z}_{MG} (see [Appendix B](#)) is now given as

$$\tilde{Z}_{MG} = B\left(\frac{L}{2}, \frac{L}{4}\right) - B\left(\frac{L}{2}, \frac{L}{4} - 1\right). \quad (24)$$

Similar to Figure 1, due to the exclusion of the zero-momentum strings, the saturation rules exhibit vanishing behavior in the thermodynamic limit. However, in contrast with the Néel case (see Figure 1), one has the behavior $Q_0 \sim 1/L$, as confirmed by the fits (dash-dotted lines in Figure 2).

Similar to the Néel case this reflects the same vanishing behavior of the eigenstates containing no zero-momentum strings \tilde{Z}_{MG}/Z_{MG} as

$$\frac{\tilde{Z}_{MG}}{Z_{MG}} = \frac{4}{4+L}. \quad (25)$$

6. Validity of the string hypothesis for overlap calculations

Here we discuss the validity of the string hypothesis when calculating the overlaps between the eigenstates of the XXX chain and the Néel state. Here we focus on the Heisenberg chain with $L = 20$ sites. Figure 3 plots the squared overlaps $|\langle \lambda | N \rangle|^2$ between the Néel state and the eigenstates of the chain. The overlaps are plotted against the eigenstate energy density $E/L \in [-\log(2), 0]$. The circles are exact diagonalization results for all the chain eigenstates (382 eigenstates), whereas the crosses denote the overlaps calculated using formula (12), and the Bethe-Gaudin-Takahashi equations. Note that only the eigenstates with no zero-momentum strings are shown (252 eigenstates) in the Figure. Panel (a) in the Figure is an overview of all the results. Panels (b)-(d) correspond to zooming to the smaller overlap values $|\langle N | \lambda \rangle| \lesssim 0.02$, $|\langle N | \lambda \rangle| \lesssim 0.002$, and $|\langle N | \lambda \rangle| \lesssim 10^{-5}$.

Clearly, the overlaps decay rapidly upon increasing the energy density. This is expected since the XXX Hamiltonian expectation value over the Néel state is $\langle N | H | N \rangle = -1/2$. Importantly, the agreement between the exact diagonalization results and the results obtained using the BGT equations (6) is excellent, confirming the validity of the string hypothesis.

7. Monte Carlo implementation of the Quench-Action approach

For the finite-size Heisenberg spin chain the Quench-Action expectation values can be obtained by sampling the eigenstates of the chain using Monte Carlo. One starts with an eigenstate of the XXX chain with N_∞ infinite rapidities and $M \equiv L/2 - N_\infty$ finite ones. The state is identified by a parity-invariant Bethe quantum number configuration \mathcal{C} and by the corresponding parity-invariant rapidities $\{\lambda\}$ as $|\lambda\rangle$. The string content associated with the finite rapidities is denoted as \mathcal{S} . The Monte Carlo procedure consists of four steps as follows:

- ① Choose a new number of finite rapidities M' with probability

$$\mathcal{P}(M') = \frac{\tilde{Z}'_{Neel}(L, M')}{\tilde{Z}_{Neel}(L)}. \quad (26)$$




Figure 3. The squared overlap $|\langle N|\lambda\rangle|^2$ between the the Neel state $|N\rangle$ and the eigenstates $|\lambda\rangle$ of the XXX chain with $L = 20$ sites. Only non-zero overlaps are shown. In all the panels the x -axis shows the eigenstate energy density E/L . The circles are the exact diagonalization results for all the non-zero overlaps. The crosses are the Bethe ansatz results obtained using the Bethe-Gaudin-Takahashi equations. The missing crosses correspond to eigenstates containing zero-momentum strings. (a) Overview of all the non-zero overlaps. (b)(c)(d) The same overlaps as in (a) zooming in the regions $[0, 0.2]$, $[0, 0.020]$, and $[0, 4 \cdot 10^{-5}]$. The discrepancies between the ED and the Bethe ansatz results are attributed to the string deviations.

- ② Choose a new a new string content $\mathcal{S}' \equiv \{s'_1, \dots, s'_{M'}\}$ with probability $\mathcal{P}'(M', \mathcal{S}')$

$$\mathcal{P}'(M', \mathcal{S}') = \frac{1}{\widetilde{Z}'_{Neel}(L, M')} \prod_{n=1}^{M'} B\left(\frac{L}{2} - \frac{1}{2} \sum_{m=1}^{M'} t_{nm} s'_m, s'_n\right). \quad (27)$$

- ③ Generate a new parity-invariant quantum number configuration \mathcal{C}' compatible with the \mathcal{S}' obtained in step ②. Solve the corresponding BGT equations (10), finding a new eigenstate $|\lambda'\rangle$.
- ④ Calculate the overlap $\langle \lambda'|N\rangle$ between the new eigenstate and the Néel state and accept the eigenstate with the Metropolis probability

$$\mathcal{P}''_{\lambda \rightarrow \lambda'} = \text{Min}\left\{1, \exp\left(-2\Re(\mathcal{E}' - \mathcal{E})\right)\right\}, \quad (28)$$

where $\mathcal{E}' \equiv -\log\langle \lambda'|N\rangle$, similarly for \mathcal{E} , and \Re denotes the real part.

Afte some necessary thermalization steps the sequence 1 – 4 generates eigenstates of the XXX chain distributed according to the Quench-Action probability distribution.

For a generic observable \mathcal{O} , its Quench-Action expectation value $\langle \mathcal{O} \rangle_{QA}$ is obtained as the arithmetic average of the expectation value over the eigenstates sampled in the Monte Carlo as

$$\langle \mathcal{O} \rangle_{QA} = \frac{1}{N_{mcs}} \sum_{\lambda} \langle \lambda | \mathcal{O} | \lambda \rangle, \quad (29)$$

with N_{mcs} being the total number of Monte Carlo steps.




Figure 4. The overlap sum rules for the Neel state: Numerical results obtained using the Hilbert space Monte Carlo sampling approach. (a) The energy sum rule $\langle N|Q_2|N\rangle/L = -1/2$, with Q_2/L the Hamiltonian density. We plot $\sum_\lambda |\langle \lambda|N\rangle|^2 Q_2(\lambda)/L = 1/2$, with $|\lambda\rangle$ the eigenstates of the XXX chain, versus the inverse chain length $1/L$. The symbols are Monte Carlo data obtained by sampling the eigenstates of the XXX chain. The dash-dotted line is the expected result. The dashed line is a fit to the behavior $-1/2 + A/L + B/L^2$, with A, B fitting parameters. (b) The energy fluctuations sum rule $\sigma^2(Q_2)/L \equiv (\langle N|Q_2^2|N\rangle - \langle N|Q_2|N\rangle^2)/L = 1/4$. The horizontal line is the expected result. (c)(d) Same as in (a)(b) for the charge Q_4 and its fluctuations.

8. Monte Carlo Quench-Action approach: Numerical results

In this section we numerically demonstrate that the only effect of neglecting the zero-momentum strings are finite-size corrections.

The validity of the Monte Carlo approach for simulating the Quench-Action results is demonstrated in Figure 4. The Figure focuses on the Néel sum rules for the conserved charges Q_2 and Q_4 . Specifically, panel (a) shows the sum rules for $\hat{Q} = \hat{Q}_2$, $\hat{Q} = \hat{Q}_4$, and the corresponding fluctuations $\sigma^2(Q_n) \equiv \langle N|\hat{Q}_n^2|N\rangle - \langle N|\hat{Q}_n|N\rangle^2$.

$$\langle N|\hat{Q}_n|N\rangle = \sum_\lambda |\langle N|\lambda\rangle|^2 Q_n. \quad (30)$$

Note that the Néel sum rule for the identity \mathbb{I} is trivially satisfied, and it corresponds to the normalization of the Monte Carlo probability. Figure 4 (a) plots the Néel sum rule for the energy density Q_2/L . The circles are Monte Carlo data obtained using the procedure outlined in 7 for several Heisenberg spin chains. The largest chain simulated is for $L = 56$ sites. The data correspond to a simulation with $\sim 10^7$ Monte Carlo step (mcs). The expected result $\langle N|\hat{Q}_2|N\rangle/L = -1/2$ is shown as dash-dotted line.

In stark contrast with Figure 1 and Figure 2, while for small chains the sum rule is violated the data strongly suggest that in the thermodynamic limit the sum rule saturates the expected theoretical value.

Furthermore, the data suggest the behavior $\propto 1/L$, as confirmed by the fit to $-1/2 + A/L + b/L^2$ (dashed line in the Figure), with A, B fitting parameters.




Figure 5. The steady-state root densities $\rho_1(x)$ (panel (a)) and $\rho_2(x)$ (panel (b)), plotted versus the root x . The histograms are Monte Carlo data for the Heisenberg chain with $L = 56$ sites and $\sim 10^7$ Monte Carlo steps. The y -axis is divided by 10^6 for convenience. The continuous lines are the analytical Quench-Action results in the thermodynamic limit.

The same qualitative behavior is demonstrated in panel (b) for the energy fluctuations. Interestingly, for $L = 56$ the Monte Carlo result is compatible within the error bar with the expected asymptotic one.

Finally, panels (c)(d) show the same results as for (a)(b) for the conserved charge Q_4 .

9. The Quench-Action root distributions

The BGT root distributions $\rho_n(\lambda)$ identifying the Quench-Action saddle point can be extracted from the Monte Carlo simulation as shown in [2].

Figure 5 displays the first two root distributions $\rho_1(\lambda)$ (panel (a)) and $\rho_2(\lambda)$ (panel (b)), plotted against the BGT root λ . The data are the histograms of the BGT roots sampled in the Monte Carlo. The data are for a chain with $L = 56$ sites and $N_{mcs} \sim 10^7$ Monte Carlo steps. All the data are divided by a factor 10^6 for plotting convenience. The width of the histogram bins $\Delta\lambda$ is $\Delta\lambda \approx 0.02$ and $\Delta\lambda \approx 0.001$ for $\rho_1(\lambda)$ and $\rho_2(\lambda)$, respectively.

The histogram fluctuations are due to N_{mcs} being finite, and are expected to vanish in the limit $N_{mcs} \rightarrow \infty$.

The continuous lines is the expected analytic result in the thermodynamic limit.

Clearly, $\rho_1(\lambda)$ is in good agreement with the Monte Carlo data in the whole range $-2 \leq \lambda \leq 2$ considered. On the other hand, larger deviations from the theoretical result are present for $\rho_2(\lambda)$. These deviations are larger on the tails of the distribution. This

is expected since large λ correspond to large quasimomenta, which are more sensitive to the finite size of the chain.

9.1. Counting of eigenstates with non-zero Neel overlap

We numerically checked that the number of states with non-zero overlap with the Neel state is given as

$$Z_{Neel} = 2^{\frac{L}{2}-1} + \frac{1}{2}B\left(\frac{L}{2}, \frac{L}{4}\right) + 1, \quad (31)$$

with $B(x, y)$ denoting the binomial coefficient. The contribution 1 accounts for the ferromagnetic state. Here we assumed that L is divisible by four. Here Z_N is obtained as the total number of parity-invariant Bethe-Gaudin-Takahashi quantum numbers.

After excluding the zero-momentum strings the total number of states with non-zero overlap with the Neel state is

$$\tilde{Z}_{Neel} = \left(\frac{L}{2}, \frac{L}{4}\right) \quad (32)$$

Importantly, the fraction of eigenstates corresponding to non-zero momentum strings is vanishing in the thermodynamic limit as

$$\frac{\tilde{Z}_{Neel}}{Z_{Neel}} \propto \frac{4}{\sqrt{\pi L}}. \quad (33)$$

Appendix A. Eigenstates with nonzero Néel overlap: eigenstates counting and string content

Here we prove that the total number of eigenstates with Néel nonzero overlap $Z_{Neel}(L)$ for a chain of length L is given as

$$Z_{Neel} = 2^{\frac{L}{2}-1} + \frac{1}{2}B\left(\frac{L}{2}, \frac{L}{4}\right) + 1. \quad (A.1)$$

For simplicity here we restrict ourselves to the situation with L divisible by four. The strategy to prove (A.1) is to count all the possible parity-invariant BGT quantum numbers configurations. Let us consider the sector with fixed number of particles M , and a generic string content $\mathcal{S} = \{s_1, s_2, \dots, s_M\}$. Here s_n is the number of n -strings, and one has the constraint $\sum_k k s_k = M$.

It is straightforward to check that total number of parity-invariant quantum number pairs $\mathcal{N}_n(L, \mathcal{S})$ in the n -string sector is given as

$$\mathcal{N}_n(L, \mathcal{S}) = \left\lfloor \frac{L}{2} - \frac{1}{2} \sum_{m=1}^M t_{nm} s_m \right\rfloor. \quad (A.2)$$

where $t_{nm} \equiv 2\text{Min}(n, m) - \delta_{n,m}$. The number of parity-invariant quantum number configurations (i.e., eigenstates) $\mathcal{N}(L, \mathcal{S})$ compatible with string content \mathcal{S} is obtained

by choosing in all the possible ways the parity-invariant quantum number pairs independently in each n -string sector, which implies that

$$\mathcal{N}(L, \mathcal{S}) = \prod_{m=1}^M B\left(\mathcal{N}_m, \left\lfloor \frac{s_m}{2} \right\rfloor\right). \quad (\text{A.3})$$

Here the product is because each string sector is treated independently, while the factor $1/2$ in $s_m/2$ is because since all quantum numbers are organized in pairs, only half of the quantum numbers have to be specified. Note that in each n -string sector only one zero momentum (i.e., zero quantum number) string is allowed, due to the Pauli principle. Moreover, s_m is odd (even) only if the zero momentum string is (not) present. The floor function $\lfloor \cdot \rfloor$ in (A.3) reflects that the quantum number of zero-momentum strings is fixed.

We now consider the string configurations with particle number $0 \leq \ell \leq M$ and fixed number of strings $1 \leq q \leq M/2$. Note that the maximul allowed string length is $M/2$ beacause of parity invariance. Note also that in determining q strings of different length are treated equally. Clearly, one has that $\sum_m s_m = q$. For a given fixed pair ℓ, q the total number of quantum number configurations is given as

$$\mathcal{N}'(L, \ell, q) = \sum_{\{\{s_m\} : \sum m s_m = \ell, \sum s_m = q\}} \mathcal{N}(L, \mathcal{S}), \quad (\text{A.4})$$

where the sum is over the content $\{s_m\}_{m=1}^M$ compatible with the constraints $\sum_m s_m = q$ and $\sum_m m s_m = \ell$. The strategy is to write a recursive relation for $\mathcal{N}'(L, \ell, q)$. To this purpose it is useful to consider the shifted string content \mathcal{S}' defined as

$$\mathcal{S}' \equiv \{s_{m+1}\} \quad \text{with } s_m \in \mathcal{S}, \forall m. \quad (\text{A.5})$$

Using the definition of t_{ij} , it is straightforward to derive that

$$t_{ij} = t_{i-1, j-1} + 2, \quad (\text{A.6})$$

which implies that $\mathcal{N}_n(L, \mathcal{S})$ (see (A.2)) satisfies the recursive equation

$$\mathcal{N}_n(L, \mathcal{S}) = \mathcal{N}_{n-1}(L - 2q, \mathcal{S}'). \quad (\text{A.7})$$

After substituting in (A.3) one obtains

$$\mathcal{N}(L, \mathcal{S}) = B\left(\mathcal{N}_1(L, \mathcal{S}), \left\lfloor \frac{s_1}{2} \right\rfloor\right) \mathcal{N}(L - 2q, \mathcal{S}'). \quad (\text{A.8})$$

Finally, after substituting (A.8) in (A.4), one obtains a recursive relation for $\mathcal{N}'(L, \ell, q)$ as

$$\mathcal{N}'(L, \ell, q) = \sum_{s=0}^{q-1} B\left(\frac{L}{2} - q + \left\lfloor \frac{s}{2} \right\rfloor, \left\lfloor \frac{s}{2} \right\rfloor\right) \mathcal{N}'(L - 2q, \ell - q, q - s), \quad (\text{A.9})$$

with the constraint that when $\ell = q$ one has

$$\mathcal{N}'(L, q, q) = B\left(\left\lfloor \frac{L - q}{2} \right\rfloor, \left\lfloor \frac{q}{2} \right\rfloor\right). \quad (\text{A.10})$$

This is obtained by observing that if $\ell = q$ only 1-strings are allowed and (A.2) gives $\mathcal{N}_n(L, \mathcal{S}) = \lfloor (L - q)/2 \rfloor$.

It is straightforward to check that for even q the ansatz

$$\mathcal{N}'(L, \ell, q) = \frac{q}{\ell} B\left(\frac{L-\ell}{2}, \frac{q}{2}\right) B\left(\frac{\ell}{2}, \frac{q}{2}\right), \quad (\text{A.11})$$

satisfies (A.9). For odd q the solution of (A.9) is

$$\mathcal{N}'(L, \ell, q) = \frac{\ell - q + 1}{\ell} B\left(\frac{L-\ell}{2}, \frac{q-1}{2}\right) B\left(\frac{\ell}{2}, \frac{q-1}{2}\right). \quad (\text{A.12})$$

The number of eigenstates in the sector with ℓ particles with nonzero Néel overlap $Z'_{\text{Neel}}(L, \ell)$ are obtained by summing over all possible values of q as

$$Z'_{\text{Neel}}(L, \ell) = \sum_{q=1}^{\ell} \mathcal{N}'(L, \ell, q). \quad (\text{A.13})$$

It is convenient to split the summation in (A.13) considering odd values of q and even q separately. For odd q one obtains

$$\sum_{k=0}^{\ell/2-1} \mathcal{N}'(L, \ell, 2k+1) = B\left(\frac{L}{2} - 1, \frac{\ell}{2} - 1\right), \quad (\text{A.14})$$

while for even q one has

$$\sum_{k=0}^{\ell/2} \mathcal{N}'(L, \ell, 2k) = B\left(\frac{L}{2} - 1, \frac{\ell}{2}\right). \quad (\text{A.15})$$

Putting everything together one obtains

$$Z'_{\text{Neel}}(L, \ell) = B\left(\frac{L}{2} - 1, \frac{\ell}{2} - 1\right) + B\left(\frac{L}{2} - 1, \frac{\ell}{2}\right). \quad (\text{A.16})$$

The total number of eigenstates with nonzero Néel overlap $Z_{\text{Neel}}(L)$ (cf. (A.1)) is obtained from (A.16) by summing over the allowed values of $\ell = 2k$ with $k = 0, 1, \dots, \ell/2$.

Note that the total number Z_{MG} of parity-invariant eigenstates having non zero overlap with the Majumdar-Ghosh state is obtained from Eq (A.16) replacing $\ell = L/2$, to obtain

$$Z_{MG} = B\left(\frac{L}{2} - 1, \frac{L}{4} - 1\right) + B\left(\frac{L}{2} - 1, \frac{L}{4}\right). \quad (\text{A.17})$$

Physically, this is due to the fact that the Majumdar-Ghosh state is invariant under $SU(2)$ rotations, which implies that only eigenstates with zero total spin $S = 0$ can have non zero overlap.

Appendix B. Excluding the zero-momentum strings

Here we demonstrate that the total number of eigenstates with nonzero Néel overlap, which do not contain zero-momentum strings, $\tilde{Z}_{\text{Neel}}(L)$ is given as

$$\tilde{Z}_{\text{Neel}}(L) = B\left(\frac{L}{2}, \frac{L}{4}\right). \quad (\text{B.1})$$

Given a generic M -particle eigenstate of the XXX chain, due to parity invariance, if one excludes the zero-momentum strings only n -strings with length $n \leq M/2$ are allowed.

Similarly, the string content is of the form $\tilde{\mathcal{S}} \equiv \{\tilde{s}_1, \dots, \tilde{s}_{M/2}\}$, i.e., $\tilde{s}_m = 0 \forall m > M/2$. Note that due to parity invariance and to the exclusion of the zero-momentum strings one has that \tilde{s}_m is always an even integer. Clearly one has $\sum_{m=1}^{M/2} m\tilde{s}_m = M$. The total number of parity-invariant quantum numbers $\tilde{\mathcal{N}}_n$ in the n -string sector is given as

$$\tilde{\mathcal{N}}_n(L, \tilde{\mathcal{S}}) = \frac{L}{2} - \frac{1}{2} \sum_{m=1}^{M/2} t_{nm} \tilde{s}_m. \quad (\text{B.2})$$

The proof now proceeds as in [Appendix A](#). One can define the total number of eigenstates with nonzero Néel overla in the sector with ℓ particles and q different strings as $\tilde{\mathcal{N}}'(L, \ell, q)$. Note that due to parity invariance and the exclusion of zero-momentum strings, q must be even. It is straightforward to show that $\tilde{\mathcal{N}}'(L, \ell, q)$ obeys the recursive relation

$$\tilde{\mathcal{N}}'(L, \ell, q) = \sum_{s=0}^{q/2-1} B\left(\frac{L}{2} - q + s, s\right) \tilde{\mathcal{N}}'\left(L - 2q, \frac{\ell - q}{2}, \frac{q}{2} - s\right), \quad (\text{B.3})$$

with the constraint

$$\tilde{\mathcal{N}}'(L, 1, 1) = \frac{L}{2} - 1. \quad (\text{B.4})$$

It is straightforward to check that the solution of [\(B.3\)](#) is given as

$$\tilde{\mathcal{N}}'(L, \ell, q) = \frac{L - 2\ell + 2}{L - \ell + 2} B\left(\frac{L - \ell}{2} + 1, q\right) B\left(\frac{\ell}{2} - 1, \frac{q}{2} - 1\right). \quad (\text{B.5})$$

After summing over the allowed values of $q = 2k$ with $k = 1, 2, \dots, \ell/2$ one obtains the total number of eigenstaets with nonzero Néel overlap at fixed number of particles ℓ $\tilde{\mathcal{Z}}'_{\text{Neel}}(L, \ell)$ as

$$\tilde{\mathcal{Z}}'_{\text{Neel}}(L, \ell) = B\left(\frac{L}{2}, \frac{\ell}{2}\right) - B\left(\frac{L}{2}, \frac{\ell}{2} - 1\right). \quad (\text{B.6})$$

Summing over ℓ one obtains [\(32\)](#). Similar to [\(A.17\)](#) the total number of eigenstates $\tilde{\mathcal{Z}}_{MG}$ with no zero-momentum strings having non-zero overlap with the Majumdar-Ghosh state is obtained from [\(B.6\)](#) replacing $\ell \rightarrow L/2$, to obtain

$$\tilde{\mathcal{Z}}_{MG} = B\left(\frac{L}{2}, \frac{L}{4}\right) - B\left(\frac{L}{2}, \frac{L}{4} - 1\right). \quad (\text{B.7})$$

Interestingly, using [\(A.17\)](#) and [\(B.7\)](#), one obtains that the ratio $\tilde{\mathcal{Z}}_{MG}/\mathcal{Z}_{MG}$ is given as

$$\frac{\tilde{\mathcal{Z}}_{MG}}{\mathcal{Z}_{MG}} = \frac{4}{4 + L}. \quad (\text{B.8})$$

Appendix C. Exact Néel and Majumdar-Ghosh overlaps for a small Heisenberg chain

In this section we provide exact diagonalization results for the overlap of both the Néel state and the Majumdar-Ghosh (MG) state with all the eigenstates of the Heisenberg spin chain with $L = 12$ sites. We also provide the corresponding results obtained using the string hypothesis and the overlap formulas [\(12\)](#) and [\(19\)](#), restricting ourselves to eigenstates with no zero-momentum strings.

Appendix C.1. Néel overlap

The overlaps between all the eigenstates of the Heisenberg spin chain and the Néel state are reported in Table C1. The first column in the Table shows the string content $\mathcal{S} \equiv \{s_1, \dots, s_M\}$, with M being the number of finite rapidities. The number of infinite rapidities $N_\infty = L/2 - M$ is also reported. Note that eigenstates containing infinite rapidities correspond to different S_z eigenvalue. The second column shows $2I_n^+$, with I_n the Bethe-Gaudin-Takahashi quantum number identifying the BGT rapidity of the n -string. Due to the parity invariance only the positive quantum numbers are reported. The total number of independent strings, i.e., $q \equiv \sum_j s_j$, is reported in the third column. The fourth column is the eigenstate's energy eigenvalue E . The last two columns show the squared Néel overlaps and the corresponding result obtained using the Bethe-Gaudin-Takahashi equations, respectively. In the last column only the case with no zero-momentum strings is considered. The deviations from the exact diagonalization results (digits with different colors) have to be attributed to the string hypothesis. Notice that the overlap between the Néel state and the $S_z = 0$ eigenstate in the sector with maximal total spin $S = L/2$ (first column in Table C1), is given analytically as $2/B(L, L/2)$, with $B(x, y)$ the Newton binomial.

Appendix C.2. Majumdar-Ghosh overlap

The overlap between the Heisenberg chain eigenstates with the Majumdar-Ghosh state are shown in Table C2. The conventions on the representation of the eigenstates is the same as in Table C1. Note that in contrast with the Néel state, only the eigenstates with zero total spin $S = 0$ have non zero overlap, i.e., no eigenstates with infinite rapidities are present, which reflect that the Majumdar-Ghosh state is unvariant under $SU(2)$ rotations.

- [1] P. Calabrese and P. Le Doussal, J. Stat. Mech. (2014) P05004.
- [2] V. Alba, arXiv:1507.06994.

Bethe states with nonzero Néel overlap ($L = 12$)						
String content	$2I_n^+$	q	E	$ \langle\lambda N\rangle ^2$ (exact)	$ \langle\lambda N\rangle ^2$ (BGT)	
6 inf	-	-	0	0.002164502165	0.002164502165	
{2,0} 4 inf	1_1	2	-3.918985947229	0.096183409244	0.096183409244	
	3_1		-3.309721467891	0.011288497947	0.011288497947	
	5_1		-2.284629676547	0.004542580506	0.004542580506	
	7_1		-1.169169973996	0.002752622983	0.002752622983	
	9_1		-0.317492934338	0.002116006203	0.002116006203	
{4,0,0,0} 2 inf	$1_1 3_1$	4	-7.070529325964	0.310133033838	0.310133033838	
	$1_1 5_1$		-5.847128730477	0.129277023687	0.129277023687	
	$1_1 7_1$		-4.570746557876	0.085992436024	0.085992436024	
	$3_1 5_1$		-5.153853093221	0.015256395523	0.015256395523	
	$3_1 7_1$		-3.916336243695	0.010091113504	0.010091113504	
	$5_1 7_1$		-2.817696043731	0.004059780228	0.004059780228	
{0,2,0,0} 2 inf	1_2	2	-1.905667167442	0.001207238321	0.001207245406	
	3_2		-1.368837200825	0.002340453815	0.002325724713	
	5_2		-0.681173793635	0.001921010489	0.001939001396	
{1,0,1,0} 2 inf	$0_1 0_3$	2	-2.668031843135	0.034959609810	-	
{6,0,0,0,0,0} 0 inf	$1_1 3_1 5_1$	6	-8.387390917445	0.153412152966	0.153412152966	
{2,2,0,0,0,0} 0 inf	$1_1 1_2$	4	-5.401838225870	0.040162686361	0.041042488913	
	$3_1 1_2$		-4.613929948329	0.004636541934	0.004730512604	
	$5_1 1_2$		-3.147465758841	0.001335522556	0.001337334035	
{3,0,1,0,0,0} 0 inf	$0_1 2_1 0_3$	4	-6.340207488736	0.052743525774	-	
	$0_1 4_1 0_3$		-5.203653009936	0.015022005621	-	
	$0_1 6_1 0_3$		-3.788693957250	0.011144489334	-	
{1,0,0,0,1,0} 0 inf	$0_1 0_5$	2	-2.444293750583	0.005887902992	-	
{0,0,2,0,0,0} 0 inf	1_3	2	-1.111855930538	0.001342476001	0.001384980817	
{0,1,0,1,0,0} 0 inf	$0_2 0_4$	2	-1.560671012472	0.000026982174	-	

Table C1. All Bethe states for $L = 12$ having nonzero overlap with the zero-momentum Néel state. The first column shows the string content of the Bethe states, including the number of infinite rapidities. The second and third column show $2I_n^+$, with I_n^+ the BGT quantum numbers identifying the different states, and the number q of independent strings. In the second column only the positive BGT numbers are shown. The fourth column is the Bethe state eigenenergy. Finally, the last two columns show the exact overlap with the Néel state and the approximate result obtained using the BGT equations. In the last column Bethe states containing zero-momentum strings are excluded. Deviations from the exact result (digits with different colors) are attributed to the string hypothesis.

Bethe states with nonzero Néel overlap ($L = 12$)					
String content	$2I_n^+$	q	E	$ \langle \lambda MG \rangle ^2$ (exact)	$ \langle \lambda MG \rangle ^2$ (BGT)
$\{6,0,0,0,0,0\}$	$1_1 3_1 5_1$	6	-8.387390917445	0.716615769224	0.716615769224
$\{2,2,0,0,0,0\}$	$1_1 1_2$	4	-5.401838225870	0.055624700196	0.054033366543
	$3_1 1_2$		-4.613929948329	0.005687428810	0.005582983043
	$5_1 1_2$		-3.147465758841	0.002107475934	0.002107086933
$\{3,0,1,0,0,0\}$	$0_1 2_1 0_3$	4	-6.340207488736	0.205891158647	-
	$0_1 4_1 0_3$		-5.203653009936	0.038832154450	-
	$0_1 6_1 0_3$		-3.788693957250	0.006019410923	-
$\{1,0,0,0,1,0\}$	$0_1 0_5$	2	-2.444293750583	0.000129601311	-
$\{0,0,2,0,0,0\}$	1_3	2	-1.111855930538	0.000011727787	0.000012785580
$\{0,1,0,1,0,0\}$	$0_2 0_4$	2	-1.560671012472	0.000330572718	-

Table C2. All Bethe states for $L = 12$ having nonzero overlap with the zero-momentum Majumdar-Ghosh (MG) state. The first column shows the string content of the Bethe states. The second and third column show $2I_n^+$, with I_n^+ the BGT quantum numbers identifying the different states, and the number q of independent strings. In the second column only the positive BGT numbers are shown. Note that, in contrast to Table C1 no states with infinite rapidities are present. The fourth column is the Bethe state eigenenergy. Finally, the last two columns show the exact overlap with the MG state and the approximate result obtained using the BGT equations. In the last column Bethe states containing zero-momentum strings are excluded. Deviations from the exact result (digits with different colors) are attributed to the string hypothesis.