

# Advanced Computational Physics - Exercise 1

Francesco Pederiva

## Scattering of H atoms on a Kr atom

In this exercise we will try to model and reproduce the total cross section of H-Kr (elastic) scattering, as measured in the paper by J.P. Toennies, W. Welz, and G. Wolf, J. Chem. Phys. **71**, 614 (1979). We will start assuming a simple form of the interaction:

$$v(r) = 4\epsilon \left[ \left( \frac{\sigma}{r} \right)^{12} - \left( \frac{\sigma}{r} \right)^6 \right],$$

where  $\epsilon = 5.9\text{meV}$ ,  $\sigma = 3.18\text{\AA}$ , and  $r$  is the interatomic H-Kr distance. Notice that in general the parameters in the potential should be *fitted* against the experiment.

In order to write a code that computes the desired cross section you will have to go through the following steps:<sup>1</sup>

1. Start writing a code solving the Schrödinger's equation for a simple one dimensional harmonic oscillator using the Numerov algorithm illustrated in class.

Use a dimensionless Hamiltonian:

$$-\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2$$

and plot the first 5 solutions (eigenvalues and eigenfunctions) in order to check the correctness of your code. Try in particular to check the dependence of your solution on the number of points in the mesh. [20 points]

---

<sup>1</sup>In the exercise report you will have to separately comment on each of the items in the list.

2. Repeat the same check for a 3-dimensional harmonic oscillator (use spherical coordinates) for values of the angular momentum  $\ell = 0, 1, 2$  and for 3 values of the principal quantum number for each  $\ell$ . [10 points]
3. Build a small program (that you will later transform into a function) to compute the Bessel functions  $j_\ell$  and  $n_\ell$  using the recursion formulae given in class:

$$j_{-1}(x) = \frac{\cos(x)}{x} \quad j_0(x) = \frac{\sin(x)}{x}$$

$$n_{-1}(x) = \frac{\sin(x)}{x} \quad n_0(x) = -\frac{\cos(x)}{x}$$

$$s_{\ell+1}(x) = \frac{2\ell+1}{x}s_\ell(x) - s_{\ell-1}(x).$$

Check the results against the exact values. [5 points]