

Jet Energy Loss at LHC

Vineet Kumar*

Nuclear Physics Division, Bhabha Atomic Research Center, Mumbai, India

Prashant Shukla

*Nuclear Physics Division, Bhabha Atomic Research Center, Mumbai, India and
Homi Bhabha National Institute, Anushakti Nagar, Mumbai, India*

(Dated: July 8, 2020)

In this work, the jet energy loss is analyzed using a monte carlo method. The data from LHC at $\sqrt{s_{NN}} = 2.76$ TeV and 5.02 TeV is used to extract the parameters for specific energy loss of jets inside QGP. Our calculations give good discription of the nuclear modification factor and asymmetry measurements at LHC.

PACS numbers: 12.38.Mh, 24.85.+p, 25.75.-q

Keywords: quark-gluon plasma, direct photon

I. INTRODUCTION

II. JET ENERGY LOSS

III. RESULTS AND DISCUSSIONS

IV. SUMMARY

-
- [1] J. Sollfrank, P. Huovinen, M. Kataja, P.V. Ruuskanen, M. Prakash and R. Venugopalan, Phys. Rev. C**55**, 392 (1997).

* vineetk@barc.gov.in




FIG. 1. (Color online) DiJet asymmetry measured by CMS compared with our calculations.




FIG. 2. (Color online) DiJet asymmetry measured by CMS compared with our calculations.




FIG. 3. (Color online) DiJet asymmetry measured by CMS compared with our calculations.




FIG. 4. (Color online) DiJet asymmetry measured by CMS compared with our calculations.