

Quarkonia production and dissociation in Pb+Pb collisions

Vineet Kumar^{1,2} and Prashant Shukla^{1,2,*}

¹*Nuclear Physics Division, Bhabha Atomic Research Center, Mumbai, India*

²*Homi Bhabha National Institute, Anushakti Nagar, Mumbai, India*

(Dated: June 16, 2015)

Abstract

We calculate the high p_T quarkonia production using NRQCD method. Different methods of quarkonia suppression are used to explain the high p_T quarkonia suppression observed by CMS in LHC

PACS numbers: 12.38.Mh, 24.85.+p, 25.75.-q

Keywords: quark-gluon plasma, quarkonia, suppression, regeneration

* pshukla@barc.gov.in




FIG. 1. (Color online) Differential production cross-section of J/ψ as a function of p_T compared with the CMS [?] data.

I. INTRODUCTION

Heavy-ion collisions at relativistic energies are performed to create and characterize quark gluon plasma (QGP), a phase of strongly-interacting matter at high energy density where quarks and gluons are no longer bound within hadrons. The quarkonia states (J/ψ and Υ) have been some of the most popular tools since their suppression was proposed as a signal of QGP formation [1]. The understanding of these probes has evolved substantially via measurements through three generations of experiments: the SPS (at CERN), RHIC (at BNL) and the LHC (at CERN) and by a great deal of theoretical activity. (For recent reviews see Refs. [2–4].) Quarkonia are produced early in the heavy-ion collisions and, if they evolve through the deconfined medium, their yields should be suppressed in comparison with those in pp collisions. The first such measurement was the ‘anomalous’ J/ψ suppression discovered at the SPS which was considered to be a hint of QGP formation. The RHIC measurements showed almost the same suppression at a much higher energy contrary to expectation [4, 5]. Such an observation was consistent with the scenario that, at higher collision energies, the expected greater suppression is compensated by J/ψ regeneration through recombination of two independently-produced charm quarks [6].

In this paper, we calculate J/ψ and Υ production and suppression

II. QUARKONIA PRODUCTION IN P+P COLLISIONS

In this section we describe the production of quarkonia at high transverse momenta in p+p collisions.

The factorization formalism of the NRQCD provides a theoretical framework for studying the heavy quarkonium production and decay. According to the NRQCD factorization formalism, the cross-section for direct production of a resonance H in a collision of particle A and B can be expressed as

$$d\sigma_{A+B \rightarrow H+X} = \sum_{a,b,n} \int dx_a dx_b G_{a/A}(x_a, \mu_F^2) G_{b/B}(x_b, \mu_F^2) \times d\sigma(a+b \rightarrow Q\bar{Q}(n)+X) < \mathcal{O}^H(n) > \quad (1)$$

where, $G_{a/A}(G_{b/B})$ is the parton distribution function (PDF) of the incoming parton $a(b)$ in the incident hadron $A(B)$, which depends on the momentum fraction $x_a(x_b)$ and the factorization scale μ_F as well as on the renormalization scale μ_R . However, as we have chosen $\mu_F = \mu_R$, in our case PDFs are function of x and μ_F only. The transverse mass of the resonance H is $m_T = \sqrt{p_T^2 + m_H^2}$, where $m_H \sim 2m_Q$ is the mass of resonance H . The short distance contribution $d\sigma(a+b \rightarrow Q\bar{Q}(n)+X)$ can be calculated within the framework of perturbative QCD (pQCD). On the other hand, $< \mathcal{O}^H(n) >$ (the state $n = {}^{2S+1}L_J^{[i]}$) are nonperturbative LDMEs and can be estimated on the basis of the comparison with experimental measurements.

The dominant processes in evaluating the differential yields of heavy mesons as a function of p_T are the $2 \rightarrow 2$ processes of the kind $g+q \rightarrow H+q$, $q+\bar{q} \rightarrow H+g$ and $g+g \rightarrow H+g$, where H refers to the heavy meson. We label the process generically as $a+b \rightarrow c+d$, where a and b are light incident partons, c refers to H and d is a light final-state parton. The differential cross-section for the short distance contribution i.e. the heavy quark pair production from the reaction of the type $a+b \rightarrow c+d$ can be written as [7]

$$\frac{d\sigma^{ab \rightarrow cd}}{dp_T dy} = \int dx_a G_{a/A}(x_a, \mu_F^2) G_{b/B}(x_b, \mu_F^2) \times 2p_T \frac{x_a x_b}{x_a - \frac{m_T}{\sqrt{s}} e^y} \frac{d\sigma}{d\hat{t}}(ab \rightarrow cd), \quad (2)$$

where, \sqrt{s} being the total energy in the centre-of-mass and y is the rapidity of the $Q\bar{Q}$ pair. In our numerical computation, we use CTEQ6M [8] for the parton distribution functions. The invariant differential cross-section is given by

$$\frac{d\sigma}{d\hat{t}} = \frac{|\mathcal{M}|^2}{16\pi\hat{s}^2}, \quad (3)$$

where \hat{s} and \hat{t} are the parton level Mandelstam variables. \mathcal{M} is the feynman amplitude for the process. Energy momentum conservation fixes value of the momentum fraction x_b as,

$$x_b = \frac{1}{\sqrt{s}} \frac{x_a \sqrt{s} m_T e^{-y} - m_H^2}{x_a \sqrt{s} - m_T e^y}. \quad (4)$$

The minimum value of x_a is

$$x_{a\min} = \frac{1}{\sqrt{s}} \frac{\sqrt{s} m_T e^y - m_H^2}{\sqrt{s} - m_T e^{-y}}. \quad (5)$$

The LDMEs are predicted to scale with a definite power of the relative velocity v of the heavy constituents inside $Q\bar{Q}$ bound states. In the limit $v \ll 1$, the production of quarkonium is based on the $^3S_1^{[1]}$ and $^3P_J^{[1]}$ ($J = 0,1,2$) CS states and $^1S_0^{[8]}$, $^3S_1^{[8]}$ and $^3P_J^{[8]}$ CO states. In our calculations, we used the expressions for the short distance CS cross-sections given in Refs. [9, 10] and the CO cross-sections given in Refs. [11, 12].

In this section we calculate the p_T distribution of J/ψ , $\psi(2S)$ and $\Upsilon(nS)$ mesons in $p-p$ collisions at LHC energies. For J/ψ production in $p-p$ collisions, three sources need to be considered: direct J/ψ production, feed-down contributions to the J/ψ from the decay of heavier charmonium states, predominantly from $\psi(2S)$, χ_{c0} , χ_{c1} and χ_{c2} and J/ψ from B hadron decays. The sum of the first two sources is called "prompt J/ψ " and the third source will be called " J/ψ from B ". On the other hand, $\psi(2S)$ has no significant feed-down contributions from higher mass states. We call this direct contribution as "prompt $\psi(2S)$ " to be consistent with the experiments. The other source to $\psi(2S)$ production is from B hadron decays and we call it " $\psi(2S)$ from B ". The sum of the prompt $J/\psi(\psi(2S))$ and $J/\psi(\psi(2S))$ from B will be called "inclusive $J/\psi(\psi(2S))$ ". Similar terminology is used for Υ states, only difference is that we do not have any contribution from open T mesons.

NRQCD provides a systematic procedure to compute any quantity as an expansion in the relative velocity v of the heavy quarks in the meson. For example, the wavefunction of the J/ψ meson (analogous expressions hold for the $\psi(2S)$, $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$) is written as

$$\begin{aligned} |J/\psi\rangle = & |Q\bar{Q}([{}^3S_1]_1)\rangle + \mathcal{O}(v)|Q\bar{Q}([{}^1S_0]_{8g})\rangle + \mathcal{O}(v^2)|Q\bar{Q}([{}^3S_1]_{8gg})\rangle \\ & + \mathcal{O}(v^1)|Q\bar{Q}([{}^3P_0]_{8g})\rangle + \mathcal{O}(v^1)|Q\bar{Q}([{}^3P_1]_{8g})\rangle + \mathcal{O}(v^1)|Q\bar{Q}([{}^3P_2]_{8g})\rangle + \dots \end{aligned} \quad (6)$$

The differential cross section for the direct production of J/ψ can be written as the sum

of the contributions,

$$\begin{aligned}
d\sigma(J/\psi) = & d\sigma(Q\bar{Q}([{}^3S_1]_1))\langle\mathcal{O}(Q\bar{Q}([{}^3S_1]_1) \rightarrow J/\psi)\rangle + d\sigma(Q\bar{Q}([{}^1S_0]_8))\langle\mathcal{O}(Q\bar{Q}([{}^1S_0]_8) \rightarrow J/\psi)\rangle \\
& + d\sigma(Q\bar{Q}([{}^3S_1]_8))\langle\mathcal{O}(Q\bar{Q}([{}^3S_1]_8) \rightarrow J/\psi)\rangle + d\sigma(Q\bar{Q}([{}^3P_0]_8))\langle\mathcal{O}(Q\bar{Q}([{}^3P_0]_8) \rightarrow J/\psi)\rangle \\
& + d\sigma(Q\bar{Q}([{}^3P_1]_8))\langle\mathcal{O}(Q\bar{Q}([{}^3P_1]_8) \rightarrow J/\psi)\rangle + d\sigma(Q\bar{Q}([{}^3P_2]_8))\langle\mathcal{O}(Q\bar{Q}([{}^3P_2]_8) \rightarrow J/\psi)\rangle + \dots,
\end{aligned}
\tag{7}$$

where the quantity in the brackets $[\]$ represents the angular momentum quantum numbers of the $Q\bar{Q}$ pair in the Fock expansion. The subscript on $[\]$ refers to the color structure of the $Q\bar{Q}$ pair, 1 being the color-singlet and 8 being the color-octet. The dots represent terms which contribute at higher powers of v . The short distance cross sections $d\sigma(Q\bar{Q})$ correspond to the production of a $Q\bar{Q}$ pair in a particular color and spin configuration, while the long distance matrix element $\langle\mathcal{O}(Q\bar{Q}) \rightarrow J/\psi\rangle$ corresponds to the probability of the $Q\bar{Q}$ state to convert to the quarkonium wavefunction. This probability includes any necessary prompt emission of soft gluons to prepare a color neutral system that matches onto the corresponding Fock component of the quarkonium wavefunction.

III. SUMMARY

IV. ACKNOWLEDGEMENT

The authors thank their CMS colleagues for the fruitful discussions, help and comments. Many of these results were presented at WHEPP and we acknowledge discussions with the participants of the meeting, in particular with D. Das, S. Datta, R. Gavai, S. Gupta and R. Sharma. The work of RV was performed under the auspices of the US Department of Energy, Lawrence Livermore National Laboratory, Contract DE-AC52-07NA27344.

-
- [1] T. Matsui and H. Satz, “ J/ψ Suppression by Quark-Gluon Plasma Formation”, Phys. Lett. B **178**, 416 (1986).
 - [2] J. Schukraft, “Heavy Ion Physics at the LHC: What’s new ? What’s next ?”, arXiv:1311.1429 [hep-ex].

- [3] L. Kluberg and H. Satz, “Color Deconfinement and Charmonium Production in Nuclear Collisions,” arXiv:0901.3831 [hep-ph].
- [4] N. Brambilla, S. Eidelman, B. K. Heltsley, R. Vogt, G. T. Bodwin, E. Eichten, A. D. Frawley and A. B. Meyer *et al.*, “Heavy quarkonium: progress, puzzles, and opportunities,” Eur. Phys. J. C **71**, 1534 (2011).
- [5] A. Adare *et al.* [PHENIX Collaboration], “ J/ψ suppression at forward rapidity in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” Phys. Rev. C **84**, 054912 (2011).
- [6] A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, “Statistical hadronization of charm in heavy ion collisions at SPS, RHIC and LHC,” Phys. Lett. B **571**, 36 (2003).
- [7] J. F. Owens, “Large Momentum Transfer Production of Direct Photons, Jets, and Particles,” Rev. Mod. Phys. **59**, 465 (1987).
- [8] H. L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin and C.-P. Yuan, “New parton distributions for collider physics,” Phys. Rev. D **82**, 074024 (2010), [arXiv:1007.2241 [hep-ph]].
- [9] R. Baier and R. Ruckl, “Hadronic Collisions: A Quarkonium Factory,” Z. Phys. C **19**, 251 (1983).
- [10] R. Gastmans, W. Troost and T. T. Wu, “Production of Heavy Quarkonia From Gluons,” Nucl. Phys. B **291**, 731 (1987).
- [11] P. L. Cho and A. K. Leibovich, “Color octet quarkonia production,” Phys. Rev. D **53**, 150 (1996), [hep-ph/9505329].
- [12] P. L. Cho and A. K. Leibovich, “Color octet quarkonia production. 2.,” Phys. Rev. D **53**, 6203 (1996), [hep-ph/9511315].