


# Verifying the Klein-Nishina prediction using Compton scattering of 661.7 keV photons in NaI scintillators

Vinh Q. Tran

# Kinematics of Compton scattering & Klein-Nishina differential cross section

---




# Kinematics of Compton scattering & Klein-Nishina differential cross section

---

- Energy of the scattered photon

$$E'_\gamma = \frac{E_\gamma}{1 + \frac{E_\gamma}{m_e c^2} (1 - \cos \theta)}$$




# Kinematics of Compton scattering & Klein-Nishina differential cross section

- Energy of the scattered photon

$$E'_\gamma = \frac{E_\gamma}{1 + \frac{E_\gamma}{m_e c^2} (1 - \cos \theta)}$$

- Klein-Nishina formula

$$\frac{d\sigma_{KN}}{d\Omega} = \frac{r_e^2}{2} \left( \frac{\lambda}{\lambda'} \right)^2 \left[ \frac{\lambda}{\lambda'} + \frac{\lambda'}{\lambda} - \sin^2 \theta \right]$$



# Kinematics of Compton scattering & Klein-Nishina differential cross section

- Energy of the scattered photon


$$E'_\gamma = \frac{E_\gamma}{1 + \frac{E_\gamma}{m_e c^2} (1 - \cos \theta)}$$

- Klein-Nishina formula

$$\frac{d\sigma_{KN}}{d\Omega} = \frac{r_e^2}{2} \left( \frac{\lambda}{\lambda'} \right)^2 \left[ \frac{\lambda}{\lambda'} + \frac{\lambda'}{\lambda} - \sin^2 \theta \right]$$


- Classical Thompson formula

$$\frac{d\sigma_T}{d\Omega} = r_e^2 \frac{1 + \cos^2 \theta}{2}$$




## The apparatus arrangement

---




## The apparatus arrangement




$\theta \in [0^\circ, 135^\circ]$  (at  $15^\circ$  intervals)

## The apparatus arrangement




$\theta \in [0^\circ, 135^\circ]$  (at  $15^\circ$  intervals)






## Circuit design & Coincidence mode



## Circuit design & Coincidence mode


- Coincidence mode registers only events detected within  $\tau = 2 \mu s$ .



## Circuit design & Coincidence mode


- Coincidence mode registers only events detected within  $\tau = 2 \mu s$ .
- Systematic error from background detection

$$n_F = 2\tau n_r n_s$$




## Circuit design & Coincidence mode

- Coincidence mode registers only events detected within  $\tau = 2 \mu s$ .
- Systematic error from background detection
$$n_F = 2\tau n_r n_s$$
- Setting threshold for the discriminator at 60 mV help reduce background rate to  $< 1$  Hz, while Compton scattering rates remain at  $\sim 20$  Hz.



# Cs137 energy spectrum in NaI scintillator



## MCA calibration

---


- Channel-energy conversion relation

$$E_i = \alpha C_i + b$$


# MCA calibration

- Channel-energy conversion relation



$$E_i = \alpha C_i + b$$



(Heath 1964)





# Spectrum analysis







## Spectrum analysis

- Apply a Gaussian filter ( $\sigma = 5$  channels) to smoothen the data.





## Spectrum analysis

- Apply a Gaussian filter ( $\sigma = 5$  channels) to smoothen the data.
- Find the local minima and maxima with derivative-base peak finding algorithms.



## Spectrum analysis



- Apply a Gaussian filter ( $\sigma = 5$  channels) to smoothen the data.
- Find the local minima and maxima with derivative-base peak finding algorithms.
- Fit peaks of interest using Gaussian functions (with optional offset).



## Spectrum analysis


- Apply a Gaussian filter ( $\sigma = 5$  channels) to smoothen the data.
- Find the local minima and maxima with derivative-base peak finding algorithms.
- Fit peaks of interest using Gaussian functions (with optional offset).
- For Compton scattering data, use Poisson-based likelihood approach

$$\chi^2 = - \sum 2 \log \frac{\hat{y}_i^{y_i} e^{-\hat{y}_i}}{y_i!} + \log 2\pi y_i$$



## Calibration result

- Example: Recoil scintillator, March 4<sup>th</sup> 2025




## Calibration result

- Example: Recoil scintillator, March 4<sup>th</sup> 2025
- Channel-energy conversion relation


$$E_i = \alpha C_i + b$$

$$\alpha = 0.390 \pm 0.014 \text{ keV}$$

$$b = -10.5 \pm 6.2 \text{ keV}$$




## Energy-angle dependency



## Energy-angle dependency

- The energies of the incident photons are well-reconstructed. Systematic errors of  $\sigma_{E_\gamma}^{\text{sys}} \sim 10$  keV.






## Energy-angle dependency

- The energies of the incident photons are well-reconstructed. Systematic errors of  $\sigma_{E_\gamma}^{\text{sys}} \sim 10$  keV.
- The energies of the scattered photons and recoil electrons follow as expected from the Compton theory

$$E'_\gamma = \frac{E_\gamma}{1 + \frac{E_\gamma}{m_e c^2} (1 - \cos \theta)}$$
$$K'_e = E_\gamma - E'_\gamma$$



## Energy-angle dependency

- The energies of the incident photons are well-reconstructed. Systematic errors of  $\sigma_{E_\gamma}^{\text{sys}} \sim 10$  keV.
- The energies of the scattered photons and recoil electrons follow as expected from the Compton theory


$$E'_\gamma = \frac{E_\gamma}{1 + \frac{E_\gamma}{m_e c^2} (1 - \cos \theta)}$$

$$K'_e = E_\gamma - E'_\gamma$$


- Initial angle offset,  $\cos \theta \rightarrow \cos(\theta - \theta_0)$

$$\theta_0 = 2.56 \pm 0.71^\circ$$

$$\sigma_{\theta_0} \sim \sigma_\theta^{\text{sys}} \sim 0.5^\circ$$




## Scattering rate



## Scattering rate

- The scattering rates follow the Klein-Nishina formula well (except for the case of forward scattering)

$$\frac{d\sigma_{KN}}{d\Omega} = \frac{r_e^2}{2} \left( \frac{\lambda}{\lambda'} \right)^2 \left[ \frac{\lambda}{\lambda'} + \frac{\lambda'}{\lambda} - \sin^2 \theta \right]$$




## Scattering rate

- The scattering rates follow the Klein-Nishina formula well (except for the case of forward scattering)

$$\frac{d\sigma_{KN}}{d\Omega} = \frac{r_e^2}{2} \left( \frac{\lambda}{\lambda'} \right)^2 \left[ \frac{\lambda}{\lambda'} + \frac{\lambda'}{\lambda} - \sin^2 \theta \right]$$

- The classical Thompson description fail to describe the observed data

$$\frac{d\sigma_T}{d\Omega} = r_e^2 \frac{1 + \cos^2 \theta}{2}$$



## Scattering rate


- The scattering rates follow the Klein-Nishina formula well (except for the case of forward scattering)

$$\frac{d\sigma_{KN}}{d\Omega} = \frac{r_e^2}{2} \left( \frac{\lambda}{\lambda'} \right)^2 \left[ \frac{\lambda}{\lambda'} + \frac{\lambda'}{\lambda} - \sin^2 \theta \right]$$

- The classical Thompson description fail to describe the observed data

$$\frac{d\sigma_T}{d\Omega} = r_e^2 \frac{1 + \cos^2 \theta}{2}$$

- Quantum relativistic treatment is necessary for the energy regime of  $E_\gamma \sim 1$  MeV.



## Conclusion

- Compton kinematics is confirmed!
- The quantum relativistic treatment of the Klein-Nishina prediction is necessary for the measured energy regime  $E_\gamma \sim 1$  MeV.

