

DETERMINANTS

ELECTRONIC VERSION OF LECTURE

HoChiMinh City University of Technology
Faculty of Applied Science, Department of Applied Mathematics



HCMC — 2020.

OUTLINE

1 DETERMINANTS

OUTLINE

- 1 DETERMINANTS
- 2 RANK OF A MATRIX BY MEANS OF DETERMINANTS

OUTLINE

- 1 DETERMINANTS
- 2 RANK OF A MATRIX BY MEANS OF DETERMINANTS
- 3 INVERSE OF AN MATRIX

OUTLINE

- 1 DETERMINANTS
- 2 RANK OF A MATRIX BY MEANS OF DETERMINANTS
- 3 INVERSE OF AN MATRIX
- 4 REAL-WORLD PROBLEMS

OUTLINE

- 1 DETERMINANTS
- 2 RANK OF A MATRIX BY MEANS OF DETERMINANTS
- 3 INVERSE OF AN MATRIX
- 4 REAL-WORLD PROBLEMS
- 5 MATLAB

DEFINITION 1.1

If $A = (a_{ij})$ is a square matrix, then the *determinant* of A is a *number*. We denote it by $\det(A)$ or $|A|$.

DEFINITION 1.1

If $A = (a_{ij})$ is a square matrix, then the *determinant* of A is a *number*. We denote it by $\det(A)$ or $|A|$.

So

$$\det: M_n(K) \rightarrow K$$

$$A \rightarrow \det A.$$

DEFINITION 1.2

If $A = (a_{ij})_{n \times n}$ is a *square matrix*, then the *minor of entry a_{ij}* is denoted by M_{ij} and is defined to be the determinant of the submatrix of order $(n - 1)$ that remains after the *i -th row* and *j -column* are deleted from A .

$$|A| = \begin{vmatrix} a_{11} & \dots & a_{1(j-1)} & a_{1j} & a_{1(j+1)} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{(i-1)1} & \dots & a_{(i-1)(j-1)} & a_{(i-1)j} & a_{(i-1)(j+1)} & \dots & a_{(i-1)n} \\ a_{i1} & \dots & a_{i(j-1)} & a_{ij} & a_{i(j+1)} & \dots & a_{in} \\ a_{(i+1)1} & \dots & a_{(i+1)(j-1)} & a_{(i+1)j} & a_{(i+1)(j+1)} & \dots & a_{(i+1)n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{n(j-1)} & a_{nj} & a_{n(j+1)} & \dots & a_{nn} \end{vmatrix}_{n \times n}$$

$$|A| = \begin{vmatrix} a_{11} & \dots & a_{1(j-1)} & a_{1j} & a_{1(j+1)} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{(i-1)1} & \dots & a_{(i-1)(j-1)} & a_{(i-1)j} & a_{(i-1)(j+1)} & \dots & a_{(i-1)n} \\ a_{i1} & \dots & a_{i(j-1)} & a_{ij} & a_{i(j+1)} & \dots & a_{in} \\ a_{(i+1)1} & \dots & a_{(i+1)(j-1)} & a_{(i+1)j} & a_{(i+1)(j+1)} & \dots & a_{(i+1)n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{n(j-1)} & a_{nj} & a_{n(j+1)} & \dots & a_{nn} \end{vmatrix}_{n \times n}$$

$$|A| = \begin{vmatrix} a_{11} & \dots & a_{1(j-1)} & a_{1j} & a_{1(j+1)} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{(i-1)1} & \dots & a_{(i-1)(j-1)} & a_{(i-1)j} & a_{(i-1)(j+1)} & \dots & a_{(i-1)n} \\ a_{i1} & \dots & a_{i(j-1)} & a_{ij} & a_{i(j+1)} & \dots & a_{in} \\ a_{(i+1)1} & \dots & a_{(i+1)(j-1)} & a_{(i+1)j} & a_{(i+1)(j+1)} & \dots & a_{(i+1)n} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{n(j-1)} & a_{nj} & a_{n(j+1)} & \dots & a_{nn} \end{vmatrix}_{n \times n}$$

$$M_{ij} =$$

$$\begin{vmatrix} a_{11} & \cdots & a_{1(j-1)} & a_{1(j+1)} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a_{(i-1)1} & \cdots & a_{(i-1)(j-1)} & a_{(i-1)(j+1)} & \cdots & a_{(i-1)n} \\ a_{(i+1)1} & \cdots & a_{(i+1)(j-1)} & a_{(i+1)(j+1)} & \cdots & a_{(i+1)n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{n(j-1)} & a_{n(j+1)} & \cdots & a_{nn} \end{vmatrix}_{(n-1) \times (n-1)}$$

$$M_{ij} =$$

$$\begin{vmatrix} a_{11} & \cdots & a_{1(j-1)} & a_{1(j+1)} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a_{(i-1)1} & \cdots & a_{(i-1)(j-1)} & a_{(i-1)(j+1)} & \cdots & a_{(i-1)n} \\ a_{(i+1)1} & \cdots & a_{(i+1)(j-1)} & a_{(i+1)(j+1)} & \cdots & a_{(i+1)n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{n(j-1)} & a_{n(j+1)} & \cdots & a_{nn} \end{vmatrix}_{(n-1) \times (n-1)}$$

DEFINITION 1.3

If $A = (a_{ij})_{n \times n}$ is a *square matrix*, then the number $C_{ij} = (-1)^{i+j} M_{ij}$ is called the *cofactor of entry* a_{ij} .

DEFINITION 1.4

If A is an $n \times n$ matrix, then the number obtained by multiplying the entries in any **row or column** of A by the corresponding cofactors and adding the resulting products is called the **determinant of A** , and the sums themselves are called **cofactor expansion of A** . That is,

$$\det(A) = \sum_{j=1}^n a_{ij}C_{ij} = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in}$$

$$\det(A) = \sum_{i=1}^n a_{ij}C_{ij} = a_{1j}C_{1j} + a_{2j}C_{2j} + \dots + a_{nj}C_{nj}$$

COFACTOR EXPANSION ALONG THE FIRST ROW

$$\begin{aligned}
 \det(A) &= \begin{vmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{nj} & \dots & a_{nn} \end{vmatrix} \\
 &= \sum_{j=1}^n a_{1j} C_{1j} = \sum_{j=1}^n a_{1j} \cdot (-1)^{1+j} M_{1j}.
 \end{aligned}$$

1 $n = 1, A = (a_{11}) \Rightarrow |A| = a_{11}.$

① $n = 1, A = (a_{11}) \Rightarrow |A| = a_{11}.$

② $n = 2, A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$

$$\Rightarrow |A| = (-1)^{1+1} a_{11} M_{11} + (-1)^{1+2} a_{12} M_{12} = a_{11} a_{22} - a_{12} a_{21}.$$

$$\textcircled{1} \quad n = 1, A = (a_{11}) \Rightarrow |A| = a_{11}.$$

$$\textcircled{2} \quad n = 2, A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

$$\Rightarrow |A| = (-1)^{1+1} a_{11} M_{11} + (-1)^{1+2} a_{12} M_{12} = a_{11} a_{22} - a_{12} a_{21}.$$

$$\textcircled{3} \quad n = 3, A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

$$\Rightarrow |A| = (-1)^{1+1} a_{11} M_{11} + (-1)^{1+2} a_{12} M_{12} + (-1)^{1+3} a_{13} M_{13}$$

$$= (-1)^{1+1} a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} + (-1)^{1+2} a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}$$

$$+ (-1)^{1+3} a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}.$$

EXAMPLE 1.1

Find the determinant $\det A$ of $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 2 & 1 \\ 3 & 1 & 5 \end{pmatrix}$

EXAMPLE 1.1

Find the determinant $\det A$ of $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 2 & 1 \\ 3 & 1 & 5 \end{pmatrix}$

Solution. Cofactor expansion along the first row:

$$|A| = 1.C_{11} + 2.C_{12} + 3.C_{13}.$$

$$C_{11} = (-1)^{1+1} \begin{vmatrix} 2 & 1 \\ 1 & 5 \end{vmatrix} = 2 \times 5 - 1 \times 1 = 9,$$

$$C_{12} = (-1)^{1+2} \begin{vmatrix} 4 & 1 \\ 3 & 5 \end{vmatrix} = -(4 \times 5 - 1 \times 3) = -17,$$

$$C_{13} = (-1)^{1+3} \begin{vmatrix} 4 & 2 \\ 3 & 1 \end{vmatrix} = 4 \times 1 - 2 \times 3 = -2.$$

Therefore, $|A| = 1 \times 9 + 2 \times (-17) + 3 \times (-2) = -31.$

SMART CHOICE OF ROW OR COLUMN

SMART CHOICE OF ROW OR COLUMN

We can find determinant using cofactor expansion along **any row**.

$$\det A = \begin{vmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \mathbf{a_{i1}} & \dots & \mathbf{a_{ij}} & \dots & \mathbf{a_{in}} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{nj} & \dots & a_{nn} \end{vmatrix} = \sum_{j=1}^n \mathbf{a_{ij}} C_{ij}$$

Determinant also can be found using cofactor expansion along **any column**.

$$\det A = \begin{vmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{nj} & \dots & a_{nn} \end{vmatrix} = \sum_{i=1}^n a_{ij} C_{ij}$$

Determinant also can be found using cofactor expansion along **any column**.

$$\det A = \begin{vmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ a_{n1} & \dots & a_{nj} & \dots & a_{nn} \end{vmatrix} = \sum_{i=1}^n a_{ij} C_{ij}$$

It will be easiest to use cofactor expansion along the row or column which has **the most zeros**.

EXAMPLE 1.2

Evaluate $\det A$ where $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 2 & 0 \\ 3 & 1 & 5 \end{pmatrix}$

EXAMPLE 1.2

Evaluate $\det A$ where $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 2 & 0 \\ 3 & 1 & 5 \end{pmatrix}$

Solution. Cofactor expansion along the second row:

$$\begin{aligned} |A| &= 0.C_{21} + 2.C_{22} + 0.C_{23} \\ &= 2.(-1)^{2+2} \begin{vmatrix} 1 & 3 \\ 3 & 5 \end{vmatrix} = 2(1 \times 5 - 3 \times 3) = -8. \end{aligned}$$

EXAMPLE 1.3

Evaluate $\det A$ where $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 0 \\ 3 & 1 & 0 \end{pmatrix}$

EXAMPLE 1.3

Evaluate $\det A$ where $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 0 \\ 3 & 1 & 0 \end{pmatrix}$

Solution. Cofactor expansion along the third column

$$\begin{aligned} |A| &= 3.C_{13} + 0.C_{23} + 0.C_{33} \\ &= 3.(-1)^{1+3} \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} = 3(2 \times 1 - 1 \times 3) = -3. \end{aligned}$$

HOW AN ELEMENTARY ROW (OR COLUMN) OPERATION ON A SQUARE MATRIX AFFECTS THE VALUE OF ITS DETERMINANT

HOW AN ELEMENTARY ROW (OR COLUMN) OPERATION ON A SQUARE MATRIX AFFECTS THE VALUE OF ITS DETERMINANT

- 1 If $A \xrightarrow{r_i \leftrightarrow r_j (c_i \leftrightarrow c_j)} B$ then $\det B = -\det A$.

HOW AN ELEMENTARY ROW (OR COLUMN) OPERATION ON A SQUARE MATRIX AFFECTS THE VALUE OF ITS DETERMINANT

- 1 If $A \xrightarrow{r_i \leftrightarrow r_j (c_i \leftrightarrow c_j)} B$ then $\det B = -\det A$.
- 2 If $A \xrightarrow{r_i \rightarrow \lambda r_i (c_i \rightarrow \lambda c_i)} B$ then $\det B = \lambda \det A$ where $\lambda \neq 0$.

HOW AN ELEMENTARY ROW (OR COLUMN) OPERATION ON A SQUARE MATRIX AFFECTS THE VALUE OF ITS DETERMINANT

- 1 If $A \xrightarrow{r_i \leftrightarrow r_j (c_i \leftrightarrow c_j)} B$ then $\det B = -\det A$.
- 2 If $A \xrightarrow{r_i \rightarrow \lambda r_i (c_i \rightarrow \lambda c_i)} B$ then $\det B = \lambda \det A$ where $\lambda \neq 0$.
- 3 If $A \xrightarrow{r_i \rightarrow r_i + \lambda \cdot r_j (c_i \rightarrow c_i + \lambda c_j)} B$ then
$$\det B = \det A, \forall \lambda \in K$$

COROLLARY 1.1

- 1 If A is a square matrix with *2 equal rows or 2 equal columns* then $\det(A) = 0$.

COROLLARY 1.1

- ① If A is a square matrix with *2 equal rows or 2 equal columns* then $\det(A) = 0$. $A \xrightarrow{r_i \leftrightarrow r_j (c_i \leftrightarrow c_j)} A$ where i, j are 2 equal rows or 2 equal columns $\det A = -\det A \Rightarrow \det A = 0$.

COROLLARY 1.1

- ① If A is a square matrix with **2 equal rows or 2 equal columns** then $\det(A) = 0$. $A \xrightarrow{r_i \leftrightarrow r_j (c_i \leftrightarrow c_j)} A$ where i, j are 2 equal rows or 2 equal columns $\det A = -\det A \Rightarrow \det A = 0$.
- ② If A is a square matrix with **2 proportional rows or 2 proportional columns** then $\det(A) = 0$.

COROLLARY 1.1

- ① If A is a square matrix with **2 equal rows or 2 equal columns** then $\det(A) = 0$. $A \xrightarrow{r_i \leftrightarrow r_j (c_i \leftrightarrow c_j)} A$ where i, j are 2 equal rows or 2 equal columns $\det A = -\det A \Rightarrow \det A = 0$.
- ② If A is a square matrix with **2 proportional rows or 2 proportional columns** then $\det(A) = 0$. Since $A \xrightarrow{r_i \rightarrow \lambda r_i (c_i \rightarrow \lambda c_i)} B$ where $\lambda \neq 0$ is the ratio of 2 rows or 2 columns, $\det B = \lambda \det A$, where $\det B = 0 \Rightarrow \det A = 0$.

EXAMPLE 1.4

Use Row Reduction to evaluate the determinant

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix}$$

EXAMPLE 1.4

Use Row Reduction to evaluate the determinant

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix}$$

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix} \xrightarrow{r_2 \rightarrow r_2 - r_1}$$

EXAMPLE 1.4

Use Row Reduction to evaluate the determinant

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix}$$

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix} \xrightarrow[r_2 \rightarrow r_2 - r_1]{=} \begin{vmatrix} 2 & 3 & -4 \\ 1 & -8 & 6 \\ 5 & 4 & 3 \end{vmatrix} \xrightarrow[r_3 \rightarrow r_3 - 5r_2]{r_1 \rightarrow r_1 - 2r_2}$$

EXAMPLE 1.4

Use Row Reduction to evaluate the determinant

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix}$$

$$\begin{vmatrix} 2 & 3 & -4 \\ 3 & -5 & 2 \\ 5 & 4 & 3 \end{vmatrix} \xrightarrow[r_2 \rightarrow r_2 - r_1]{=} \begin{vmatrix} 2 & 3 & -4 \\ 1 & -8 & 6 \\ 5 & 4 & 3 \end{vmatrix} \begin{matrix} r_1 \rightarrow r_1 - 2r_2 \\ r_3 \rightarrow r_3 - 5r_2 \end{matrix}$$

$$\begin{vmatrix} 0 & 19 & -16 \\ 1 & -8 & 6 \\ 0 & 44 & -27 \end{vmatrix} \quad \text{Cofactor expansion along the first column}$$

$$= 1.(-1)^{2+1} \cdot \begin{vmatrix} 19 & -16 \\ 44 & -27 \end{vmatrix} = -191.$$

DETERMINANT OF A MATRIX PRODUCT

THEOREM 1.1

If A, B are square matrices of the same size, then

$$\det(AB) = \det(A).\det(B) \quad (1)$$

EXAMPLE 1.5

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & -2 & 6 \\ 2 & 8 & 9 \end{pmatrix}, B = \begin{pmatrix} 7 & 8 & 9 \\ 4 & -3 & 6 \\ -1 & 2 & 3 \end{pmatrix}$$

$$AB = \begin{pmatrix} 12 & 8 & 30 \\ 14 & 50 & 42 \\ 37 & 10 & 93 \end{pmatrix}$$

Verify that

$$\det(A).\det(B) = (-6).(-246) = \det(AB) = 1476$$

COROLLARY 1.2

If A, B are square matrices of the same size

❶ $\det(A^k) = (\det A)^k.$

COROLLARY 1.2

If A, B are square matrices of the same size

$$\textcircled{1} \quad \det(A^k) = (\det A)^k. \text{ Indeed, } \det(A^k) = \det(\underbrace{A.A \dots A}_{k \text{ times}}) = \underbrace{\det A. \det A \dots \det A}_{k \text{ times}} = (\det A)^k.$$

COROLLARY 1.2

If A, B are square matrices of the same size

- ① $\det(A^k) = (\det A)^k$. Indeed, $\det(A^k) = \det(\underbrace{A.A \dots A}_{k \text{ times}}) = \underbrace{\det A. \det A \dots \det A}_{k \text{ times}} = (\det A)^k$.
- ② $\det(\alpha AB) = \alpha^n. \det A. \det B$.

COROLLARY 1.2

If A, B are square matrices of the same size

$$\textcircled{1} \quad \det(A^k) = (\det A)^k. \text{ Indeed, } \det(A^k) = \det(\underbrace{A.A \dots A}_{k \text{ times}}) = \underbrace{\det A. \det A \dots \det A}_{k \text{ times}} = (\det A)^k.$$

$$\textcircled{2} \quad \det(\alpha AB) = \alpha^n. \det A. \det B.$$

Indeed,

$$\det(\alpha AB) = \det(\alpha A). \det B = \underbrace{\alpha. \alpha \dots \alpha}_{n \text{ times}} \det A. \det B$$

EXAMPLE 1.6

Evaluate $\det(X)$ if X satisfies

$$\begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{pmatrix} X = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & -1 \\ 3 & 5 & 2 \end{pmatrix}$$

EXAMPLE 1.6

Evaluate $\det(X)$ if X satisfies

$$\begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{pmatrix} X = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & -1 \\ 3 & 5 & 2 \end{pmatrix}$$

We have $\begin{vmatrix} 1 & 2 & 1 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{vmatrix} \cdot \det(X) = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & -1 \\ 3 & 5 & 2 \end{vmatrix}$

$$\Rightarrow$$

EXAMPLE 1.6

Evaluate $\det(X)$ if X satisfies

$$\begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{pmatrix} X = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & -1 \\ 3 & 5 & 2 \end{pmatrix}$$

We have $\begin{vmatrix} 1 & 2 & 1 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{vmatrix} \cdot \det(X) = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & -1 \\ 3 & 5 & 2 \end{vmatrix}$

$$\Rightarrow 1 \cdot \det(X) = 3 \Rightarrow \det(X) = 3.$$

EXAMPLE 1.7

If $A = \begin{pmatrix} -1 & 0 & 0 \\ 2 & 1 & 0 \\ 4 & 3 & 1 \end{pmatrix}$, then evaluate $\det(A^{2011})$.

EXAMPLE 1.7

If $A = \begin{pmatrix} -1 & 0 & 0 \\ 2 & 1 & 0 \\ 4 & 3 & 1 \end{pmatrix}$, then evaluate $\det(A^{2011})$.

We have

$$\det(A^{2011}) = (\det A)^{2011} = (-1)^{2011} = -1.$$

EXAMPLE 1.8

If $A = \begin{pmatrix} 3 & -2 & 6 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 2 & 5 \\ 1 & -2 & 7 \end{pmatrix}$, then evaluate $\det(2AB)$.

EXAMPLE 1.8

If $A = \begin{pmatrix} 3 & -2 & 6 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 2 & 5 \\ 1 & -2 & 7 \end{pmatrix}$, then evaluate $\det(2AB)$.

We have

$$\det(2AB) = 2^3 \cdot \det A \cdot \det B = 8 \times 3 \times 2 = 48.$$

DEFINITION 2.1

Let A be an $m \times n$ matrix. *A minor of A of order k is a determinant of a $k \times k$ submatrix of A .*

DEFINITION 2.1

Let A be an $m \times n$ matrix. *A minor of A of order k is a determinant of a $k \times k$ submatrix of A .*

EXAMPLE 2.1

Find the minors of order 3 of the matrix

$$A = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 2 & 4 & 2 \\ 0 & 2 & 2 & 1 \end{pmatrix}$$

SOLUTION. We obtain the determinants of order 3 by keeping all the rows and deleting one column from A . So there are 4 different minors of order 3. We compute one of them to illustrate:

$$\begin{vmatrix} 1 & 0 & 2 \\ 0 & 2 & 4 \\ 0 & 2 & 2 \end{vmatrix} = 1 \times (-4) + 2 \times 0 = -4.$$

SOLUTION. We obtain the determinants of order 3 by keeping all the rows and deleting one column from A . So there are 4 different minors of order 3. We compute one of them to illustrate:

$$\begin{vmatrix} 1 & 0 & 2 \\ 0 & 2 & 4 \\ 0 & 2 & 2 \end{vmatrix} = 1 \times (-4) + 2 \times 0 = -4.$$

The minors of order 3 are called **the maximal minors of A** , since there are no 4×4 submatrices of A .

PROPOSITION 2.1

*Let A be an $m \times n$ matrix. The rank of A is **maximal order of a non-zero minor of A** .*

COMPUTING THE RANK

Start with the minors of maximal order k . If there is one that is non-zero then $\text{rank}(A) = k$. If all maximal minors are zero, then $\text{rank}(A) < k$, and we continue with the minors of order $k - 1$ and so on, until we find a minor that is non-zero. If all minors of order 1 (i.e. all entries in A) are zero, then $\text{rank}(A) = 0$.

EXAMPLE 2.2

Find the rank of the matrix $A = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 2 & 4 & 2 \\ 0 & 2 & 2 & 1 \end{pmatrix}$

EXAMPLE 2.2

Find the rank of the matrix $A = \begin{pmatrix} 1 & 0 & 2 & 1 \\ 0 & 2 & 4 & 2 \\ 0 & 2 & 2 & 1 \end{pmatrix}$

The maximal minors have order 3, and we found that the one obtained by deleting the last column is $-4 \neq 0$. Hence $\text{rank}(A) = 3$.

EXAMPLE 2.3

Find the rank of the matrix $A = \begin{pmatrix} 1 & 2 & 1 & -1 \\ 9 & 5 & 2 & 2 \\ 7 & 1 & 0 & 4 \end{pmatrix}$

EXAMPLE 2.3

Find the rank of the matrix $A = \begin{pmatrix} 1 & 2 & 1 & -1 \\ 9 & 5 & 2 & 2 \\ 7 & 1 & 0 & 4 \end{pmatrix}$

The maximal minors have order 3, so we compute the 4 minors of order 3. The first one is

$$\begin{vmatrix} 1 & 2 & 1 \\ 9 & 5 & 2 \\ 7 & 1 & 0 \end{vmatrix} = 7 \times (-1) + (-1) \times (-7) = 0.$$

The other 3 are also zero. Since all minors of order 3 are zero, $\text{rank}(A) < 3$. We continue to look at the minors of order 2. The first one is

$$\begin{vmatrix} 1 & 2 \\ 9 & 5 \end{vmatrix} = 1 \times 5 - 2 \times 9 = -13 \neq 0.$$

The other 3 are also zero. Since all minors of order 3 are zero, $\text{rank}(A) < 3$. We continue to look at the minors of order 2. The first one is

$\begin{vmatrix} 1 & 2 \\ 9 & 5 \end{vmatrix} = 1 \times 5 - 2 \times 9 = -13 \neq 0$. It is not necessary to compute any more minors, and we conclude that $\text{rank}(A) = 2$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① A square matrix A is invertible if and only if $\det A \neq 0$. Since $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = \det I = 1 \Rightarrow \det A \neq 0$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① A square matrix A is invertible if and only if $\det A \neq 0$. Since
 $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = \det I = 1 \Rightarrow \det A \neq 0$.
- ② $\det(A^{-1}) = \frac{1}{\det A}$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① A square matrix A is invertible if and only if $\det A \neq 0$. Since
 $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = \det I = 1 \Rightarrow \det A \neq 0$.
- ② $\det(A^{-1}) = \frac{1}{\det A}$. Since
 $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = 1$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① A square matrix A is invertible if and only if $\det A \neq 0$. Since $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = \det I = 1 \Rightarrow \det A \neq 0$.
- ② $\det(A^{-1}) = \frac{1}{\det A}$. Since $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = 1$.
- ③ If A, B are invertible, then AB is also invertible and $(AB)^{-1} = B^{-1}A^{-1}$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① A square matrix A is invertible if and only if $\det A \neq 0$. Since
 $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = \det I = 1 \Rightarrow \det A \neq 0$.
- ② $\det(A^{-1}) = \frac{1}{\det A}$. Since
 $A.A^{-1} = I \Rightarrow \det A.\det(A^{-1}) = 1$.
- ③ If A, B are invertible, then AB is also invertible and $(AB)^{-1} = B^{-1}A^{-1}$. Since
 $(B^{-1}A^{-1}).(AB) = B^{-1}(A^{-1}.A)B = B^{-1}B = I$

SOME PROPERTIES OF INVERSE OF AN MATRIX

① If A is invertible and $\alpha \neq 0$, then $(\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

① If A is invertible and $\alpha \neq 0$, then $(\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$.

Since $\left(\frac{1}{\alpha} A^{-1}\right) \cdot (\alpha A) = I$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① If A is invertible and $\alpha \neq 0$, then $(\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$.

$$\text{Since } \left(\frac{1}{\alpha} A^{-1} \right) \cdot (\alpha A) = I.$$

- ② If A is invertible, then A^{-1}, A^T are also invertible and $(A^{-1})^{-1} = A$, $(A^T)^{-1} = (A^{-1})^T$.

SOME PROPERTIES OF INVERSE OF AN MATRIX

- ① If A is invertible and $\alpha \neq 0$, then $(\alpha A)^{-1} = \frac{1}{\alpha} A^{-1}$.

$$\text{Since } \left(\frac{1}{\alpha} A^{-1}\right) \cdot (\alpha A) = I.$$

- ② If A is invertible, then A^{-1}, A^T are also invertible and $(A^{-1})^{-1} = A$, $(A^T)^{-1} = (A^{-1})^T$. Indeed, $A^{-1} \cdot A = I$, $(A^{-1})^T \cdot A^T = (A \cdot A^{-1})^T = I^T = I$.

- 1 If A is an $n \times n$ square matrix and $\det(A) \neq 0$; B is an $n \times p$ matrix, then $AX = B$ has an unique solution $X = A^{-1}B$.

- ① If A is an $n \times n$ square matrix and $\det(A) \neq 0$; B is an $n \times p$ matrix, then $AX = B$ has an unique solution $X = A^{-1}B$.
- ② If A is an $n \times n$ square matrix and $\det(A) \neq 0$; B is an $p \times n$ matrix, then $XA = B$ has an unique solution $X = BA^{-1}$.

- ① If A is an $n \times n$ square matrix and $\det(A) \neq 0$; B is an $n \times p$ matrix, then $AX = B$ has an unique solution $X = A^{-1}B$.
- ② If A is an $n \times n$ square matrix and $\det(A) \neq 0$; B is an $p \times n$ matrix, then $XA = B$ has an unique solution $X = BA^{-1}$.
- ③ If A is an $n \times n$ square matrix and $\det(A) \neq 0$; B is an $m \times m$ square matrix and $\det(B) \neq 0$; C is an $n \times m$ matrix, then $AXB = C$ has an unique solution $X = A^{-1}CB^{-1}$.

EXAMPLE 3.1

Find matrix X which satisfies

$$\begin{pmatrix} 0 & -8 & 3 \\ 1 & -5 & 9 \\ 2 & 3 & 8 \end{pmatrix} X = \begin{pmatrix} -25 & 23 & -30 \\ -36 & -2 & -26 \\ -16 & -26 & 7 \end{pmatrix}$$

EXAMPLE 3.1

Find matrix X which satisfies

$$\begin{pmatrix} 0 & -8 & 3 \\ 1 & -5 & 9 \\ 2 & 3 & 8 \end{pmatrix} X = \begin{pmatrix} -25 & 23 & -30 \\ -36 & -2 & -26 \\ -16 & -26 & 7 \end{pmatrix}$$

$$\begin{aligned} X &= \begin{pmatrix} 0 & -8 & 3 \\ 1 & -5 & 9 \\ 2 & 3 & 8 \end{pmatrix}^{-1} \cdot \begin{pmatrix} -25 & 23 & -30 \\ -36 & -2 & -26 \\ -16 & -26 & 7 \end{pmatrix} = \\ &= \begin{pmatrix} 1 & 5 & 7 \\ 2 & -4 & 3 \\ -3 & -3 & -2 \end{pmatrix} \end{aligned}$$

EXAMPLE 3.1

Find matrix X which satisfies

$$\begin{pmatrix} 0 & -8 & 3 \\ 1 & -5 & 9 \\ 2 & 3 & 8 \end{pmatrix} X = \begin{pmatrix} -25 & 23 & -30 \\ -36 & -2 & -26 \\ -16 & -26 & 7 \end{pmatrix}$$

$$\begin{aligned} X &= \begin{pmatrix} 0 & -8 & 3 \\ 1 & -5 & 9 \\ 2 & 3 & 8 \end{pmatrix}^{-1} \cdot \begin{pmatrix} -25 & 23 & -30 \\ -36 & -2 & -26 \\ -16 & -26 & 7 \end{pmatrix} = \\ &= \begin{pmatrix} 1 & 5 & 7 \\ 2 & -4 & 3 \\ -3 & -3 & -2 \end{pmatrix} \end{aligned}$$

EXAMPLE 3.2

Find matrix X which satisfies $X \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix}$

EXAMPLE 3.2

Find matrix X which satisfies $X \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix}$

Solution.

$$X = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix} \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix}^{-1} =$$

EXAMPLE 3.2

Find matrix X which satisfies $X \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix}$

Solution.

$$X = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix} \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix}^{-1} = \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix}$$

EXAMPLE 3.2

Find matrix X which satisfies $X \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix} = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix}$

Solution.

$$X = \begin{pmatrix} -1 & 2 \\ -5 & 6 \end{pmatrix} \cdot \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix}^{-1} = \begin{pmatrix} 3 & -2 \\ 5 & -4 \end{pmatrix}$$

EXAMPLE 3.3

Find matrix X which satisfies

$$\begin{pmatrix} 3 & -1 \\ 5 & -2 \end{pmatrix} \cdot X \cdot \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix} = \begin{pmatrix} 14 & 16 \\ 9 & 10 \end{pmatrix}$$

EXAMPLE 3.3

Find matrix X which satisfies

$$\begin{pmatrix} 3 & -1 \\ 5 & -2 \end{pmatrix} \cdot X \cdot \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix} = \begin{pmatrix} 14 & 16 \\ 9 & 10 \end{pmatrix}$$

Solution.

$$\begin{aligned} X &= \begin{pmatrix} 3 & -1 \\ 5 & -2 \end{pmatrix}^{-1} \cdot \begin{pmatrix} 14 & 16 \\ 9 & 10 \end{pmatrix} \cdot \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \end{aligned}$$

EXAMPLE 3.3


Find matrix X which satisfies

$$\begin{pmatrix} 3 & -1 \\ 5 & -2 \end{pmatrix} \cdot X \cdot \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix} = \begin{pmatrix} 14 & 16 \\ 9 & 10 \end{pmatrix}$$

Solution.


$$\begin{aligned} X &= \begin{pmatrix} 3 & -1 \\ 5 & -2 \end{pmatrix}^{-1} \cdot \begin{pmatrix} 14 & 16 \\ 9 & 10 \end{pmatrix} \cdot \begin{pmatrix} 5 & 6 \\ 7 & 8 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \end{aligned}$$

EVALUATING THE AREA OF THE TRIANGLE



$$S = \frac{1}{2}abs|[\overrightarrow{AB}, \overrightarrow{AC}]| = \frac{1}{2}abs \begin{vmatrix} 2,5 & 1 & 1 \\ 3 & 2 & 1 \\ 1 & 3 & 1 \end{vmatrix} = \frac{5}{4}$$

EVALUATING THE VOLUME OF THE PARALLELEPIPED



$$\vec{a} = (a_1, a_2, a_3);$$

$$\vec{b} = (b_1, b_2, b_3); \vec{c} = (c_1, c_2, c_3)$$

$$\Rightarrow V = \text{abs}([\vec{a} \times \vec{b}], \vec{c}) = \text{abs} \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

MATLAB

- **Evaluating determinant of matrix A : $\det(A)$**
- **Finding inverse of Matrix A :**
 A^{-1} or $\text{inv}(A)$

THANK YOU FOR YOUR ATTENTION