Supplementary Material for: City charges for competing delivery carriers: exact best responses under probability-box travel times and leader pricing without equilibrium search

Quang-Vinh Dang

British University Vietnam, Hung Yen, Vietnam

Abstract

This document accompanies the main manuscript. It holds the full model with all assumptions, the complete statements of the algorithms and theorems, the extended data and calibration description with the parameter tables, the extended computational results with the per-city tables and the further experiments (the zone-market-clearing negative result, the details of the criterion-robustness study and of the BRL benchmark), all technical proofs, and the description of the Lean verification. Sections, theorems, tables, figures and equations of this document carry the prefix S; the main text refers to them by section number.

Contents and relation to the main text

The main text states every result with a proof sketch and reports the headline experiments. This document contains the following parts; the main text cites them as "Supplementary Material, Section Sn".

S1 Model and notation: the model of the main text with Assumptions A1–A3 (finite plans, affine coupling, carrier oracle).

S2 Algorithms: complete statements of BRL, branch-and-price details, the equilibrium engine, DIP and ARP, with Algorithms 1–3.

S3 Equilibrium theory: complete statements on imprecise beliefs, existence, credal absorption, complexity and the rounding algorithm.

S4 Data and calibration: the extended description of the instances, the parameters and their grades, and the simplifications.

S5 Computational study: the extended results, including the BRL benchmark details, the criterion comparison, the engine comparison with the ZMC negative result, the price of anarchy, imprecision, DIP, ARP and the criterion-robustness study.

S6 Additional tables and figures: the per-city and per-seed tables, the parameter table, the sensitivity analysis and the theory and ZMC checks.

S7 –[S10] Proofs for the equilibrium theory, BRL, DIP and ARP.

S11 Zone-market clearing (ZMC), including its guarantee and proofs.

S12 Machine-checked statements (Lean 4).

Numbers in the text are generated from the result files of the repository, as in the main text. The theorem numbers of this document are independent of those of the main text.

Email address: vinh.dq4@buv.edu.vn (Quang-Vinh Dang)

S1. Model and notation

This section repeats the model of the main text in full, with all assumptions, so that the statements and proofs below are self-contained.

Notation: $[n] = \{1, \dots, n\}$; for $v \in \mathbb{R}^d$ and $w \in \mathbb{R}^d_{>0}$, $\|v\|_{w,\infty} = \max_k |v_k|/w_k$ and $\|v\|_M^2 = \langle v, Mv\rangle$. Greek letters with related roles are kept apart: ρ_j is an overtime rate, $\bar\rho$ a belief radius and ϱ the contraction rate of ARP. A zone charge t_z carries no carrier index, whereas $t_i(p)$ is the traffic exposure of a plan (Assumption A1).

S1.1. Carriers, plans and congestion

The city is divided into zones $z \in [Z]$. There are N carriers. Carrier i has a finite plan set $\mathcal{P}_i$; a plan is a set of depot-to-depot routes over the carrier's customer zones. A plan p determines deliveries $q_{iz}(p)$, vehicles $n_{iz}(p)$ and vehicle-hours $\tau_{iz}(p)$ in each zone, and a routing cost $C_i(p)$ (Section S1.2). Write $Q_z = \sum_i q_{iz}$ and $T_z = \sum_i \tau_{iz}$. Inverse demand in zone z is $\alpha_z - b_z Q_z$ with $b_z > 0$, and a vehicle-hour costs $\gamma(\xi_z + c_z X_z)$, where $\gamma > 0$ is the hourly cost, ξ_z the free-flow travel-time factor with mean μ_z, $c_z > 0$ the congestion slope and X_z the congestion variable. The city charges s_z per delivery and t_z per unit of the congestion variable. Carrier i's profit is

$$u_i = \sum_z \big[q_{iz}(\alpha_z - b_z Q_z) - \gamma\tau_{iz}(\xi_z + c_z X_z) - s_z q_{iz} - t_z x_{iz}\big] - C_i(p). \tag{S1}$$

Two congestion regimes are distinguished.

- **D1 (exposure-based):** $x_{iz} = \tau_{iz}$ and $X_z = \sum_i \tau_{iz}$.
- **D2 (vehicle-count):** $x_{iz} = n_{iz}$ and $X_z = \sum_i n_{iz}$.

Under D1 the carriers' interaction depends on the same quantity that determines the carrier's own congestion cost. Under D2 the externality depends on a different quantity (counts) than the carrier's own exposure (hours), which breaks the potential structure (Section S3).

For the theory it is convenient to write the interaction abstractly. Let $d = 2Z$ with coordinates (z, Q) and (z, X).

Assumption A1 (Finite plans)**.** Plan $p \in \mathcal{P}_i$ has a contribution $a_i(p) \in \mathbb{R}^d_{\geq 0}$ to the aggregate, an exposure $\varphi_i(p) \in \mathbb{R}^d_{\geq 0}$ to the aggregate, a traffic exposure $t_i(p) \in \mathbb{R}^Z_{\geq 0}$ and a private profit $\pi_i(p)$. At a pure profile σ, $A(\sigma) = \sum_i a_i(\sigma_i)$ and $A_{-i} = A - a_i(\sigma_i)$.

Assumption A2 (Affine coupling)**.** There are a diagonal $M \geq 0$ and $\gamma > 0$ such that, for a rival aggregate $m \in \mathbb{R}^d_{\geq 0}$ and traffic factor $\xi \in \mathbb{R}^Z_{\geq 0}$, carrier i obtains $u_i(p; m, \xi) = \pi_i(p) - \langle \varphi_i(p), Mm\rangle - \gamma\langle t_i(p), \xi\rangle$.

Equation (S1) has this form with $a_{i,(z,Q)} = \varphi_{i,(z,Q)} = q_{iz}$, $M_{(z,Q)} = b_z$, $a_{i,(z,X)} = x_{iz}$, $\varphi_{i,(z,X)} = \tau_{iz}$, $M_{(z,X)} = \gamma c_z$ and $t_{iz} = \tau_{iz}$. The game is *of type D1* if $\varphi_i = a_i$ for all carriers and plans, and *of type D2* otherwise. Social welfare, with charges treated as transfers, is

$$W(\sigma) = \sum_z \big[\alpha_z Q_z - \tfrac{b_z}{2} Q_z^2 - \gamma\mu_z T_z - \gamma c_z T_z X_z\big] - \sum_i C_i(\sigma_i), \tag{S2}$$

that is, consumer surplus minus the congestion cost that carriers impose on each other, minus routing costs; in the experiments the traffic term is part of the routing cost and the social cost is $\mathrm{SC} = -W$. The affine coupling excludes power-law congestion such as BPR curves, which we linearise (Section S4).

S1.2. The routing layer

For every arc a the carrier knows a probability box: distribution functions $\underline{F}_a \le \overline{F}_a$ on a bounded support $[0, B_a]$, for example a Dvoretzky–Kiefer–Wolfowitz band around the empirical distribution of n observed days, $\underline{F}_a(x) = \max\{F_n(x) - \varepsilon, 0\}$ with $\varepsilon = \sqrt{\ln(2/\alpha)/(2n)}$ [1, 2]. The dependence between arcs is unknown, so the joint credal set is the Fréchet class $\mathcal{M}$ of all laws whose a-th marginal lies between $\underline{F}_a$ and $\overline{F}_a$. A route of duration $T_r = \sum_{a \in r} X_a$ incurs a convex piecewise-linear loss

$$\phi(T) = c_0 + \gamma T + \sum_{j=1}^{H} \rho_j \, (T - \Theta_j)^+ , \qquad \gamma \ge 0, \ \rho_j > 0, \ 0 \le \Theta_1 < \cdots < \Theta_H. \tag{S3}$$

In the experiments $H = 1$: a fixed cost, an hourly cost and an overtime penalty after the shift length. The robust route cost is the upper prevision $\overline{\mathbb{E}}[\phi(T_r)] = \sup_{P \in \mathcal{M}} \mathbb{E}_P \phi(T_r)$, and $C_i(p) = \sum_{r \in p} \overline{\mathbb{E}}[\phi(T_r)]$. A carrier's best response to prices on its deliveries and hours is therefore a prize-collecting routing problem in which every route is priced by $\overline{\mathbb{E}}[\phi]$. We call this problem the *carrier oracle*:

Assumption A3 (Oracle)**.** For each i there is an oracle that, for price vectors $\lambda^a, \lambda^\varphi \in \mathbb{R}^d$, $\nu \in \mathbb{R}^Z$ and a diagonal $Q \succeq 0$, returns a plan in $\arg\max_{p \in \mathcal{P}_i} \{\pi_i(p) + \frac{1}{2}\langle a_i(p), Q a_i(p)\rangle - \langle \lambda^a, a_i(p)\rangle - \langle \lambda^\varphi, \varphi_i(p)\rangle - \langle \nu, t_i(p)\rangle\}$.

A best response at the belief (m, ξ) is the call $(0, Mm, \gamma\xi, 0)$. Section S2.1 gives an exact implementation of the oracle. We count *oracle calls* as the cost unit of every method below.

S1.3. The leader's problem

The city chooses charges u from an admissible set $\mathcal{U}$. For a charge u let $\mathrm{NE}(u)$ be the set of pure Nash equilibria of the carriers' game. With optimistic selection the leader's problem is

$$\min_{u \in \mathcal{U}, \ \sigma} \ \mathrm{SC}(\sigma) \qquad \text{s.t.} \qquad \sigma \in \mathrm{NE}(u). \tag{S4}$$

In the D1 instance used for computation (Section S2.4), each carrier serves its customer zones in all-or-nothing lots, and the charges reach the game through one number per zone. The three methods compared in Section S5 differ in what the leader may use: the *grid* and DIP use the carriers' routing oracle, ARP uses only the zone aggregates the market produces after each posted charge.

S2. Algorithms, full statements

The algorithms form a stack. BRL (Section S2.1) and branch-and-price (Section S2.2) implement the carrier oracle exactly. An equilibrium engine (Section S2.3) calls the oracle to compute a pure equilibrium for a given charge. DIP (Section S2.4) and ARP (Section S2.5) choose the charges; the grid baseline calls the engine once per grid point.

S2.1. Budget-resource labelling (BRL)

In words. A route is hard to price when the delays on its arcs could all be bad at the same time. Under unknown dependence the worst case is exactly that: every arc is equally unlucky at once. The route's cost then splits arc by arc once a time budget is allocated to each arc. BRL treats this budget as a resource in a shortest-path search. A partial route keeps its cost as a function of the budget, and it is dropped only if another partial route is at least as cheap for every budget that a completion can still use. That condition is why the pruning cannot discard the best route (Theorem S4).

Let $\overline{Q}_a(u) = \inf\{x : \underline{F}_a(x) \ge u\}$ be the upper quantile of arc a, $\bar{x}_a = \int_0^1 \overline{Q}_a(u)\, du$ and $\overline{\Pi}_a(d) = \int_0^1 \left(\overline{Q}_a(u) - d\right)^+ du$, the stop-loss transform of the lower distribution function. $\overline{\Pi}_a$ is convex and nonincreasing with slopes in $[-1, 0]$.

Theorem S1 (Upper prevision and budget decomposition)**.** *For every route r,*

$$\overline{\mathbb{E}}[\phi(T_r)] = \int_0^1 \phi\Big(\sum_{a\in r} \overline{Q}_a(u)\Big)du = c_0 + \sum_{a\in r} \gamma \bar{x}_a + \sum_{j=1}^{H} \rho_j \min\Big\{\sum_{a\in r} \overline{\Pi}_a(d_a^j) : \sum_{a\in r} d_a^j \le \Theta_j\Big\}.$$

The supremum is attained by the comonotone coupling of the lower CDFs; each minimum is attained, at an integer vector when quantiles and thresholds are integers.

The first equality and the stop-loss identity are due to Meilijson and Nádas [3] (see also 4, 5), and the lower CDF as worst marginal is standard [6]; we re-prove the statement in Section S8 because the algorithm relies on the inequality-budget form and on integrality (see "Relation to prior work" below). With a common quantile grid of L levels (weights ω_l), the route profile $S_{r,l} = \sum_{a\in r} q_{a,l}$ gives $\overline{\mathbb{E}}[\phi(T_r)] = c_0 + f_r(T_{\max})$ for $H = 1$, where $f_r(D) = \gamma \sum_l \omega_l S_{r,l} + \rho \sum_l \omega_l \left(S_{r,l} - D\right)^+$.

Theorem S2 (Budget-resource reformulation)**.** *Let y be dual prices and define the arc costs*

$$c_a(d^1, \dots, d^H) = \gamma \bar{x}_a + \sum_j \rho_j \overline{\Pi}_a(d^j).$$

The dependence-robust pricing problem

$$\min_{r\in\mathcal{R}} \Big\{\overline{\mathbb{E}}[\phi(T_r)] - \sum_{i\in r} y_i\Big\}$$

equals an elementary shortest path problem in which every arc chooses a consumption d_a^j of each of H budget resources, paid through the separable convex cost c_a and subject to the budget constraints $\sum_{a\in r} d_a^j \le \Theta_j$, $j \le H$. The number of resources is H; it does not depend on L, the sample size or the number of arcs.

A scenario view of the same problem, which is what carrying the comonotone reduction of Ghosal et al. [7] into a pricing problem would give, needs the L level sums $(S_{p,l})_l$ as resources and compares labels pointwise. BRL instead compares labels through their cost functions, but only where a completion can still reach. For a partial path (label) p ending at v with level profile S_p, collected duals $y(p)$, visited set $\mathcal{V}(p)$ and load $\ell(p)$, concatenation with a completion q is an inf-convolution, $f_{p\oplus q}(T_{\max}) = \min_D\{f_p(D) + f_q(T_{\max} - D)\}$ (Lemma S46). Let $C_{\min}(v)$ be a lower bound on the duration of any completion from v and define the completion window $\mathcal{W}(p) = [\underline{D}(p), \overline{D}(p)]$ with $\overline{D}(p) = \min\{S_{p,L}, T_{\max} - C_{\min}(v)\}$ and $\underline{D}(p) = \min\{S_{p,1}, \overline{D}(p)\}$.

Definition S3 (Functional dominance)**.** Label p_1 functionally dominates p_2, both ending at v, if $\mathcal{V}(p_1) \subseteq \mathcal{V}(p_2)$, $\ell(p_1) \le \ell(p_2)$ and $(f_{p_1}(D) - y(p_1)) - (f_{p_2}(D) - y(p_2)) \le 0$ for all $D \in \mathcal{W}(p_2)$.

The difference is piecewise linear with breakpoints at the level sums, so the test needs at most $2L + 2$ evaluations.

Theorem S4 (Validity and exactness)**.** *Functional dominance implies that every completion of p_2 is a completion of p_1 with no larger reduced cost. The labelling algorithm that discards dominated labels (Algorithm 1) returns the minimum reduced cost exactly, with a route attaining it.*

The window is what makes the rule exact and strong at the same time. The slope of f_p lies in $[-\rho, 0]$ and that of $D \mapsto f_q(T_{\max} - D)$ in $[0, \rho]$, so an optimal split never lies above $\overline{D}(p)$, and below $S_{p,1}$ the difference of two label functions is nondecreasing (Lemma S47). Quantile-vector dominance and sector dominance are special cases of functional dominance, and neither converse holds (Proposition S48). Under independent arc times concatenation is a convolution of laws

Algorithm 1 BRL pricing for the dependence-robust routing problem ($H = 1$)

Require: graph, demands, capacity; level weights ω; arc level profiles $q_{a,\cdot}$; $\gamma, \rho, T_{\max}, c_0$; duals y

$C_{\min}(v) \leftarrow$ shortest path from v to the depot with arc lengths $q_{a,1}$
buckets $\mathcal{B}_v \leftarrow \emptyset$; queue $\leftarrow$ {depot label}
while queue nonempty **do**
    pop p; record the closed route $p \oplus (v, 0)$ with reduced cost $c_0 + f_{p\oplus(v,0)}(T_{\max}) - y(p)$
    for each customer $j \notin \mathcal{V}(p)$ that fits the capacity **do**
        $p' \leftarrow$ extension of p to j ($S_{p'} = S_p + q_{(v,j),\cdot}$, $y(p') = y(p) + y_j$)
        if some $o \in \mathcal{B}_j$ functionally dominates p' **then** discard p'
        else delete the labels of $\mathcal{B}_j$ dominated by p'; insert and push p'
return recorded routes sorted by reduced cost

rather than an inf-convolution and the method does not apply. The natural shortcut of fixing a single risk level for all routes gives a lower bound but is not exact (Proposition S49).

On the city instances the pricing runs over ng-routes (Section S2.2). Column generation [8, 9] with BRL pricing returns the exact LP bound of the dependence-robust routing problem when no negative reduced cost remains. An integer program over the generated columns (price-and-branch) gives a feasible plan but is not an exact integer method; Section S2.2 makes the integer step exact.

Relation to prior work. The upper-prevision formula and the stop-loss decomposition are due to Meilijson–Nádas and Tchen; Theorem S1 is re-proved in Section S8 because the algorithm uses its inequality-budget form and its integrality. The comonotone worst case for a sum along a route is used in routing by Ghosal et al. [7] (their Theorem 4, from Long–Qi–Zhang) for ambiguous demand in a capacity constraint with known travel times, solved by branch-and-cut on a vehicle-flow formulation; the supermodularity argument applies verbatim to arc times, and we claim no novelty for the reduction. Their paper has no pricing problem, budget resource or label dominance. BRL contributes the per-arc budget as a decision resource of the pricing problem (Theorem S2) and the completion-window functional dominance with its exactness proof (Theorem S4). Resource extension functions [10] are the general language for such pricing problems. With the L level sums as resources, the classical dominance for non-decreasing functions is our quantile-vector baseline, and for integer values a chosen arc consumption is a bundle of parallel arcs with classical functions. In that framework labels are resource vectors joined by composition; BRL labels are functions of the budget joined by inf-convolution, and the window rests on the bounded slope of the loss (Lemma S47), which the theory of resource extension functions does not supply. Cost-function labels compared over a range of resource values are used for time-window problems with linear node costs [11], so we claim the window and its exactness proof for this loss, not the idea of comparing cost functions.

S2.2. Branch-and-price with BRL pricing

In words. Price-and-branch solves the integer problem only over the routes found at the root, so it can miss the integer optimum. Branch-and-price keeps generating routes inside a branching tree, so the plan it returns is provably optimal, and its lower bound gives every plan an optimality gap.

We embed the BRL pricer in branch-and-price. The master is a set-partitioning model for the routing benchmark, or a prize-collecting set-packing model for the carrier oracle, with one column pool shared by all nodes. Each node solves its LP by column generation with the exact BRL pricer, restricted to the node's allowed arcs with $C_{\min}$ recomputed on them; functional dominance remains valid because the restrictions are local.

- *Branching* is on the aggregated arc flow f_{ij}: $f_{ij} = 0$ deletes the arc; $f_{ij} = 1$ deletes the other arcs leaving i and entering j. Integral arc flows imply an integral plan.

- *Fleet size.* When every customer must be served, a row for the minimum number of routes is added at the root and a fractional route count is branched on first.
- *ng-routes* [12]. The visited set is replaced by the ng-memory, and Theorem S4 holds verbatim with the weaker relaxation. If a node solution contains a non-elementary column, the ng-sets along its cycle are enlarged; reported plans are elementary.
- *Bounds.* The node LP value and the Lagrangian bound during column generation are valid lower bounds, marked invalid if a pricing call hits the label limit. Upper bounds come from integral nodes and from an integer program over the generated elementary columns. Nodes are chosen by best bound; the method stops at a relative gap of 10^{-6}.

Since ng-routes relax elementary routes, a zero gap proves optimality for the elementary problem. This is standard branch-and-price machinery; its purpose here is to remove the inexact integer step, not to add a branching rule.

S2.3. The equilibrium engine

In words. For a given charge, the engine starts from an approximate market-clearing guess and then lets the carriers best-respond one at a time. Under exposure-based congestion every improving move raises a single shared potential, so the sequence cannot cycle and ends at a pure equilibrium (Proposition S26).

Phase one is zone-market clearing (ZMC): every carrier's oracle is queried at a common zone-price vector for eight rounds, using the sign of the clearing residual to update the prices (Section S11). A restricted master over the collected plans and derandomised rounding give a start profile. In phase two, sequential best responses (one oracle call each) run until no carrier improves; since D1 has an exact potential (Proposition S26) this terminates at a pure Nash equilibrium. Each carrier's column pool is kept across calls, because route costs do not depend on prices. In the city experiments a carrier chooses node-disjoint routes from its depot over 22 customer zones; each served zone adds its whole lot (all-or-nothing) together with prize, own Cournot and congestion terms, charges and prices as node costs, and each route costs its BRL upper expectation.

Remark S5 (ZMC and its negative result). ZMC has a clearing identity that bounds the potential gap and each carrier's regret by the clearing error, and a price-bracketing rate that does not depend on d (Section S11). On synthetic games with non-separable plan sets, it gave the smallest potential gap in 17 of 20 instances at an equal call budget (Table S20). On the nine city games it did not beat plain best response: it used 36–65 calls, against 8–17 for best response, and was strictly better than all other methods in 0 cities (Section S5.3). We use it as the starting phase of the engine and make no claim for it.

S2.4. Dual implementation pricing (DIP)

In words. A city that could see everything would first pick the outcome it wants, the social optimum, and then look for charges under which the carriers choose it voluntarily. DIP does exactly this. It finds the charges by repeatedly asking each carrier what it would do instead (one best response) and adding a constraint that rules that deviation out. Because each question is one routing solve at a fixed target, no equilibrium has to be computed for any candidate charge.

A grid of charges needs one equilibrium solve per grid point (Section S5.5.1). DIP replaces the search by an implementation step: it computes the profile the city wants and then computes charges under which that profile is an equilibrium, using the carriers' own oracle to separate candidate charges. No equilibrium is solved for the leader's decision.

Setting. We use the D1 instance of Section S1 in cost form. Carrier i serves a subset of its customer zones with all-or-nothing lots $d_{iz} > 0$, so $q_{iz} \in \{0, d_{iz}\}$ and $q_{iz}^2 = d_{iz} q_{iz}$; vehicle-hours are $\tau_{iz} = r\, q_{iz}$ for one ratio $r > 0$ (assumption (C3)); $R_i(p) \geq 0$ is the robust routing cost of plan p; $g_z = \gamma c_z$. The city posts s_z per delivery and t_z per vehicle-hour. Carrier i's cost and the social cost (charges are transfers) are

$$J_i(p; \sigma_{-i}) = R_i(p) + \sum_z \big[q_{iz}(-\alpha_z + b_z(q_{iz} + Q_{-i,z}) + s_z) + \tau_{iz}(g_z(\tau_{iz} + X_{-i,z}) + t_z) \big],$$

$$\mathrm{SC}(\sigma) = \sum_i R_i(\sigma_i) + \sum_z \big[-\alpha_z Q_z + \tfrac{b_z}{2} Q_z^2 + g_z X_z^2 \big].$$

Write $\kappa_z = g_z r^2$ (congestion slope in delivery units), $m_z = b_z + \kappa_z$ (the slope the carriers face) and $\bar{d}_z$ for the mean lot in zone z.

Proposition S6 (One instrument per zone)**.** *(a) For every plan, $\sum_z (s_z q_{iz} + t_z \tau_{iz}) = \langle u, q_i \rangle$ with $u_z = s_z + r t_z$, so the game depends on the charges only through $u \in \mathbb{R}^Z$. (b) Every u is realised by a uniform per-delivery charge $s = \min_z u_z$ over zones with lots and non-negative zone congestion charges $t_z = (u_z - s)/r$. (c) The grid of Section S5.5 is the two-parameter family $u = s\mathbf{1} + \vartheta\, r\, g \odot X^{\mathrm{NE}}$.*

Under u the game has the exact potential $\Phi_u(\sigma) = \sum_i R_i(\sigma_i) + \sum_z [-\alpha_z Q_z + \frac{m_z}{2}(Q_z^2 + \sum_i q_{iz}^2) + u_z Q_z]$ (Proposition S26).

Lemma S7 (Potential–welfare identity)**.** *Fix $Q^\circ \in \mathbb{R}^Z$, let $u_z^{\tan} = \kappa_z Q_z^\circ - \frac{m_z}{2} \bar{d}_z$ and $v = u - u^{\tan}$. For every pure profile σ,*

$$\Phi_u(\sigma) = \mathrm{SC}(\sigma) - \sum_z \tfrac{\kappa_z}{2}(Q_z - Q_z^\circ)^2 + L(\sigma) + \langle v, Q \rangle + \sum_z \tfrac{\kappa_z}{2}(Q_z^\circ)^2, \qquad L(\sigma) = \sum_{i,z} \tfrac{m_z}{2}(d_{iz} - \bar{d}_z) q_{iz}.$$

The *tangent charge* $u^{\tan}$ is the classical corrective tax of a Cournot market with an externality: the marginal external damage at the target aggregate minus a market-power subsidy [13, 14]. Two details are specific to all-or-nothing lots: the subsidy is half a lot, $\frac{m_z}{2}\bar{d}_z$ per delivery, and it includes the carriers' own congestion. With carrier-specific charges $U_{iz} = \kappa_z Q_{-i,z}^* - \frac{b_z}{2} d_{iz}$ every carrier's cost differences equal the social-cost differences, so every social optimum is an equilibrium; this is the classical marginal-contribution charge [15, 16] (Proposition S50 in Section S9). A city cannot charge per carrier, and the next results measure what flat zone charges lose. In the convexified (mixed) game let x^* minimise the convexified social cost $\overline{\mathrm{SC}}$, with aggregate $\bar{Q}^*$.

Theorem S8 (Loss under asymmetric lots)**.** *Let $u = \kappa \odot \bar{Q}^* - \frac{m}{2} \odot \bar{d}$. Every convexified equilibrium x^u satisfies $0 \leq \overline{\mathrm{SC}}(x^u) - \overline{\mathrm{SC}}(x^*) \leq (1 + \varrho)E$, where $\varrho = \max_z \kappa_z/(b_z + \kappa_z) < 1$ and $E = \sum_{i,z} \frac{m_z}{2} |d_{iz} - \bar{d}_z|\, d_{iz}$ measures how unequal the lots in a zone are.*

Theorem S9 (Exact implementation with symmetric lots)**.** *If all lots in a zone are equal, the tangent charge makes x^* a convexified equilibrium, and every convexified equilibrium under it has aggregate $\bar{Q}^*$ and social cost $\overline{\mathrm{SC}}(x^*)$.*

Sketch. The identity holds for the convexified potential and social cost with the same algebra. Because $\overline{\mathrm{SC}}$ is linear plus a quadratic in $\bar{Q}$, its expansion around x^* is exact; comparing it with the identity at a minimiser of the potential bounds the aggregate deviation by the lot-asymmetry term L, and $|L| \leq E$. With equal lots $L \equiv 0$.

The pure game. For a pure target σ^* and an admissible box $\mathcal{U}$ of charges let

$$P(\sigma^*) = \big\{ u \in \mathcal{U} :\ \langle q_i(p) - q_i(\sigma_i^*), u \rangle \geq J_i^0(\sigma_i^*; \sigma_{-i}^*) - J_i^0(p; \sigma_{-i}^*)\ \ \forall i,\ \forall p \in \mathcal{P}_i \big\},$$

Algorithm 2 DIP: dual implementation pricing

Target. Column generation on the convexified social cost (the carriers' oracle at social prices), rounding and coordinate polish $\Rightarrow$ pure profile σ^* with aggregate Q^*
Reference. $u^{\mathrm{ref}} \leftarrow \kappa \odot Q^* - \frac{m}{2} \odot \bar{d}$ (tangent charge); cuts $\mathcal{C} \leftarrow \emptyset$; $u \leftarrow u^{\mathrm{ref}}$
repeat
  for each i: $b_i \leftarrow$ best response to σ^*_{-i} under u ▷ N routing calls, no equilibrium solve
  for each i with $J_i^u(b_i) < J_i^u(\sigma_i^*)$: add $\langle q_i(b_i) - q_i(\sigma_i^*), u\rangle \geq J_i^0(\sigma_i^*) - J_i^0(b_i) + \varepsilon$ to $\mathcal{C}$
  if no cut was added: **return** u ▷ σ^* is an exact pure equilibrium under u
  $u \leftarrow \arg\min\{\|u - u^{\mathrm{ref}}\|_1 : u \in \mathcal{U} \text{ satisfies } \mathcal{C}\}$ (LP); **if** infeasible: **break**
until
Fallback (only if $P(\sigma^*) = \emptyset$): best-response dynamics from σ^* under the least-violating charge and under u^{ref}; keep the better equilibrium
Instruments. $s = \min_z u_z$, $t_z = (u_z - s)/r$ (Proposition S6)

with J_i^0 the cost at zero charges. Since $J_i^u = J_i^0 + \langle u, q_i\rangle$, $P(\sigma^*)$ is exactly the set of charges under which σ^* is a pure Nash equilibrium: a polyhedron, possibly empty. Toll polyhedra of this kind are classical for nonatomic routing [17, 18] and inverse optimisation [19].

Theorem S10 (Separation and finite termination)**.** *(a) $u \in P(\sigma^*)$ if and only if no carrier has a strictly improving best response to σ^*_{-i} under u; an improving best response gives that carrier's most violated constraint. (b) With $\varepsilon = 0$, Algorithm 2 stops after at most $1 + \sum_i(|\mathcal{P}_i| - 1)$ rounds, either with $u \in P(\sigma^*)$ or with an infeasible LP, which happens only if $P(\sigma^*) = \emptyset$. Each round costs N oracle calls and no equilibrium solve.*

Proposition S11 (Single-level form)**.** *With optimistic selection, the leader's value is $V^\star = \min\{\mathrm{SC}(\sigma) : \sigma \text{ pure},\ P(\sigma) \neq \emptyset\} \geq \min_\sigma \mathrm{SC}(\sigma)$. If Algorithm 2 ends in its main loop, the leader's loss is at most the suboptimality of the target, $\mathrm{SC}(\sigma^*) - \min_\sigma \mathrm{SC}(\sigma)$; in particular DIP is at least as good as any grid over any subfamily of charges when its target is a social optimum.*

Theorem S12 (Loss of the fallback)**.** *For any charge u with $v = u - u^{\tan}$ and $Q^\circ = Q^*$, the equilibrium σ^u reached from σ^* by improving moves satisfies $\mathrm{SC}(\sigma^u) - \mathrm{SC}(\sigma^*) \leq \sum_z \frac{\kappa_z}{2}(Q_z^u - Q_z^*)^2 + L(\sigma^*) - L(\sigma^u) + \langle v, Q^* - Q^u\rangle$.*

Corollary S13 (Oracle calls)**.** *DIP uses the calls of the social target, N calls per cutting-plane round, and, only if $P(\sigma^*)$ is empty, the calls of the fallback dynamics. A grid of G points with an equilibrium solve of C_{eq} calls per point uses GC_{eq}.*

Theorem S14 (Imprecision shifts the charge)**.** *Under Γ-maximin credal boxes of width δ (Lemma S25), the charges that implement σ^* are $P(\sigma^*) - \hat{s}(\delta)$ with $\hat{s}_z(\delta) = \delta(b_z w_{(z,Q)} + r\, g_z w_{(z,X)})$. One run of DIP therefore gives $u(\delta) = u(0) - \hat{s}(\delta)$ for every width at no further oracle cost, and the leader attains the same social cost at every width as long as $u(\delta)$ is admissible. For E-admissibility, maximality and interval dominance, $u(0)$ works at every width.*

Proposition S15 (When one instrument suffices)**.** *A uniform per-delivery charge implements the convexified optimum under equal lots if the tangent charge is the same in every active zone. With the calibration of Section S4 ($\alpha_z = 1.5$ fee, $b_z = \alpha_z/(2Q_z^{\mathrm{full}})$) and n_z carriers with equal lots in zone z, the congestion-free tangent charge is $u_z = -0.375\,\mathrm{fee}/n_z$: a subsidy of 37.5% of the fee in single-carrier zones and 19% in two-carrier zones. The congestion charge contributes only spatial differentiation.*

Proofs are in Section S9. Theorem S10(a) is what makes the method cheap: the separation oracle is one best response per carrier at the fixed target, not an equilibrium solve. Selection remains optimistic: the target is an exact equilibrium under the returned charges, but other equilibria may coexist.

Algorithm 3 ARP flat rule

Require: slopes b, g, r; damping θ

$u \leftarrow 0$; observe (Q, n) of the unpriced equilibrium

repeat

  update the lot estimates $\bar{d}_z$ from (Q, n) and the previous observations

  $u_z \leftarrow (1-\theta)u_z + \theta[\kappa_z Q_z - \frac{m_z}{2}\bar{d}_z]$ for all z

  post u; the carriers best-respond from their current plans until nobody moves; observe (Q, n)

until $u = \kappa Q - \frac{m}{2}\bar{d}$ (fixed point) or the posted charge and the market state repeat (cycle)

Relation to prior work. DIP combines the Pigou–Cournot corrective tax [13, 14], marginal-contribution utilities [15, 16], toll polyhedra for implementing a target [17, 18, 20, 21], and an inverse-optimisation cutting plane [19, 22, 23] with best-response cuts as used for integer programming games [24]; implementation of a target by incentives and taxes in atomic congestion games is studied by Monderer and Tennenholtz [25] and Caragiannis et al. [26]. The separation step of Theorem S10 is one best response per carrier at the fixed target, so no equilibrium is solved, whereas the closest leader-side method [27] solves $2B$ approximate equilibria per outer step. New here are the exact potential–welfare identity of the lumpy Cournot–congestion game with loss bounds in the lot-asymmetry number E, the translation of the implementing charges by the Γ-maximin surcharge, and the competitor-count rule.

S2.5. Aggregate-response pricing (ARP)

In words. ARP asks what a city can do without the carriers' routing models. It posts a charge, watches the zone totals that the carriers then deliver, and moves each zone's charge towards the corrective tax that those totals imply. In the convexified game this update is a contraction, so the iteration settles at a rate that depends only on the demand and congestion slopes, not on the number of carriers or on how hard their routing problems are.

DIP and the grid need the carriers' routing oracle, which a real city does not have. ARP asks how much of the leader's gain is reachable from what the city can observe. The city knows its own demand and congestion slopes (b, g, r) and hence $\kappa_z = g_z r^2$ and $m_z = b_z + \kappa_z$. It posts a per-delivery-equivalent charge $u \in \mathbb{R}^Z$ (realised, as in Proposition S6, by a uniform per-delivery charge plus non-negative zone congestion charges), the carriers settle in a pure equilibrium by their own best responses, and the city observes the deliveries Q_z and the number n_z of active carriers in each zone, nothing else.

The flat rule. With an estimate $\bar{d}_z$ of the mean lot, the city updates

$$u_z^{t+1} = (1-\theta_z)u_z^t + \theta_z\big[\kappa_z Q_z^t - \tfrac{m_z}{2}\bar{d}_z\big], \qquad \theta_z \in (0, 1]. \tag{S5}$$

With $\theta = 1$ this is DIP's tangent charge evaluated at the observed aggregate instead of a computed target. The lot estimate is Q_z/n_z; when n_z changes by one between observations, the jump of Q_z is the lot of the carrier that entered or left (Lemma S53); for a zone never served nothing is learned and the city-wide mean observed lot is used.

The marginal tariff (ARP-M). The flat rule charges each delivery the externality at the observed aggregate. The exact externality of carrier i's lot on the others is $\kappa_z d_{iz} Q_{-i,z}$, and its market-power wedge is $\frac{b_z}{2} d_{iz}^2$; both are functions of the carrier's own deliveries and of the zone total, so an anonymous nonlinear tariff can bill them exactly, in the limit of re-tangenting at every transaction.

Theorem S16 (Marginal tariff; the mechanism is classical)**.** *Let every carrier pay, in every zone,* $T_z(q_{iz}, Q_z) = \kappa_z q_{iz}(Q_z - q_{iz}) - \frac{b_z}{2} q_{iz}^2$. *Then (a)* $J_i^T(p; \sigma_{-i}) = \mathrm{SC}(p, \sigma_{-i}) - \mathrm{SC}(\emptyset_i, \sigma_{-i})$ *for all* i, p

and σ_{-i}, so SC *is an exact potential of the tariff game; (b) the pure equilibria are exactly the profiles that no single carrier can improve socially, every social optimum is a pure equilibrium, and best-response dynamics from any profile never increase* SC *and stop at such an equilibrium; (c) given the rivals, the tariff is linear in the carrier's own service indicators, so a best response is one oracle call with the zone prices shifted by $\kappa_z Q_{-i,z} - \frac{b_z}{2} d_{iz}$.*

The mechanism is the marginal-contribution tariff of Wolpert and Tumer [15], Sandholm [28] and Marden and Wierman [16]; what is stated here is its lumpy form in the city game and the consequence that DIP's implementation step is unnecessary under it. The tariff needs billing on own deliveries and the zone total. It leaves equilibrium selection open: the carriers' dynamics stop at a unilateral social optimum, which may be a local optimum of SC. In the experiments we run it with the carriers' own best responses from the status quo (tariff+BR) and with the potential-minimising solver of the engine on the social-cost potential (tariff+SO).

Theory of the flat rule. Convexify the plan sets: let $\mathcal{X} = \prod_i \Delta(\mathcal{P}_i)$, $\bar{q}_i(x_i) = \mathbb{E}_{x_i} q_i$, $\bar{Q}(x) = \sum_i \bar{q}_i$ and $\mathcal{Q} = \bar{Q}(\mathcal{X})$. With a reference lot vector $\bar{d}$ define the carriers' aggregate value function $G(Q) = \min\{\sum_i \mathbb{E}_{x_i}[R_i + \sum_z \frac{m_z}{2}(d_{iz} - \bar{d}_z) q_{iz}] : \bar{Q}(x) = Q\}$ and $F(Q) = G(Q) + \sum_z [-\alpha_z Q_z + \frac{m_z}{2} Q_z^2 + \frac{m_z}{2} \bar{d}_z Q_z] + \iota_{\mathcal{Q}}(Q)$, where R_i is the routing cost. Let $L_{\bar{d}}(\sigma) = \sum_{i,z} \frac{m_z}{2}(d_{iz} - \bar{d}_z) q_{iz}$ be the lot-asymmetry term of Section S2.4 with reference lots $\bar{d}$, $E_{\bar{d}} = \sum_{i,z} \frac{m_z}{2} |d_{iz} - \bar{d}_z| d_{iz}$, and $g(Q) = \sum_z [\frac{\kappa_z}{2} Q_z^2 - \frac{m_z}{2} \bar{d}_z Q_z]$. All statements below are proved in Section S10; $\|v\|_w^2 = \sum_z w_z v_z^2$.

Theorem S17 (The response is a proximal map)**.** *F is 1-strongly convex in $\|\cdot\|_m$. For every charge u, all convexified equilibria share one aggregate $Q(u) = \arg\min_Q \{F(Q) + \langle u, Q\rangle\} = \nabla F^*(-u)$, and $\|Q(u_1) - Q(u_2)\|_m \le \|u_1 - u_2\|_{m^{-1}}$. The leader's problem $\min \overline{\mathrm{SC}} + L_{\bar{d}}$ is the Fenchel dual $\min_u D(u)$, $D(u) = F^*(-u) + \sum_z (u_z + \frac{m_z}{2} \bar{d}_z)^2 / (2\kappa_z)$, whose gradient $\kappa^{-1} \odot (u + \frac{m}{2} \bar{d}) - Q(u)$ is observable from the aggregate alone. Its unique minimiser u^* is the unique fixed point of $u \mapsto \kappa \odot Q(u) - \frac{m}{2} \odot \bar{d}$. With symmetric lots within zones the fixed point implements a convexified social optimum; in general it is within $E_{\bar{d}}$ of one.*

Theorem S18 (Instance-free contraction)**.** *Let $\varrho = \max_z \kappa_z / m_z < 1$ over zones with a lot. In the convexified game, (a) the plain rule ($\theta = 1$) satisfies $\|u^{t+1} - u^*\|_{\kappa^{-1}} \le \varrho \|u^t - u^*\|_{\kappa^{-1}}$; (b) the damped rule with $\theta = 2/(2+\varrho)$ contracts at ratio $\varrho/(2+\varrho) < 1/3$. The ratios depend on the instance only through ϱ, not on Z, N, the number of customers, the lots, the routing costs or the probability boxes.*

The rate comes from the Cournot slope b_z: without the demand side ($b = 0$), $\varrho = 1$ and there is no contraction, which is why the nonatomic literature uses averaging [29]. The flat rule is gradient descent on D in the κ^{-1} geometry; this identification, and the resulting rate in a lumpy NP-hard routing game, is the new part.

Theorem S19 (Pure game and identification)**.** *(a) If σ is a pure equilibrium under $u = \kappa \odot Q(\sigma) - \frac{m}{2} \odot \bar{d} + v$, then for every carrier i and plan p, with $\sigma' = (p, \sigma_{-i})$ and $\Delta Q = Q(\sigma') - Q(\sigma)$, $\mathrm{SC}(\sigma') \ge \mathrm{SC}(\sigma) + [L_{\bar{d}}(\sigma) - L_{\bar{d}}(\sigma')] + \sum_z \frac{\kappa_z}{2} \Delta Q_z^2 - \langle v, \Delta Q \rangle$: at an exact fixed point with symmetric lots, no carrier can lower the social cost unilaterally. (b) If the observed aggregate Q^t differs from the convexified response by e^t, then $\|u^{t+1} - u^*\|_{\kappa^{-1}} \le \varrho \|u^t - u^*\|_{\kappa^{-1}} + \|e^t\|_\kappa$, hence $\limsup \|u^t - u^*\|_{\kappa^{-1}} \le \sup \|e^t\|_\kappa / (1 - \varrho)$; if the carriers settle in potential minimisers, $\|e^t\|_\kappa \le \sqrt{2\varrho\eta_t}$ with η_t the rounding gap of the potential. (c) The convexified response map determines F up to a constant, so the convexified optimum is all that an observation-only scheme can target; the pure social optimum is not identified (Example S52).*

Proposition S20 (Imprecise carriers)**.** *Under Γ-maximin beliefs of width δ the response to u is the precise response to $u + \hat{s}(\delta)$ (Lemma S25), so the flat rule needs no knowledge of δ and contracts at the same rates; it only moves the fixed point, and the loss in $\overline{\mathrm{SC}} + L_{\bar{d}}$ against the precise fixed point is at most $\frac{1}{2} \|\hat{s}(\delta)\|_{m^{-1}}^2$. Under E-admissibility of width δ, a pure target is*

implemented exactly once the posted charge is within the surcharge band $|u_z - u_z^*| \le \hat{s}_z(\delta)$ *of an implementing charge (with a side condition on rival volumes, Section S10), after at most* $\lceil \log(\cdot)/\log(1/\varrho) \rceil$ *rounds.*

Relation to prior work. ARP follows trial-and-error marginal-cost pricing [29, 30, 31], evolutionary implementation of pricing [28] and adaptive incentive design [32, 33], whose fixed points are socially optimal, together with marginal-contribution tariffs. The Lipschitz stability of the equilibrium load in the leader's parameter is standard [cf. 27, Lemma 2.6], whose model-free zeroth-order outer loop gives stationarity rather than a rate. New here are the identification of the equilibrium response as a proximal map with instance-free rates in the lumpy routing game (Theorems S17 and S18), the fixed-point and identification results for the pure game (Theorem S19; part (b) is conditional on potential minimisation), and the absorption of the leader's error by the carriers' credal width (Proposition S20).

S3. Equilibrium theory, full statements

This section states the results on pure and imprecise equilibria; complete proofs are in Section S7. Results labelled *classical* are known and cited.

In words. When carriers choose among a few discrete plans, an equilibrium in pure plans, the kind a city can observe, need not exist. We show when it does (exposure-based congestion) and when deciding existence is hard (vehicle-count congestion). We also show that a carrier who is unsure about its rivals' activity by an explicit margin accepts a rounded version of a mixed equilibrium as an exact equilibrium of its imprecise game.

S3.1. Imprecise beliefs

Assumption A4 (Traffic box)**.** $\Xi = \prod_z [\underline{\mu}_z, \overline{\mu}_z] \subset \mathbb{R}^Z_{\ge 0}$ with widths $\varpi_z = \overline{\mu}_z - \underline{\mu}_z$; a nominal mean $\hat{\mu} \in \Xi$ is fixed and the true mean μ lies in Ξ.

Assumption A5 (Credal sets, clipped and consistent)**.** Fix weights $w \in \mathbb{R}^d_{>0}$ and a width $\delta \ge 0$. For $c \in \mathbb{R}^d_{\ge 0}$ let $K^\delta(c) = \{m \in \mathbb{R}^d_{\ge 0} : |m_k - c_k| \le \delta w_k \ \forall k\}$. At a pure profile σ, carrier i's credal set $\mathcal{M}_i^\delta(\sigma)$ is the set of Borel probability laws P on $\mathbb{R}^d_{\ge 0} \times \mathbb{R}^Z_{\ge 0}$ with finite first moments such that $\mathbb{E}_P[m] \in K^\delta(A_{-i}(\sigma))$ and $\mathbb{E}_P[\xi] \in \Xi$.

The box is centred at the realised rival aggregate, which is the consistency requirement of equilibrium in beliefs [34], and it is clipped at zero, so every belief is a feasible aggregate. The width δ measures how much a carrier is unsure about its rivals' activity, in units w_k.

Definition S21 (Decision criteria; 35)**.** For a finite option set $\mathcal{P}$ and a set $\mathcal{M}$ of laws, with $\underline{\mathbb{E}} = \inf_{P \in \mathcal{M}} \mathbb{E}_P$ and $\overline{\mathbb{E}} = \sup_{P \in \mathcal{M}} \mathbb{E}_P$, an option p is *E-admissible* if it maximises $\mathbb{E}_P[u(\cdot)]$ for some $P \in \mathcal{M}$; *maximal* if no p' has $\underline{\mathbb{E}}[u(p') - u(p)] > 0$; *interval-undominated* (ID) if $\overline{\mathbb{E}}[u(p)] \ge \max_{p'} \underline{\mathbb{E}}[u(p')]$; and Γ-*maximin* if $\underline{\mathbb{E}}[u(p)] = \max_{p'} \underline{\mathbb{E}}[u(p')]$.

Definition S22 (Imprecise credal Nash equilibrium)**.** For $C \in \{E, M, ID, \Gamma\}$ and $\delta \ge 0$, $\mathrm{ICNE}_C(\delta)$ is the set of pure profiles σ in which every σ_i is C-admissible for $(\mathcal{P}_i, \mathcal{M}_i^\delta(\sigma))$ with payoff u_i. The existence threshold is $\delta_0^C = \inf\{\delta \ge 0 : \mathrm{ICNE}_C(\delta) \ne \emptyset\}$.

S3.2. Reduction to mean boxes and the criteria

Write $\bar{u}_i(p; m, \xi)$ for the payoff of Assumption A2 evaluated at the means (m, ξ) and $K_i^\delta(\sigma) = K^\delta(A_{-i}(\sigma)) \times \Xi$.

Lemma S23 (Reduction of the credal set to its mean box)**.** *Under Assumptions A1–A5: (a) for every* $P \in \mathcal{M}_i^\delta(\sigma)$ *and every* p, $\mathbb{E}_P[u_i(p)] = \bar{u}_i(p; \mathbb{E}_P m, \mathbb{E}_P \xi)$*; (b)* $P \mapsto (\mathbb{E}_P m, \mathbb{E}_P \xi)$ *maps* $\mathcal{M}_i^\delta(\sigma)$ *onto* $K_i^\delta(\sigma)$*; (c) hence each criterion of Definition S21 coincides with the same criterion evaluated over the compact box* $K_i^\delta(\sigma)$*, and infima and suprema are attained.*

Lemma S24 (Nesting and monotonicity)**.** *For every $\delta \geq 0$,*

$$\mathrm{ICNE}_E(\delta) \subseteq \mathrm{ICNE}_M(\delta) \subseteq \mathrm{ICNE}_{ID}(\delta) \quad \textit{and} \quad \mathrm{ICNE}_\Gamma(\delta) \subseteq \mathrm{ICNE}_M(\delta).$$

For $C \in \{E, M, ID\}$ and $\delta \leq \delta'$, $\mathrm{ICNE}_C(\delta) \subseteq \mathrm{ICNE}_C(\delta')$. For $C = \Gamma$ the monotonicity fails (Example S40 in Section S7).

The inclusions between criteria are due to Troffaes [35]. Monotonicity in δ says that more imprecision can only enlarge the set of E-admissible, maximal or interval-undominated equilibria; Γ-maximin behaves differently because the worst case moves with δ.

Lemma S25 (Γ-maximin surcharge)**.** *If $\varphi_i(p), t_i(p) \geq 0$, the lower expectation over $K^\delta(c) \times \Xi$ is attained, for every plan simultaneously, at the upper corner $(c + \delta w, \overline{\mu})$ and is linear in mixtures. Consequently* $\mathrm{ICNE}_\Gamma(\delta)$ *is the set of pure Nash equilibria of the precise game G^Γ_δ with private profits*

$$\pi^\Gamma_i(p) = \pi_i(p) - \delta\langle \varphi_i(p), Mw\rangle - \gamma\langle t_i(p), \overline{\mu}\rangle,$$

the same coupling, and traffic set to zero.

The surcharge $\delta\langle \varphi_i(p), Mw\rangle$ is what a cautious carrier adds to the cost of activity in zone k because it fears up to δw_k more rival activity than it observes. Section S5 uses it to compute Γ-maximin equilibria of the city games.

S3.3. Existence under exposure-based congestion

Proposition S26 (Existence under D1 at every width)**.** *Under D1, for every $\hat{\xi} \in \Xi$,*

$$\Psi_{\hat{\xi}}(\sigma) = \textstyle\sum_i \left[\pi_i(\sigma_i) + \frac{1}{2}\|a_i(\sigma_i)\|^2_M - \gamma\langle t_i(\sigma_i), \hat{\xi}\rangle\right] - \frac{1}{2}\|A(\sigma)\|^2_M$$

is an exact potential of the precise game at traffic $\hat{\xi}$. Every maximiser lies in $\mathrm{ICNE}_E(\delta)$ *for all $\delta \geq 0$, so $\delta^E_0 = \delta^M_0 = \delta^{ID}_0 = 0$, and* $\mathrm{ICNE}_\Gamma(\delta) \neq \emptyset$ *for all $\delta \geq 0$.*

Sketch. For a unilateral change of carrier i from contribution a to a' with rival aggregate R, the quadratic terms give $\frac{1}{2}\|a'\|^2_M - \frac{1}{2}\|a\|^2_M - \frac{1}{2}\|R + a'\|^2_M + \frac{1}{2}\|R + a\|^2_M = -\langle a' - a, MR\rangle$, which equals the change of the coupling term when $\varphi_i = a_i$. A maximiser of an exact potential is a pure equilibrium [36], and the centre of every credal box is the realised aggregate. For Γ, G^Γ_δ is again of type D1.

S3.4. Credal absorption

Let x^* be a mixed equilibrium of the precise game at traffic $\hat{\xi} \in \Xi$ (it exists by 37; under affine coupling, mixed equilibria are exactly the profiles in which every plan in the support is a best response to the mean rival aggregate, Proposition S41). Write B_j for carrier j's best-response set at x^*, $\bar{a}_j = \sum_p x^*_j(p) a_j(p)$, $\bar{A}^*_{-i} = \sum_{j \neq i} \bar{a}_j$, and define the spreads $s_{jk} = \max_{p,p' \in B_j} |a_{jk}(p) - a_{jk}(p')|$ and $r_{jk} = \max_{p \in B_j} |a_{jk}(p) - \bar{a}_{jk}|$, with $\bar{\sigma} = \max_{j,k} s_{jk}/w_k$, $\bar{\rho} = \max_{j,k} r_{jk}/w_k$, $d_L = \dim \operatorname{span} \bigcup_j (a_j(B_j) - a_j(B_j))$ and $k^\sharp = \min\{N, d_L\}$. Let $\sigma_{\max} = \max_{i,k} \operatorname{range}_{p \in \mathcal{P}_i} a_{ik}(p)/w_k$, so that $\bar{\rho} \leq \bar{\sigma} \leq \sigma_{\max}$. The *anchored radius* of a pure profile $p \in \prod_j B_j$ is $\hat{\delta}(p) = \max_i \|A_{-i}(p) - \bar{A}^*_{-i}\|_{w,\infty}$.

Theorem S27 (Credal absorption)**.** *Assume A1–A5 and let x^* be a mixed equilibrium at $\hat{\xi} \in \Xi$.*

(a) Absorption. *For every $p \in \prod_j B_j$ and every $\delta \geq \hat{\delta}(p)$, $p \in \mathrm{ICNE}_E(\delta) \subseteq \mathrm{ICNE}_M(\delta) \subseteq \mathrm{ICNE}_{ID}(\delta)$.*

(b) Radius. $\min_{p \in \prod_j B_j} \hat{\delta}(p) \leq \min\left\{ \max_i \max_k \sum_{j \neq i} r_{jk}/w_k,\ k^\sharp \bar{\sigma} + \bar{\rho},\ \bar{\sigma}\sqrt{k^\sharp \ln(2d)/2} + \bar{\rho} \right\}$.

(c) Uniform existence. $\delta^{ID}_0 \leq \delta^M_0 \leq \delta^E_0 \leq \sigma_{\max} \min\{N - 1,\ d + 1,\ 1 + \sqrt{d \ln(2d)/2}\}$.

(d) Construction. *A profile achieving the second bound is obtained from x^* by one linear program (a vertex of the polytope of Shapley–Folkman decompositions) followed by an arbitrary choice for at most $k^\sharp$ carriers; one achieving the third bound by independent sampling of these carriers, with success probability at least $1 - 2d\exp(-2t^2/(k^\sharp\bar{\sigma}^2))$ per draw for target $t + \bar{\rho}$; one achieving the first bound is any $p \in \prod_j B_j$.*

Sketch. Every carrier in $p \in \prod_j B_j$ plays a best response to the *anchor* $\bar{A}^*_{-i}$, which is nonnegative. If $\delta \geq \hat{\delta}(p)$, the anchor lies in the clipped box around the realised aggregate $A_{-i}(p)$, so the Dirac law at the anchor belongs to the credal set and witnesses E-admissibility. The radius bounds follow from the decomposition $A_{-i}(p) - \bar{A}^*_{-i} = \sum_{j\in J}(a_j(p_j) - y_j) - (a_i(p_i) - \bar{a}_i)$, where J is the set of at most $k^\sharp$ carriers left mixed at a vertex of the Shapley–Folkman polytope (Theorem S39); the sum is bounded deterministically by $k^\sharp\bar{\sigma}$ or, after independent sampling, by Hoeffding's inequality [38] and a union bound. The own term $a_i(p_i) - \bar{a}_i$ cannot be removed (it is the "+1" in the bound of 39).

Corollary S28 (ε-Nash in the precise game; cf. 39)**.** *A profile p as in Theorem S27(a) is an ε-Nash equilibrium of the precise game at any $\mu \in \Xi$ with $\varepsilon_i \leq \hat{\delta}(p)L_i(p) + \gamma\sum_z \varpi_z T_{iz}(p)$, where $L_i(p) = \max_{p'}\sum_k w_k M_k|\varphi_{ik}(p') - \varphi_{ik}(p_i)|$ and $T_{iz}(p) = \max_{p'}|t_{iz}(p') - t_{iz}(p_i)|$.*

Remark S29 (Width, regime and relation to prior work). *Width.* Above the explicit belief width $\sigma_{\max}\min\{N-1,\ d+1,\ 1+\sqrt{d\ln(2d)/2}\}$, a mixed equilibrium rounds to an exact pure imprecise equilibrium (Theorem S27(c)).

Regime. On the city instances $N \leq 6$ and $d = 2Z \geq 84$, so $k^\sharp = N$ and the Shapley–Folkman step selects nothing. The applicable term is the support-selection bound $(N-1)\bar{\rho}$, which grows with N, and its $N-1$ factor is attained at $N = 2$ (Proposition S34). The terms $d+1$ and $1+\sqrt{d\ln(2d)/2}$ do not grow with N but are smaller than $N-1$ only when $N > d$, for example with per-vehicle or crowdsourced players; the N-uniform bound is not shown to be attained.

Relation to Liu–Oudjane–Wan. Corollary S28 restates in our notation the approximate equilibrium of Liu et al. [39], who also apply a Shapley–Folkman argument to the generators of a convexified equilibrium, and part (a) follows from their Lemma 5.6; we claim no novelty for either. The addition is the explicit belief box and the reading that, under E-admissibility, the belief width absorbs the rounding error, so that the rounded profile is an exact imprecise equilibrium.

Remark S30 (Γ-maximin). By Lemma S25, $\mathrm{ICNE}_\Gamma(\delta)$ is the pure-equilibrium set of a *precise* game, so absorption does not apply to it, and under D2 that set can be empty (Proposition S34 at $\delta = 0$).

Two further groups of results are stated in Section S7.1. The *tightness* results show that the $N-1$ term of Theorem S27 is attained at $N = 2$ (Proposition S34) and that a dependence on the number of zones is necessary (Proposition S35). The *efficiency* results show that imprecise equilibria are approximate equilibria of the precise game (Lemma S36), that the price of anarchy and of stability are monotone in the belief width (Proposition S37), and that under D1 a potential maximiser bounds the price of stability (Proposition S38). A smoothness-based bound on the price of anarchy also transfers to imprecise equilibria (Proposition S43), but it is conditional, since we have no smoothness constants for the Cournot–congestion game with arbitrary private routing costs; the price-of-anarchy values in Section S5 are therefore empirical.

S3.5. Complexity under vehicle-count congestion

Theorem S31 (NP-completeness under D2)**.** *Consider the problem of deciding* $\mathrm{ICNE}_C(\delta) \neq \emptyset$ *for a game given by explicit plan lists with rational data, M, w, a degenerate traffic box and a rational $\delta \geq 0$. For each $C \in \{E, M, ID, \Gamma\}$ the problem is in NP. For every digraph D with maximum out-degree $\Delta^+ \geq 1$ one constructs in polynomial time a D2 instance with two plans per carrier and one zone per arc such that, for all $\delta \in [0, 1/(2\Delta^+ + 1))$,* $\mathrm{ICNE}_E(\delta) = \mathrm{ICNE}_M(\delta) =$

$\mathrm{ICNE}_{ID}(\delta) = \mathrm{ICNE}_{\Gamma}(\delta)$ *is the set of profiles that play "in" exactly on a kernel of* D*. Hence the problem is strongly NP-complete for each criterion, already at* $\delta = 0$*, and for every fixed* $\delta \in [0, 1/5)$*.*

Sketch. Deciding whether a digraph has a kernel (an independent set that absorbs every other vertex) is NP-complete [40], also for planar digraphs with in- and out-degree at most two [41]. Each vertex becomes a carrier with an "out" plan of zero data and an "in" plan that puts one vehicle on each out-arc zone and two vehicles on each in-arc zone, with exposure one on out-arcs and a small exposure on in-arcs. The gain of "in" is positive if no out-neighbour plays "in" and at most -1 otherwise, and the sign is constant on every credal box of width below $1/(2\Delta^+ + 1)$, so all four criteria select the realised strict best response.

Corollary S32 (Inapproximability of the existence threshold)**.** *Unless* P = NP*, no polynomial algorithm approximates* δ_0^E *of a D2 instance within any fixed factor, or within an additive error below* $1/5$*.*

Pure-equilibrium existence is known to be NP-complete in polymatrix games and integer programming games [42, 43]; Theorem S31 is a direct kernel reduction specialised to the rank-one D2 structure and extended to four credal criteria and a range of widths. Together with Proposition S26 it gives a complexity split between the two congestion regimes: charging or modelling congestion by exposure (vehicle-hours in a zone) yields a potential game with guaranteed pure imprecise equilibria, while counting vehicles does not. Under D2 the mixed equilibrium that Theorem S27 rounds is itself hard to compute: every bimatrix game embeds as a D2 instance with $N = 2$, so the step is PPAD-hard [44, 45] (Proposition S42).

S3.6. Exact thresholds and the rounding algorithm

For any pure profile p, the exact E-threshold of carrier i,

$$A_i^E(p) = \min\{\delta : p_i \text{ is E-admissible at width } \delta\},$$

is the value of a linear program over the box, so $p \in \mathrm{ICNE}_E(\delta)$ if and only if $\delta \geq \max_i A_i^E(p)$ (Theorem S44). Theorem S27 and this threshold computation define an algorithm with four phases: relax (a mixed equilibrium by Frank–Wolfe on the concave relaxed potential under D1, or by a double oracle under D2; 46), sparsify (a vertex of the Shapley–Folkman polytope), round, and compute the exact width. Under D2 its output satisfies the bounds of Theorem S27(b) (Theorem S45), but its relaxation phase is PPAD-hard in general. Under D1 the relaxation-and-rounding technique is classical [47, 48, 49]. The full rounding algorithm was not run on the city games, which use the D1 engine of Section S2; the exact thresholds of Theorem S44 were used in the tests of Section S5.7.

Remark S33 (Nonaffine congestion)*.* With a power congestion term the expected payoff under mixed rival strategies is no longer a function of the mean aggregate; Lemma S23 and the exact credal statements fail, although relaxed equilibria still exist. We do not treat this case.

S3.7. What is machine-checked

The repository contains a Lean 4 development with Mathlib (`lean/CCU`, seven modules) that checks the finite or algebraic cores of several results: the credal results, the exact potential identity under D1, parts of DIP, the marginal tariff, ARP and BRL. It does not formalise probability boxes, convex conjugates, the Hoeffding term, the NP-completeness statement or any computation. Section S12 lists what is checked in full, what only as a core, and what is not formalised.

S4. Data and calibration, extended description

Road networks and zones. For each of nine cities (New Delhi, Cairo, Moscow, Buenos Aires, New York, Mumbai, London, Paris, Berlin) we use an OpenStreetMap extract [50] distributed by BBBike (replication timestamp 16 October 2026). The drivable major-road network (motorway to tertiary, with links) is kept as an undirected graph with class-based free-flow speeds and lane counts (OSM `lanes` tag where present, class defaults otherwise). Population comes from the GHS-POP R2023A grid, epoch 2025, at 30 arc-seconds [51]. Zones are blocks of 4×4 raster cells (about 3.7 km) inside the extract's bounding box with at least one inhabitant; zone-to-zone free-flow times are shortest paths between population-weighted centroids plus a fixed access time. Points of interest from the same extracts give a depot score per zone used to place carrier depots. Table S6 (Section S6) summarises the instances; Z ranges from 42 to 593.

Operating parameters. Per-city parameters were collected from public web sources on 23 October 2026 and graded A (official or primary statistic), B (industry report or country-level proxy) or C (rough estimate or proxy from elsewhere). Table S17 in Section S6 lists the values used; the repository file `data/parameters/SOURCES.md` gives source, URL, grade and note for each number. Congestion levels are those of the TomTom Traffic Index 2025 [52]; volume-delay curves use the BPR form with $\alpha = 0.15$, $\beta = 4$ [53], linearised at the observed congestion level.

Weak numbers. Parcels per capita are country-level figures. Driver wages are minimum-wage floors or job-site averages, not courier payroll. Drops per vehicle and van capacities outside the United States are proxies. Delivery fees in Cairo, Moscow and Buenos Aires are consumer or reference rates, not carrier revenue per parcel, and the European fees are regulator revenue per parcel. Moscow is absent from the 2025 TomTom index and uses a 2020 value. Market shares are available only for Paris and Berlin; elsewhere carriers have equal shares. The sensitivity of the equilibria to the five weakest parameters is in Table S18.

Calibration rules and simplifications. Each carrier serves 22 customer zones sampled around its depot with weight proportional to population times a gravity decay in travel time, so carriers overlap in some zones. A lot is 10–50% of a vehicle's daily drops, proportional to the zone's parcel demand. Inverse demand has intercept $\alpha_z = 1.5\times$ fee and slope $b_z = \alpha_z/(2Q_z^{\text{full}})$. The congestion slope c_z is the derivative of the BPR curve at the TomTom level, with capacity from lane-kilometres, scaled by a slice multiplicity ς_z (the ratio of the city's real delivery vehicle-hours to the modelled ones) so that the slice faces the congestion of the whole city. Travel-time multipliers of nine sectors follow a lognormal law matching the TomTom all-day and peak congestion; each carrier observes 30 days, from which DKW boxes at confidence $1 - \alpha$ are built. The route loss (S3) uses the vehicle-hour cost (wage plus fuel) per hour, twice that as fixed cost, three times that per overtime hour, and the shift length as deadline. The simplifications are:

(C1) all-or-nothing lots: a carrier serves a customer zone fully or not at all, which makes the Cournot game lumpy;

(C2) the integer routing step: the main game experiments use branch-and-price with a time limit per call; the engine comparison and some checks use price-and-branch (Section S5);

(C3) vehicle-hours are proportional to the lot, so within a zone a congestion charge on hours and a delivery charge are collinear;

(C4) linear congestion (D1);

(C5) oracle exactness at city scale is checked against an independent oracle, not proved.

Further proxies are the sampled customer zones instead of whole-city demand, equal market shares in seven cities, the slice multiplicity, and a *broad* social cost that adds the delay imposed on background traffic (population times vehicle ownership times one hour per day, valued at the wage, times the added slowdown, divided by ς_z). The broad objective is reported but not used for conclusions.

Instances as a benchmark. The pipeline (extract download, zones, travel times, depots, parameters with grades, p-box construction, game instances with seeds) is in the repository and regenerates all instances from the raw extracts. The instances are small (22 customer zones per carrier, four to six carriers) so that exact oracles are possible; they are intended as a common test bed for algorithms on multi-carrier routing games, not as validated city models.

S5. Computational study, extended results

Research questions.

RQ1 Is BRL pricing exact, what does its completion window save, and how far does branch-and-price scale? (Section S5.1)

RQ2 Why probability boxes: what does the p-box criterion cost and buy out of sample, compared with sample-average, Wasserstein and CVaR criteria? (Section S5.2)

RQ3 How do equilibria computed with an exact oracle differ from those with price-and-branch, and how do the engine's components compare? (Section S5.3)

RQ4 What do the equilibria, the grid leader, the price of anarchy and the imprecision effects look like on the nine cities, and how stable are they over seeds? (Section S5.4)

RQ5 How do DIP and ARP compare with the grid, and with the social optimum, in oracle calls and gain? (Section S5.5)

RQ6 Are the findings robust to the carriers' criterion? (Section S5.6)

RQ7 Do the theoretical bounds hold on the city equilibria, and does vehicle-count congestion break pure equilibria in practice? (Section S5.7)

Protocol and seeds. All numbers are generated from the result files of the repository by `make_tables.py`. A seed redraws the depots, the customer zones of each carrier, the travel-time sectors and the 30 observed days, hence the probability boxes; calibration inputs are the same in every seed. Seeds 0–2 were run for all nine cities and the exact-oracle study was extended to seeds 3–9, which used 55 046 oracle calls, of which 1.1% were not proven optimal. Coverage is as follows.

- New Delhi, Buenos Aires, London: ten seeds for everything, including the full 63-point exact grid and ARP.
- Cairo, Moscow, New York, Mumbai, Paris, Berlin: 9 seeds (0–8) for the unpriced equilibrium, the social optimum, the price of anarchy and DIP. Grid-derived quantities (congestion-only, delivery-only and combined gains, delivery share, priced gap, DIP against the grid) still have three seeds and a reduced exact grid (7–17 screened points per seed), so they are lower bounds (marked *). ARP was run for Mumbai (five seeds) and not for the other five.
- Imprecision sweeps (three seeds): all 12 widths for New Delhi, Buenos Aires, Mumbai, London; 6 widths for Cairo, Moscow, New York, Paris, Berlin.
- Still computed with price-and-branch: the engine comparison (Table S5), the ZMC checks, the theory tests (Section S5.7), the tiny-game verification, the DIP comparisons for Berlin seed 2, Moscow seed 2, New York seed 2 (price-and-branch grids), the social-optimum benchmark of the broad objective in New York, Moscow and Berlin, and the p-box reference runs of the criterion study (Section S5.6).

Means over seeds are given with percentile bootstrap 95% confidence intervals (10,000 resamples of the seeds). They describe the spread over sampled slices of a city, not the uncertainty in the calibration inputs, which Table S18 examines separately. Of the 41,469 oracle calls of the seeds 0–2 study, 99.17% were proven optimal, and the remaining gap of an unproven call was at most 14 USD. Every reported equilibrium is a pure Nash equilibrium of the game with the exact oracle up to unproven calls (Section S5.8).

S5.1. BRL and branch-and-price

Exactness. The upper-prevision formula agrees with a linear program over all joint laws on 40 random instances to a relative error of $4.5 \cdot 10^{-14}$; the comonotone worst case exceeds the value under independent worst marginals by 4.5% on average and up to 14.1%. BRL pricing equals complete enumeration on every test instance, including 18 of 18 sub-instances built from the nine cities. Fixing one risk level for all routes is suboptimal in 9 of 20 small exact comparisons, by up to 6.42% (Proposition S49).

Public instances: setup. We also ran BRL on Solomon (25, 50, 100 customers) and Augerat A and B instances. They are treated as capacitated routing with a soft duration limit and four sectors with lognormal common-factor uncertainty. This is an emulation. It is not a reproduction of published branch-and-cut results, and the instances of Zhang et al. and Adulyasak and Jaillet could not be obtained.

What BRL does well: label counts. Table S23 compares labelling rules on random instances ($L = 33$ levels, nine sectors). BRL takes 0.004–0.065 seconds. Sector dominance times out from the smallest size, and quantile-vector and scenario labelling from $n = 15$. These random instances are favourable to BRL, and the reductions on real data are smaller (Table S24).

With the duals of the root column generation, the window rule keeps 1.8–2.9 times fewer labels than the same stop-loss comparison without the window. It keeps 2.8–6.3 times fewer than quantile-vector dominance on the metro instances (22–80 customer zones). On the public benchmark instances the factors are 1.6–14 and 2.7–23. The ratios shrink as n grows and are lower bounds where a rule hit its comparison cap. On the public instances scenario labelling keeps 61 123–98 930 labels where BRL keeps 13 175–15 455.

We have not run a general resource-extension-function or function-label labelling code [54] against the completion window; Irnich [10] reports no computational study.

What BRL does well: proof rates. On the nine metro instances with 22 customer zones, branch-and-price proved optimality in 9 of nine cities, in at most 18 minutes on one core (Moscow, 7487 nodes; Table S13). Against complete enumeration on 800 random formulations, branch-and-price returned the optimum in 800 of 800. Price-and-branch was strictly suboptimal in 39 of them, by up to 7.3%.

On the public instances, over 51 cells (instance, number of training days, draw), the BRL solve was proven optimal in 59% of cases. The figures are 35% for the deterministic plan, 20% for SAA, 3.9% for the Wasserstein criterion and 0% for mean-CVaR (90 s limit, same branch-and-price). BRL was proven on Solomon-25 in a median of 3.8 s (27 cells).

What BRL does well: scale. The scale study (Table S7) gives proofs for Cairo, London and New York at 30 zones, and for London at 40 zones (and for New York at 40 zones with the device configuration). Heuristic-first pricing gave a median 2.2-fold speed-up with an identical LP bound; the completion bound gave none.

What BRL does not do: held-out cost. On held-out days from the same law, BRL does not beat the sample-average plan (Table S25). Its plans cost 0.30–5.0% more on the mean and 1.2–2.8% more at the 95th percentile after polishing. After a local-search polish of the best plan of any arm, all arms are within 0.17% of the best known plan under their own criterion. Only in a stress test, with a law shifted away from the training days and ten training days, is BRL's 95th percentile 2.3% lower and its $\mathrm{CVaR}_{0.9}$ 2.5% lower, at 3.3% more on the mean. The advantage vanishes at 30 or more days. BRL's advantage is thus proof speed and label counts against scenario and quantile labelling, not out-of-sample cost.

Table S1: Out-of-sample comparison of criteria on the nine metro instances (mean over 20 draws and cities). Regret: expected cost of the plan above the best known plan for the true law, %. Other columns at $n = 20$: probability that at least one route runs overtime on a day; $\mathrm{CVaR}_{0.95}$ of the daily plan cost above the best known expected cost; criterion value minus realised expected cost.

Criterion	regret % $n = 10$	$n = 20$	$n = 60$	P(overtime day), $n = 20$	$\mathrm{CVaR}_{0.95}$ above best, % ($n = 20$)	criterion minus true cost, % ($n = 20$)
SAA	0.38	0.27	0.16	0.15	38.8	-0.086
Wasserstein, cross-validated radius	0.43	0.26	0.15	0.14	39.1	3.2
Wasserstein, theory radius	1.5	1.1	0.51	0.16	41.1	9.5
$\mathrm{CVaR}_{0.9}$	2.4	2.5	2.9	0.063	34.6	19
p-box, unknown dependence, 0.80	6.0	4.6	1.6	0.069	33.9	29
p-box, unknown dependence, 0.95	7.9	5.1	2.6	0.070	34.5	35
p-box, unknown dependence, 0.99	8.9	6.3	4.1	0.077	35.8	40
p-box, independent, 0.95	5.9	3.8	1.6	0.053	31.9	29
comonotone, empirical marginals (no band)	0.31	0.090	0.051	0.13	37.5	5.3

What BRL does not do: Augerat instances and larger sizes. BRL was proven in only 0 of 15 Augerat cells, and not for Augerat A-n32-k5 (gap 0.25% with 63 047 columns and no cuts). Optimality was not proven at London with 60 or 80 zones or New Delhi with 60. The method should therefore be read as exact where proven (22 customer zones in all nine cities, 30 to 40 customers in London and New York) and not beyond about 40 customers.

Conservatism. On the metro instances the robust optimum is not the best plan under the calibrated law: evaluated under it, the branch-and-price plan is worse than the sample-average plan in Buenos Aires (+7.0%), New York (+4.8%), Berlin (+6.8%), and its worst-case cost lies between 10% and 45% above its expected cost under the law. The next subsection places this in context.

S5.2. Why probability boxes

A natural question is why to use DKW probability boxes with unknown dependence instead of sample-average approximation (SAA), a type-1 Wasserstein ball or CVaR. We compared the criteria out of sample on the nine metro instances (seed 0, 22 customer zones), with $n = 10, 20, 30, 60$ training days, 20 independent draws, and a held-out set of 20,000 days of the instance's law. The planner is a heuristic (Clarke–Wright, local search, ruin and recreate, and a set-partitioning step over all evaluated routes), validated against branch-and-price: optimal in 8 of 9 instances, at most 0.10% above in the others (Paris). Table S1 gives the results.

The p-box loses in expected cost. At every $n \leq 60$ its regret is 1.6–8.9% (for confidence 0.80 to 0.99, falling with n), against 0.27% for SAA at $n = 20$ and 0.16% at $n = 60$. Wasserstein with a cross-validated radius reproduces SAA (0.26% and 0.15%); at the theory radius it costs 1.1% and 0.51%; $\mathrm{CVaR}_{0.9}$ costs 2.5% and 2.9%. The cost comes from the DKW band, not from the unknown-dependence worst case: the comonotone criterion with empirical marginals and no band has regret 0.090% and 0.051%.

It buys tail protection and a bound. At $n = 20$ the probability of an overtime day is 0.15 for SAA and 0.069 for the p-box with confidence 0.80 ($\mathrm{CVaR}_{0.9}$: 0.063), and the $\mathrm{CVaR}_{0.95}$ of the daily cost is 3.6–5.0 percentage points lower than under SAA at every n. Under a common additive congestion shock at test time, the p-box plan with confidence 0.80 has the same expected cost as the SAA plan at a shock of about 0.50 ($n = 60$) to 0.60 ($n = 20$) multiplier units (multipliers of the instances are about 1.4–1.6) and is cheaper beyond (2.7% at a shock of 1.0, $n = 60$). Its criterion value is an upper bound on the true expected cost whenever the DKW band holds: the criterion exceeds the realised cost by 20–40% for the p-box, by 3.2% for the cross-validated Wasserstein criterion and by -0.086% for SAA at $n = 20$. This is the conservatism referred to above.

What the criterion provides. P-boxes do not minimise expected cost. They provide a guaranteed upper bound that shrinks with the sample size, and tail protection, at an expected-cost price that is 1.6–8.9% for $n \leq 60$. Which criterion the carriers use matters for the game results; Section S5.6 quantifies that.

S5.3. Equilibrium engine and exactness of the oracle

Effect of the inexact integer step. The seeds 0–2 experiments were first computed with price-and-branch. Tested against a fresh exact oracle, those base equilibria were exact in New Delhi, Buenos Aires, Mumbai, London in every seed but not in 10 of the 27 city-seeds, where a carrier could gain at most 4.0 USD per day with a plan the pooled integer program had not generated. City-mean welfare changes by at most 0.14% outside Buenos Aires and by -1.4% in Buenos Aires, where seed 0 moves to a different equilibrium (-5.4%); single seeds elsewhere change by up to 0.53% with both signs. The exact oracle changes which equilibrium is reached rather than improving welfare systematically; it takes three to twelve times longer.

Engine components. On the city games (Table S5, price-and-branch oracle) ZMC uses 36–65 calls, about ten per carrier, and its potential gap coincides with those of Frank–Wolfe and column generation with eight rounds to three digits in most cities. Plain best response needs 8–17 calls and is as good as ZMC (within 10^{-3}) in Cairo, Moscow, New York, Paris, Berlin and worse in New Delhi, Buenos Aires, Mumbai, London. ZMC therefore does not beat best response on these games: only 32–89 coordinates are active, so the dimension that favours ZMC in the synthetic instances is absent. Its dimension independence is not tested at city scale.

S5.4. Unpriced equilibria and the price of anarchy

The cities fall into two groups (Table S8, seeds 0–2, exact oracle). In Cairo, Moscow, New York, Paris, Berlin the carriers serve at least 85% of customer demand; in New Delhi, Buenos Aires, Mumbai, London they serve 18–45%, because the delivery fee is low relative to the vehicle-hour cost or the shift is short. The added slowdown caused by delivery vehicles is 0.77–17.6% (Moscow highest).

Price of anarchy over ten seeds. The social optimum is approximated by column generation on the convexified welfare problem followed by rounding and polish. The equilibria considered are those found by the engine and by best response from the empty and full-service profiles and from three random price vectors. The price of anarchy is therefore *empirical*: a ratio of the rounded optimum to the equilibria found, not a bound over all equilibria. Table S2 gives the ten-seed results. Among the engine's equilibria the price of anarchy is at least 1.10 in some seed only in New Delhi, Buenos Aires, London. Its mean is 1.53 in London, 1.38 in Buenos Aires and 1.18 in New Delhi, with ranges 1.25–1.94, 1.28–1.46 and 1.09–1.31. In the other cities it is at most 1.06 in mean (Mumbai's worst equilibrium found has mean 1.10 [1.08, 1.13]). The means are ordered London > Buenos Aires > New Delhi, but the ordering holds seed by seed only in 7 of 10 seeds: London is below Buenos Aires in seeds 4, 6, 7, and Buenos Aires exceeds New Delhi in 10 of 10. With the grid's best charges the equilibrium is much closer to the optimum, with a priced gap of at most 7.4% in any seed. The relaxation bound exceeds the rounded optimum by up to 25% (London), so the true price of anarchy could be larger. Table S10 (Section S6) gives the seeds 0–2 detail.

Imprecision. Two kinds of imprecision are varied (seeds 0–2; Table S11, Figure S2). Widening the travel-time probability boxes from $\varepsilon = 0$ to $\varepsilon = 0.41$ acts through the BRL route costs: deliveries fall by 0–78% and welfare by 2.7–82% at the widest box. Cairo loses no deliveries, Moscow and New York a few per cent, Paris and Berlin somewhat more, Mumbai more, and New Delhi, Buenos Aires and London about half or more; the low-service magnitudes vary by seed (London: -60.0 / -56.1 / -78.2% in seeds 0, 1, 2). Widening the credal width about rivals under Γ-maximin from $\delta = 0$ to $\delta = 0.80$ lowers deliveries by 31–100% and welfare by 22–100%. Deliveries are not always monotone in the width: 3 of 54 series contain a rise, all in Buenos Aires; Proposition S37 does not apply to Γ-maximin or to a single equilibrium that can switch. Imprecision acts like a tax on activity; because vehicle-hours are proportional to deliveries (C3), the model cannot separate a congestion effect from a delivery effect. The imprecision sweeps were not extended to ten seeds.

Table S2: Ten-seed summary (exact oracle). n_{eq}, n_{grid}: seeds for the equilibrium quantities and for grid-derived quantities; *: reduced exact grid, grid quantities are lower bounds. PoA of the engine's equilibrium, mean with bootstrap 95% CI. Gains are % of |social cost| of the best exact grid point with the congestion charge only or the delivery charge only: mean (maximum over seeds). Delivery share: delivery-only gain over the combined gain per seed. Priced gap: shortfall of the equilibrium under the grid's best charges from the rounded optimum, %.

City	n_{eq}	n_{grid}	PoA mean [95% CI]	cong. only % mean (max)	deliv. only % mean (max)	delivery share [95% CI]	priced gap % mean (max)
New Delhi	10	10	1.18 [1.14, 1.23]	0.32 (2.9)	12.1 (21.2)	0.814 [0.687, 0.920]	2.9 (6.7)
Cairo*	9	3	1.00 [1.00, 1.00]	0 (0)	0 (0)	–	0 (0)
Moscow*	9	3	1.00 [1.00, 1.00]	0 (0)	0.215 (0.387)	–	0.0041 (0.012)
Buenos Aires	10	10	1.38 [1.33, 1.42]	2.4 (6.5)	34.6 (41.9)	0.978 [0.963, 0.991]	1.8 (3.2)
New York*	9	3	1.00 [1.00, 1.00]	0.0045 (0.013)	0.352 (0.460)	–	0 (0)
Mumbai*	9	3	1.06 [1.04, 1.07]	0 (0)	2.09 (3.31)	0.877 [0.464, 1.38]	1.0 (1.5)
London	10	10	1.53 [1.41, 1.66]	1.1 (5.4)	39.8 (69.7)	0.816 [0.752, 0.880]	3.4 (7.4)
Paris*	9	3	1.03 [1.03, 1.04]	0.041 (0.12)	3.47 (4.72)	0.970 [0.910, 1.00]	0.16 (0.23)
Berlin*	9	3	1.02 [1.02, 1.03]	0 (0)	1.85 (2.17)	0.943 [0.904, 1.00]	0.19 (0.46)

S5.5. Leader methods: grid, DIP and ARP

The grid. The baseline leader chooses a congestion charge per vehicle-hour, $t_z = \vartheta \gamma c_z X_z^{\text{NE}}$ with $\vartheta \in \{0, 0.5, 1, 2, 4, 8, 16\}$, and a per-delivery charge $s \in \{-0.6, \ldots, 0.6\}$ times the fee (negative: subsidy), a two-parameter family of 63 points with one warm-started equilibrium each. Table S2 and Table S9 (Section efapp:tables, seeds 0–2) give the reduction of social cost for the best exact point using only the congestion charge, only the delivery charge, or both.

Congestion charge alone. It reaches at least 1.7% of the social cost in 4 of 10 seeds in Buenos Aires (maximum 6.5%), 2 of 10 in London (maximum 5.4%) and 1 of 10 in New Delhi (maximum 2.9%). In the seeds 0–2 exact run the Buenos Aires seed-0 case (6.51%) arose because the exact oracle selected a worse unpriced equilibrium (welfare 252.2 against 266.7 USD with price-and-branch). The comparison is stable: the mean congestion-only gain is 0.32–2.4% in the three full-grid cities, against 12.1–39.8% for the delivery charge alone.

Per-delivery instrument. In every case with a material gain the best point has a subsidy. The delivery charge alone achieves 0.81 [0.69, 0.92] of the combined gain in New Delhi, 0.98 [0.96, 0.99] in Buenos Aires and 0.82 [0.75, 0.88] in London (ratio of the per-seed shares; bootstrap 95% CI), and below one half only in New Delhi seed 0. Mumbai's share is not established (3 seeds, CI 0.46–1.4). The gain is material only where the unpriced equilibrium under-delivers; in Cairo, Moscow and New York it is at most 0.46%. In London the combined gain ranges widely over seeds. The mechanism visible in the model is under-provision caused by fixed route costs and market power, which a subsidy corrects, while the congestion externality among carriers is small; within a zone the two instruments are collinear (C3). Subsidies cost the city money; since charges are transfers in the welfare definition, this fiscal cost is not part of the social cost. With the broad objective the leader's best response in New Delhi, Mumbai, London is to stop deliveries, an artifact of the proxy that supports no conclusion (Table S19, Section efapp:tables).

S5.5.1. DIP against the grid and against the social optimum

DIP was run with the exact oracle on all nine cities, seeds 0–8 or 0–9 (Table S3). Calls count every routing-oracle query of each method with the same counter. DIP searches a richer family than the grid (one charge per zone against two parameters), so a gain share above one against the grid is not a like-for-like win; restricted to the grid's family, DIP reaches on average 1.0 of the grid gain in the seeds 0–2 runs, with the worst case 0.50 (Mumbai, seed 0, a reduced grid). Table S3 therefore also reports every method's gain as a share of the gain of the rounded social optimum.

Gain and cost, ten seeds. Ten seeds are available for the three full-grid cities, New Delhi, Buenos Aires and London (Table S3). DIP reaches 1.23, 1.07 and 1.14 of the best grid point's gain in the three cities. It uses 5.7%, 6.8% and 8.3% of a 63-point grid's calls.

Table S3: Leader methods, mean over seeds (exact oracle). DIP / grid: DIP gain over the best exact grid point's gain; calls: DIP's oracle calls as a percentage of a 63-point grid at the same calls per point. Gain shares against the social optimum (SO): best grid point, DIP, ARP flat; ARP / DIP gain ratio; n_{ARP} seeds with an ARP run. *: reduced exact grid (lower bounds for the grid column). Shares are not reported where the optimum's gain is below 0.5% of |social cost|.

City	DIP / grid	DIP calls % of 63-pt grid	grid / SO	DIP / SO	ARP / SO	ARP / DIP	n_{ARP}
New Delhi	1.23	5.69	0.815	1.00	0.933	0.933	10
Cairo*	–	2.33	–	–	–	–	0
Moscow*	–	2.53	–	–	–	–	0
Buenos Aires	1.07	6.85	0.938	1.00	0.995	0.995	10
New York*	–	2.73	–	1.00	–	–	0
Mumbai*	1.91	6.13	0.711	1.05	0.947	0.884	5
London	1.14	8.33	0.905	1.02	0.926	0.912	10
Paris*	1.04	4.69	0.959	1.00	–	–	0
Berlin*	1.05	3.55	0.949	1.00	–	–	0

Against the social optimum, ten seeds. The best grid point reaches 0.815, 0.938 and 0.905 of the social optimum's gain in the same three cities. DIP reaches 1.00, 1.00 and 1.02. Values above one occur where DIP's multi-start target is better than the benchmark. The grid's shortfall is the price of its two-parameter family. DIP's near-optimality is by construction the quality of its social target.

Gain and cost, seeds 0–2. In the same three full-grid cities, three seeds give 104–135% of the grid's gain with 4.9–9.4% of its calls (105–221 calls against 1 921–2 419). In Cairo no charge improves on the unpriced equilibrium.

Equilibrium check, seeds 0–2. In all 27 runs the cutting-plane loop ended with charges under which the target is an exact pure equilibrium (fresh-oracle regret at most $9.1 \cdot 10^{-13}$ USD). The tangent charge alone sufficed in 18. The uniform subsidy is 0.37–0.50 of the fee, and the zone charges follow the competitor-count rule of Proposition S15.

Qualifiers, cost. The per-run detail for seeds 0–2 is in Table S12 and Figure S5 (Section S6). Against the reduced exact grids, DIP uses more than 10% of their calls in most runs (Cairo 8.5–12%, Moscow 13%, New York 23%, Mumbai 24–31%, Paris 21–23%, Berlin 15–17%).

Qualifiers, equilibrium selection. Wall-clock time is not always lower (median ratio 0.22; London seed 0: 247 s against 157 s). Selection is optimistic: under DIP's charges the grid's protocol reaches the target in 21 of 27 runs, and in 210 enumerated tiny games the equilibrium under DIP's charges was unique in only 35% of the games. In the tiny games DIP reached the exact leader optimum over all zone charges in 93% (100% with the exact social optimum as target), so each miss was a miss of the social solver.

Qualifiers, Γ*-maximin.* With Γ-maximin carriers, the shifted charges of Theorem S14 kept the target an exact equilibrium in every run at $\delta = 0.2$ and 0.8 (largest regret $9.1 \cdot 10^{-13}$ USD), but at $\delta = 0.8$ the required uniform subsidy is 0.98–3.6 times the fee, which is not a realistic policy.

S5.5.2. ARP: charges from observed aggregates

ARP was run with the exact oracle for ten seeds in New Delhi, Buenos Aires and London and five in Mumbai. The carriers best-respond day to day from their current plans after each posted charge, and all their oracle calls are counted; the status-quo equilibrium (40–60 calls) is common to all methods and not counted, as for DIP. Table S26 gives the seeds 0–2 runs in detail.

City runs, seeds 0–2. ARP reaches 90.8% to 293% of the grid's gain (mean 125%) after 3–7 posted charges and 22–67 calls. DIP reaches 104–325% (mean 134%) with 105–221 calls. In the three full-grid cities the three-seed figures are 90.8–126% for ARP and 104–135% for DIP.




Figure S1: Leader methods on four cities and three seeds (exact oracle): gain relative to the best exact grid point against routing-oracle calls (log scale).

City runs, ten seeds. In the three full-grid cities over ten seeds, ARP's gain is 0.933, 0.995 and 0.926 of the social optimum's gain (Table S3). That is 0.93, 0.99 and 0.91 of DIP's gain. ARP reached an exact fixed point in 15 of 30 runs.

Where ARP falls short, seeds 0–2. ARP matches DIP within two percentage points in 5 of 12 runs. In the other 7 it is below DIP by up to 32 points of the grid's gain. The largest gaps occur where the iteration cycles or the grid is reduced. The clear miss is New Delhi seed 0 (90.8% against DIP's 123%), where the iteration cycles between two equilibria and $\varrho = 0.40$, the largest of the sample.

Convergence, seeds 0–2. The flat rule reached an exact fixed point in 6 runs and cycled in 6. Successive averages do not remove the cycles. The relative change of the charge vector falls to 0.37–0.63 of its initial value after the first round and to 0.044–0.24 after the second.

Marginal tariff, seeds 0–2. The marginal tariff with the carriers' own best responses reaches 98.7–293% (mean 125%) with 8–14 calls and no city-side computation, and never raised the social cost along the way (12 of 12 runs), as Theorem S16 says; with the potential solver it reaches 92.5–263% with 39–51 calls. Every returned profile is an exact pure equilibrium of the posted charges (regret at most $1.1 \cdot 10^{-13}$ USD for ARP and $1.4 \cdot 10^{-14}$ USD for the tariff). ARP's wall-clock time is 1.9–41 s, a median of 0.041 of DIP's.

Tiny games and contraction constants. On 210 exhaustively enumerated games (136 with a positive leader gain; Table S14, Section efapp:tables) the flat rule reached a fixed point in 166 games and cycled in 44, almost all at strong congestion; its mean share of the exact optimal gain is 0.72 / 0.64 / 0.23 at congestion multipliers 1, 20, 60 (tariff 0.93 / 0.79 / 0.58, DIP 1.00 / 0.96 / 0.93). Its failures are entry into zones that the optimum leaves unserved, attracted by the subsidy based on the city-mean lot, and two-charge cycles. Knowing the lots does not help (share 0.63): the limit is what the aggregate says about which profile is socially good (Theorem S19(c)). Under the tariff the pure equilibria are exactly the unilateral social optima (the best tariff equilibrium is the social optimum in 210 of 210 games). On 48 convexified tiny games with ϱ from 0.0062 to 0.81 the Lipschitz bound of Theorem S17, the plain-step bound and the damped-step bound of Theorem S18 (tight, observed maximum 1.00 of the bound) held in all games. ARP is not better than DIP in solution quality; it is better in the information it needs and in cost.

Table S4: Game results under four carrier criteria, three seeds, mean. PoA with the true expected routing cost and with the carriers' own (model) cost. DIP gain as % of |social cost| under each cost. Best uniform per-delivery charge on the coarse grid as a fraction of the fee. Weighted DIP charge: u_z/fee weighted by deliveries.

City	carriers' criterion	served share	PoA (true cost)	PoA (model cost)	DIP gain %, true cost	DIP gain %, model cost	best uniform charge / fee, true cost	same, model cost	DIP charge u/fee (weighted)
New Delhi	p-box	0.26	1.35	1.18	34.0	18.2	-0.53	-0.20	-0.17
New Delhi	SAA	0.48	1.19	1.21	18.4	20.7	-0.17	-0.23	-0.18
New Delhi	Wass. CV	0.46	1.19	1.20	18.5	19.4	-0.20	-0.20	-0.18
New Delhi	Wass. theory	0.41	1.25	1.20	24.6	19.2	-0.27	-0.20	-0.18
Buenos Aires	p-box	0.37	1.49	1.36	51.2	35.9	-0.77	-0.30	-0.24
Buenos Aires	SAA	0.67	1.16	1.19	16.0	18.5	-0.30	-0.30	-0.25
Buenos Aires	Wass. CV	0.58	1.27	1.23	26.6	23.5	-0.37	-0.23	-0.25
Buenos Aires	Wass. theory	0.57	1.31	1.28	30.4	27.9	-0.30	-0.40	-0.25
London	p-box	0.18	2.06	1.66	108	65.8	-0.80	-0.30	-0.25
London	SAA	0.38	1.52	1.55	51.1	53.2	-0.33	-0.33	-0.26
London	Wass. CV	0.32	1.69	1.65	68.0	63.8	-0.47	-0.40	-0.26
London	Wass. theory	0.31	1.75	1.66	71.8	65.4	-0.40	-0.30	-0.26

S5.6. Robustness to the carriers' criterion

All game results above describe carriers who plan against the p-box worst-case route cost. To test how much this matters we recomputed the unpriced equilibrium, the price of anarchy, DIP and a coarse 16-point grid for carriers who plan with the sample-average cost of the same 30 days, with a Wasserstein ball with cross-validated radius, or with the theory radius, for all nine cities and 3 seeds. The SAA and Wasserstein carriers use a heuristic oracle without an optimality proof (equilibria verified against an independent fresh pool, regret at most $5 \cdot 10^{-13}$); the p-box reference uses price-and-branch; the coarse grid is easier to match than the 63-point grid; the cross-validated Wasserstein criterion was run for one seed only in Paris and Berlin. Table S4 gives the three low-service cities.

What is robust. Under all four criteria: the delivery charge alone achieves 0.88–1.0 of the best grid gain (0.74–1.00 for the theory radius); the congestion charge alone is small, though marginal for SAA in Buenos Aires (3.3% against 18% for the delivery charge); the price of anarchy is at least 1.10 only in London, Buenos Aires and New Delhi (at most 1.06 elsewhere with SAA carriers); DIP's per-delivery charge follows $-0.375\,\text{fee}/n_z$ (slopes 0.361–0.379, correlation at least 0.984 in absolute value); DIP matches the coarse-grid gain (ratio at least 0.99) with 5.0–10% of the calls of a 63-point grid (21–43% of the coarse grid's); the optimal charge is a subsidy in every city; and cities with complete service are unchanged.

What is not robust. The levels. With SAA carriers the unpriced served share roughly doubles (London 0.18 to 0.38, Buenos Aires 0.37 to 0.67, New Delhi 0.26 to 0.48) and the price of anarchy measured with the true cost falls (London 2.06 to 1.52, Buenos Aires 1.49 to 1.16, New Delhi 1.35 to 1.19); the model-cost price of anarchy moves little. The ordering of the price of anarchy is not preserved: with SAA carriers it is 1.16 and 1.19 in Buenos Aires and New Delhi. The leader's gain under the true cost falls by half or more (London 108% to 51.1%, Buenos Aires 51.2% to 16.0%, New Delhi 34.0% to 18.4%), and the best uniform charge for the true-cost objective shrinks by 58–69% (p-box 0.80, 0.77, 0.53 of the fee, beyond the -0.6 end of the grid axis, against 0.33, 0.30, 0.17 for SAA), while DIP's zone charges themselves are almost unchanged. *All levels of the price of anarchy, served share and gain in this paper describe carriers planning against a guaranteed upper bound; the orderings and the policy directions are robust, the levels are not.*

S5.7. Theory against data

The city games are of type D1, so the engine's profile is an exact pure equilibrium with E-threshold zero and Theorem S27 holds trivially for it. To test the theorems non-trivially we computed the exact thresholds of Theorem S44 on mixed equilibria with belief weights equal to the realised aggregate (price-and-branch oracle; each threshold is relaxed by the largest improvement observed at a verified equilibrium). Table S15 and Figure S6 are in Section S6.

Credal absorption. Non-trivial mixed equilibria were constructed from pairs of distinct pure equilibria by solving the two-plan indifference conditions and checking that every support plan is a best response over all plans. This succeeded in 14 of the 27 city-seed games; Cairo, Moscow

have none, because almost all demand is served. On all 274 tested profiles no statement of Theorem S27 was violated (0 violations). The bound is loose: the largest empirical threshold is 0.28–0.60 times the anchored bound (median 0.45). As Remark S29 states, on these games $N \leq 6 < d$, so the $(N-1)\bar{\rho}$ term is the smallest of the three in every case; the N-free terms are not tested by the city data. Without the best-response polish, the rounded profile of the first engine phase is E-admissible at a finite width in only 10 of 27 games. Limits: mixed equilibria were found only with two-plan supports, and thresholds are exact only up to the oracle gap.

Γ-*maximin.* In all 108 (city, seed, δ) cases the surcharge equilibrium differs from the precise equilibrium, and in 106 of them the precise equilibrium is not Γ-admissible. The share of demand served drops by 0–0.082 at $\delta = 0.1$ and by 0.17–0.77 at $\delta = 1$.

Vehicle-count congestion. The gadgets behave as proved: Proposition S34 has no pure equilibrium and threshold exactly $1/w$ (3 of 3 weights), and the kernel construction reproduces the kernel sets for all four criteria on 223 of 223 digraphs. Among random bilinear D2 games with three players, 6 of 400 have no pure equilibrium, against 0 of 150 D1 controls. Small routing-integrated D2 games always had a pure equilibrium (0 of 540), and on the city games of Cairo, Mumbai and Paris with a vehicle-count externality scaled by 1, 10, 100, 1 000, 10 000, all 180 best-response runs converged. Non-existence required adversarial plan menus; the D2 calibration is an assumption introduced for this test.

S5.8. Verification

On 40 tiny games solved by exhaustive enumeration, the engine's profile is a pure equilibrium in every case and reaches the global potential minimum in 40 of 40 (best response 35, Frank–Wolfe with rounding 38, column generation 39; Table S21). At city scale every base equilibrium, every verified grid best point (27 city-seeds) and every equilibrium of the price-of-anarchy experiment has regret at most $9.1 \cdot 10^{-13}$ USD against an independent fresh exact oracle. Because some oracle calls were not proven optimal, the guaranteed regret bound is nonzero for 10 profiles, at most 4.84 USD. These checks rely on the same branch-and-price method as the oracle; the pricer was checked against enumeration (Section S5.1).

S6. Additional tables

Table S5: Methods on the nine city games (seed 0, price-and-branch oracle). d_{eff}: active coordinates. Oracle calls and relative potential gap to the convexified optimum, in %, for best response (BR), ZMC with eight rounds, Frank–Wolfe and column generation with eight rounds (FW, CG8) and column generation to convergence (CG); all methods end with the same best-response polish.

City	N	Z	d_{eff}	BR	ZMC	FW	CG8	CG	BR	ZMC	FW	CG8	CG
New Delhi	5	120	64	14	45	45	50	65	13.8	10.2	10.2	10.2	10.2
Cairo	4	42	32	8	36	24	16	16	$1.0 \cdot 10^{-5}$	$1.0 \cdot 10^{-5}$	$1.0 \cdot 10^{-5}$	$1.0 \cdot 10^{-5}$	$1.0 \cdot 10^{-5}$
Moscow	5	160	71	12	45	45	45	60	0.220	0.220	0.220	0.220	0.220
Buenos Aires	4	208	62	10	39	37	40	91	17.9	16.3	16.6	16.0	16.0
New York	4	234	64	8	36	36	36	76	0.451	0.451	0.437	0.437	0.437
Mumbai	5	143	55	14	48	48	48	60	4.60	1.98	1.98	1.98	1.98
London	5	348	89	15	49	49	50	70	20.0	18.0	18.0	18.0	18.0
Paris	5	198	68	13	50	50	50	108	2.74	2.78	2.74	2.74	2.74
Berlin	6	593	82	17	65	59	59	143	2.32	2.43	2.32	2.46	2.46




Figure S2: Imprecision, exact oracle, mean over the seeds (line) with the minimum–maximum band. Top: probability-box half-width (route costs through BRL). Bottom: Γ-maximin credal width.




Figure S3: Gain of the grid leader's best exact charge over no charge, by instrument: mean over the seeds with standard-deviation bars and minimum/maximum ticks. Left: main objective. Right: broad objective with the background-delay proxy (shut-down optima are an artifact). Cairo, Moscow, New York, Mumbai, Paris, Berlin use a reduced exact grid (lower bounds).

Table S6: City instances. Z: zones; population from GHS-POP inside the zones; major road and lane kilometres inside the zones; nodes of the major-road graph; N: number of carriers.

City	Z	pop. (M)	major road km	lane-km	graph nodes	N
New Delhi	120	25.0	5 556	8 590	138 542	5
Cairo	42	14.9	2 649	4 498	119 027	4
Moscow	160	12.6	4 475	12 325	146 234	5
Buenos Aires	208	13.3	5 764	12 032	130 526	4
New York	234	12.1	8 798	17 770	382 014	4
Mumbai	143	22.0	4 259	7 092	109 804	5
London	348	11.3	7 110	13 628	350 931	5
Paris	198	9.02	5 990	10 854	309 175	5
Berlin	593	4.65	5 490	11 147	192 577	6




Figure S4: Unpriced equilibrium zone prices divided by the city's delivery fee, exact oracle, seed 0 (light grey: zones without deliveries).

Table S7: Full branch-and-price as the instance grows (1800 s limit). Metro instances with n customer zones, one Solomon and one Augerat instance. Gap: relative gap at the limit; no valid bound: a pricing call hit the comparison cap. Devices: ng-4 start with cycle-driven enlargement, heuristic-first pricing, completion bound.

instance	n	ng-8, exact pricing	devices (ng-4 start, heuristic-first pricing)
augerat_A/A-n32-k5	31	limit, gap 0.25%	limit, gap 0.30%
metro/Cairo	30	optimal, 17.2 s	optimal, 20.2 s
metro/Cairo	40	limit, gap 0.0041%	limit, gap 0.0032%
metro/London	30	optimal, 12.9 s	optimal, 14.4 s
metro/London	40	optimal, 559 s	optimal, 234 s
metro/London	60	limit, no valid bound	limit, no valid bound
metro/London	80	limit, no valid bound	limit, no valid bound
metro/NewDelhi	60	limit, no valid bound	limit, no valid bound
metro/NewYork	30	optimal, 36.0 s	optimal, 14.8 s
metro/NewYork	40	limit, no valid bound	optimal, 1 387 s
solomon25/R101	25	optimal, 8.39 s	optimal, 1.72 s

Table S8: Unpriced equilibria with the exact oracle (mean over three seeds; seed standard deviation in parentheses). Served: share of customer demand delivered; slowdown: added mean travel-time slowdown; price: mean zone price (USD per parcel); veh.-h: vehicle-hours per day; welfare W in USD per day of the slice; calls: oracle calls; regret^{LP}: guaranteed upper bound on any carrier's gain from deviating (USD), nonzero only where an oracle call was not proven optimal; η_{rel}: upper bound on the relative potential gap.

City	N	Z	served	slowdown %	price	veh.-h	welfare	calls	regret^{LP}	η_{rel}
New Delhi	5	120	0.26 (0.01)	4.2	0.803	49.1	82.7 (7.0)	48	$7.1 \cdot 10^{-15}$	0.12
Cairo	4	42.0	1.0 (0)	3.6	0.885	125	1 066 (20)	39	0.0020	$-2.4 \cdot 10^{-4}$
Moscow	5	160	0.99 (0.004)	18	2.76	170	6 756 (154)	47	1.2	0.0036
Buenos Aires	4	208	0.36 (0.03)	0.77	0.941	43.6	272 (19)	39	$2.1 \cdot 10^{-14}$	0.24
New York	4	234	0.99 (0.002)	7.8	6.82	112	20 003 (1 441)	39	4.8	0.0050
Mumbai	5	143	0.45 (0.02)	4.6	0.751	83.1	153 (10)	47	$1.8 \cdot 10^{-14}$	0.046
London	5	348	0.18 (0.03)	6.7	4.67	24.8	570 (101)	48	$1.1 \cdot 10^{-13}$	0.44
Paris	5	198	0.85 (0.04)	5.0	4.66	105	5 332 (437)	50	0.76	0.039
Berlin	6	593	0.89 (0.01)	3.7	3.36	127	8 619 (256)	58	0.90	0.029

Table S9: Grid leader, main objective, exact oracle (3 seeds). Gain of the best charge over no charge (% of |social cost|), mean [min, max] over seeds, with the congestion charge only, the delivery charge only, or both. Delivery share: gain with the delivery charge alone divided by the gain with both, per seed, shown where the combined gain is at least 0.5%. Subsidy: seeds in which the best point with both instruments has $s < 0$. Served share and city revenue (USD/day; negative is a net subsidy) at that point. †: only a subset of the grid solved exactly, so gains are lower bounds and shares are ratios of lower bounds.

City	cong. only	deliv. only	both	deliv. share by seed	subsidy	served	revenue
New Delhi	0	10.3 [5.0, 17]	14.5 [12, 17]	0.41 / 0.97 / 0.65	3/3	0.39 [0.36, 0.42]	-54.7 [-64, -48]
Cairo†	0	0	0	–	0/3	1.0	0
Moscow†	0	0.215 [0.13, 0.39]	0.215 [0.13, 0.39]	–	3/3	1.00 [1.00, 1.0]	-2 411 [-3 375, -747]
Buenos Aires	2.49 [0.43, 6.5]	35.4 [27, 40]	35.4 [27, 40]	1.0 / 1.0 / 1.0	3/3	0.74 [0.65, 0.79]	-413 [-528, -226]
New York†	0.00448 [0, 0.013]	0.352 [0.27, 0.46]	0.352 [0.27, 0.46]	–	3/3	1.0	-4 024 [-8 085, -1 891]
Mumbai†	0	2.09 [0.37, 3.3]	2.16 [0.81, 3.3]	0.46 / 1.4 / 0.79	3/3	0.55 [0.49, 0.58]	-64.4 [-91, -20]
London	0.446 [0, 1.3]	49.3 [31, 67]	59.7 [34, 90]	0.92 / 0.89 / 0.74	3/3	0.49 [0.39, 0.54]	-1 359 [-1 803, -808]
Paris†	0.0412 [0, 0.12]	3.47 [2.8, 4.7]	3.57 [2.8, 4.7]	1.0 / 0.91 / 1.0	3/3	0.95 [0.89, 0.99]	-2 712 [-4 160, -849]
Berlin†	0	1.85 [1.6, 2.2]	1.98 [1.6, 2.4]	0.93 / 1.0 / 0.90	3/3	0.94 [0.93, 0.97]	-3 334 [-5 258, -1 921]

Table S10: Empirical price of anarchy and stability, main objective, exact oracle (3 seeds). W_{SO}: rounded social optimum (USD/day), mean [min, max]; #NE: distinct equilibria found; PoA of the engine's equilibrium, of the worst equilibrium found, and PoS (best found), per seed; priced gap: shortfall of the equilibrium under the grid's best charges from W_{SO} (%), per seed.

City	W_{SO}	#NE	PoA (ZMC eq.)	PoA (worst NE)	PoS	priced gap %
New Delhi	97.7 [86, 106]	4–5	1.15 / 1.23 / 1.16	1.15 / 1.24 / 1.20	1.12 / 1.23 / 1.16	2.4 / 4.9 / 1.6
Cairo	1 066 [1 046, 1 086]	2–3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00	0 / 0 / 0
Moscow	6 771 [6 639, 6 929]	2	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00	0.012 / 0 / 0
Buenos Aires	375 [361, 390]	3–6	1.43 / 1.42 / 1.29	1.43 / 1.54 / 1.31	1.32 / 1.41 / 1.27	2.5 / 1.3 / 1.7
New York	20 072 [18 604, 21 466]	1–3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00	0 / 0 / 0
Mumbai	158 [148, 170]	4–6	1.02 / 1.04 / 1.04	1.11 / 1.08 / 1.04	1.02 / 1.04 / 1.04	1.5 / 1.2 / 0.39
London	927 [882, 991]	5–6	1.57 / 1.45 / 1.94	1.61 / 1.67 / 1.98	1.52 / 1.42 / 1.94	1.4 / 7.4 / 2.2
Paris	5 530 [5 023, 5 879]	3–5	1.03 / 1.03 / 1.05	1.03 / 1.03 / 1.05	1.03 / 1.02 / 1.05	0.047 / 0.19 / 0.23
Berlin	8 800 [8 570, 9 137]	2–4	1.02 / 1.02 / 1.03	1.02 / 1.02 / 1.03	1.02 / 1.02 / 1.02	0.054 / 0.054 / 0.46

Table S11: Imprecision, exact oracle (3 seeds; mean [min, max]). Left block: widest probability box ($\varepsilon = 0.41$) against $\varepsilon = 0$, change in deliveries and welfare (%). Right block: widest Γ-maximin width ($\delta = 0.80$) against $\delta = 0$. Monotone series: (seed, sweep) series in which deliveries never rise with the width. †: three exact widths per sweep.

City	Δdeliv. %	ΔW %	Δdeliv. %	ΔW %	monotone series
New Delhi	-53.5 [-55, -51]	-60.4 [-61, -59]	-62.2 [-75, -49]	-51.2 [-65, -39]	6/6
Cairo†	0	-2.88 [-3.0, -2.7]	-31.4 [-33, -31]	-22.8 [-24, -22]	6/6
Moscow†	-1.35 [-2.1, -0.95]	-10.5 [-11, -9.9]	-63.7 [-68, -62]	-55.1 [-61, -52]	6/6
Buenos Aires	-52.3 [-57, -49]	-52.9 [-57, -47]	-64.9 [-65, -64]	-54.6 [-56, -52]	3/6
New York†	-2.62 [-3.0, -2.1]	-10.9 [-11, -11]	-63.2 [-69, -57]	-54.4 [-61, -47]	6/6
Mumbai	-29.4 [-34, -23]	-30.3 [-32, -27]	-59.0 [-62, -56]	-44.7 [-48, -41]	6/6
London	-64.8 [-78, -56]	-74.8 [-82, -70]	-89.5 [-100, -81]	-86.5 [-100, -76]	6/6
Paris†	-10.2 [-12, -6.9]	-17.3 [-18, -16]	-71.3 [-74, -67]	-60.4 [-64, -54]	6/6
Berlin†	-8.68 [-9.2, -8.1]	-17.8 [-18, -17]	-71.2 [-72, -70]	-59.6 [-61, -58]	6/6

Table S12: DIP against the grid, per city and seed (0 / 1 / 2). Grid pts: exact grid points available (pb: the 63-point price-and-branch grid). Gain share: DIP gain divided by the best grid gain (n/a: no charge improves on the unpriced equilibrium). Calls: DIP's oracle calls as a percentage of the calls of the grid run, and of a 63-point grid at the same calls per point. Rounds: cutting-plane rounds (1: the tangent charge implemented the target without any cut). Exact: runs in which the target is an exact pure equilibrium under the returned charges. PoA of the grid's best point and of DIP's priced equilibrium.

City	grid pts	gain share	calls % (grid run)	calls % (63 pts)	rounds	exact	PoA grid	PoA DIP
New Delhi	63 / 63 / 63	1.23 / 1.35 / 1.14	5.6 / 4.9 / 5.5	5.6 / 4.9 / 5.5	2 / 1 / 2	3/3	1.02 / 1.05 / 1.02	1.00 / 1.00 / 1.00
Cairo	13 / 17 / 13	n/a / n/a / n/a	11 / 8.5 / 12	2.3 / 2.3 / 2.4	1 / 1 / 1	3/3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00
Moscow	12 / 10 / 63^{pb}	1.10 / 1.00 / 1.00	13 / 13 / 2.9	2.6 / 2.1 / 2.9	1 / 1 / 1	3/3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00
Buenos Aires	63 / 63 / 63	1.09 / 1.05 / 1.08	6.3 / 6.3 / 5.4	6.3 / 6.3 / 5.4	1 / 1 / 1	3/3	1.03 / 1.01 / 1.02	1.00 / 1.00 / 1.00
New York	7 / 7 / 63^{pb}	0.999 / 0.999 / 1.04	24 / 23 / 2.9	2.6 / 2.6 / 2.9	1 / 1 / 1	3/3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.000
Mumbai	15 / 14 / 14	3.25 / 1.20 / 1.28	24 / 31 / 26	5.6 / 6.9 / 5.8	1 / 2 / 2	3/3	1.02 / 1.00 / 1.00	0.997 / 0.997 / 0.995
London	63 / 63 / 63	1.04 / 1.31 / 1.06	7.2 / 9.4 / 8.7	7.2 / 9.4 / 8.7	3 / 3 / 3	3/3	1.01 / 1.08 / 1.02	1.00 / 1.00 / 0.995
Paris	14 / 13 / 13	1.02 / 1.06 / 1.05	22 / 23 / 21	4.9 / 4.8 / 4.4	3 / 1 / 2	3/3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00
Berlin	13 / 13 / 63^{pb}	1.03 / 1.03 / 1.13	17 / 15 / 4.1	3.5 / 3.2 / 4.1	1 / 1 / 1	3/3	1.00 / 1.00 / 1.00	1.00 / 1.00 / 1.00

Table S13: Branch-and-price with BRL pricing on the nine metro routing instances (seed 0, 22 customer zones, robust objective in driver-hours). Root LP: column-generation bound; P&B plan: price-and-branch plan; B&P optimum and remaining gap; nodes and time on one core; difference of the optimum from the best of three heuristics under the robust objective, and from the sample-average heuristic plan under the calibrated law. *Moscow congestion imputed.

City	root LP	P&B plan	B&P optimum	gap %	nodes	time s	vs best heur. %	vs SAA, true law %
New Delhi	63.03	63.18	63.18	0	65	1.63	0	0
Cairo	40.39	40.42	40.42	0	215	5.30	0	0
Moscow*	42.01	43.65	43.17	0	7487	1 063	-1.1	-0.57
Buenos Aires	47.29	48.37	48.13	0	1659	166	-2.4	7.0
New York	46.10	46.10	46.10	0	1	8.44	-10	4.8
Mumbai	53.82	54.08	53.91	0	177	3.59	-0.32	-0.31
London	51.04	52.03	51.38	0	59	21.2	0	0
Paris	37.73	38.32	38.25	0	4305	715	-0.10	-0.026
Berlin	51.85	52.67	52.06	0	3	5.63	-0.92	6.8




Figure S5: Leader's social-cost gain against routing-oracle calls: best grid point found so far (lines, exact grid rows in solve order) and DIP (one marker per run); 100% is the best grid point. Six cities have reduced exact grids; for Berlin seed 2, Moscow seed 2, New York seed 2 the grid is the price-and-branch grid.

Table S14: Tiny games with exhaustive enumeration, games with a positive leader gain. Mean share of the exact optimal gain, share of games in which the optimum is reached, games worse than the unpriced equilibrium, and mean oracle calls. For tariff rows, "worse than unpriced" compares with the potential-minimising unpriced equilibrium of the enumeration.

Method	city's information	mean share	optimum hit %	worse than unpriced	calls
ARP flat (plain)	(Q_z, n_z)	0.56	54	9	13
ARP flat (successive averages)	(Q_z, n_z)	0.64	57	8	51
ARP flat, lots known	(Q_z, n_z) + lots	0.63	54	15	12
tariff + carriers' best responses	own q, Q_z	0.80	71	3	5.4
tariff + potential solver	oracle	0.94	89	1	24
DIP	oracle	0.97	92	0	39
grid	oracle	0.83	67	0	1 492

Table S15: Theorems on the city games (three seeds). Mixed: seeds with a non-trivial mixed equilibrium. For those seeds (means): $\bar{\rho}$; the anchored three-term bound of Theorem S27(b); the N-uniform bound of Theorem S27(c); the largest empirical E-threshold over the tested profiles; and the ratio of the latter to the anchored bound. Γ-maximin (Lemma S25): drop in the share of demand served from the precise to the surcharge equilibrium at $\delta = 0.1$ and $\delta = 1$, and the maximality threshold of the surcharge equilibrium divided by δ at $\delta = 1$.

City	mixed	$\bar{\rho}$	anchored bd.	N-unif. bd.	max δ_E	ratio	drop δ=0.1	drop δ=1	M-thr./δ
New Delhi	3/3	1.0	1.61	10.7	0.64	0.41	0.0075	0.18	0.89
Cairo	0/3	–	–	–	–	–	0	0.47	0.99
Moscow	0/3	–	–	–	–	–	0.0021	0.73	0.97
Buenos Aires	1/3	0.96	1.76	8.80	0.83	0.47	0.082	0.31	0.93
New York	1/3	0.82	1.58	3.68	0.82	0.52	0.0090	0.71	0.98
Mumbai	2/3	0.74	1.23	10.6	0.53	0.43	0.049	0.34	0.92
London	3/3	0.85	1.56	13.6	0.82	0.53	0.033	0.17	0.76
Paris	3/3	0.88	1.39	8.43	0.69	0.49	0.036	0.74	0.96
Berlin	1/3	0.94	0.937	7.97	0.41	0.44	0.050	0.77	0.97

Table S16: Coverage of the exact-oracle game experiments per city and seed: exact grid points of the leader experiment (out of 63), price-of-anarchy experiment computed (yes/–), and exact widths of the imprecision experiment (out of 12); – means not computed with the exact oracle. The base equilibria are exact for all city-seeds.

City	E2 s0	E3 s0	E4 s0	E2 s1	E3 s1	E4 s1	E2 s2	E3 s2	E4 s2
New Delhi	63/63	yes	12/12	63/63	yes	12/12	63/63	yes	12/12
Cairo	13/63	yes	6/12	17/63	yes	6/12	13/63	yes	6/12
Moscow	12/63	yes	6/12	10/63	yes	6/12	11/63	yes	6/12
Buenos Aires	63/63	yes	12/12	63/63	yes	12/12	63/63	yes	12/12
New York	7/63	yes	6/12	7/63	yes	6/12	9/63	yes	6/12
Mumbai	15/63	yes	12/12	14/63	yes	12/12	14/63	yes	12/12
London	63/63	yes	12/12	63/63	yes	12/12	63/63	yes	12/12
Paris	14/63	yes	6/12	13/63	yes	6/12	13/63	yes	6/12
Berlin	13/63	yes	6/12	13/63	yes	6/12	13/63	yes	6/12

Table S17: Calibration per city (base values; ranges and sources in `data/parameters/SOURCES.md`). Parcels per capita are country-level; TomTom congestion is the 2025 city value (Moscow: 2020); drops per vehicle and van capacity are proxies outside the United States; fees and wages are graded C for most cities.

City	parcels/cap./day	cong. %	drops/veh.	shift h	van cap.	fee USD	wage USD/h	fuel USD/L	N	veh./1000
New Delhi	0.00964	60.2	40.0	10	40.0	0.690	1.07	1.15	5	400
Cairo	0.00561	40.5	40.0	8.4	40.0	1.18	0.610	0.440	4	220
Moscow	0.156	54.0	70.0	8.4	126	3.67	4.53	0.900	5	297
Buenos Aires	0.0150	48.1	70.0	8.4	126	0.980	1.28	1.45	4	331
New York	0.184	48.8	120	8.4	226	9.09	22.3	1.22	4	242
Mumbai	0.00964	63.2	40.0	10	40.0	0.690	1.07	1.15	5	330
London	0.166	51.6	72.0	7.3	126	4.21	18.8	2.34	5	288
Paris	0.0650	40.0	70.0	8.4	126	6.04	15.7	2.42	5	575
Berlin	0.143	44.6	126	8.4	206	4.40	18.8	2.39	6	330

Table S18: Sensitivity of unpriced welfare (% change, low to high value of each weak parameter, seed 0). The scenarios use the exact oracle; the reference base case is the price-and-branch equilibrium of the same seed, so differences of the order of the oracle effect (Section S5.1) are not meaningful.

City	fee	wage	drops	N	parcels
New Delhi	-58 to 317	-53 to 43	-90 to 294	-38 to 53	-9 to 0
Cairo	19 to 19	-5 to 2	-31 to 59	-25 to 24	-0 to 0
Moscow	-75 to 48	-26 to 6	-43 to 105	-62 to 21	-0 to 1
Buenos Aires	-59 to 452	-48 to -48	-87 to 474	-13 to 72	-1 to 6
New York	-41 to 55	-8 to 1	-57 to 9	-27 to 23	-0 to 0
Mumbai	-43 to 208	-36 to 26	-72 to 169	-19 to 29	-5 to -1
London	-95 to 363	-3 to 28	-95 to 795	-26 to 107	-2 to 4
Paris	-58 to 83	-4 to 10	-57 to 149	-16 to 27	-0 to 1
Berlin	-58 to 24	-5 to 3	-82 to 4	-3 to -3	-0 to 2

Table S19: City leader, broad objective with the background-delay proxy (3 seeds). Gains as in Table S9 (mean [min, max]); combined gain per seed; served share at the best point with both instruments. Shut-down optima (served share 0) are an artifact of the proxy.

City	cong. only	deliv. only	both	both by seed	served
New Delhi	77.1 [64, 85]	100	100	100 / 100 / 100	0
Cairo†	0	2.06 [0, 3.3]	4.59 [2.5, 7.5]	2.47 / 7.50 / 3.81	0.67 [0.65, 0.69]
Moscow†	3.67 [3.1, 4.1]	0.0230 [0, 0.069]	4.87 [3.6, 6.3]	3.59 / 4.72 / 6.31	0.92 [0.91, 0.93]
Buenos Aires	2.41 [1.4, 4.0]	0	3.71 [1.1, 5.4]	4.62 / 5.42 / 1.09	0.35 [0.29, 0.46]
New York†	0.00636 [0, 0.019]	0.320 [0.18, 0.52]	0.344 [0.26, 0.52]	0.259 / 0.518 / 0.256	1.00 [0.99, 1.0]
Mumbai†	66.7 [65, 71]	100	100	100 / 100 / 100	0
London	90.9 [78, 100]	100	100	100 / 100 / 100	0
Paris†	37.7 [32, 44]	74.2 [72, 76]	99.1 [98, 100]	99.3 / 100 / 98.1	0.042 [0, 0.085]
Berlin†	4.07 [2.6, 5.9]	2.39 [1.5, 3.2]	3.10 [2.1, 4.5]	2.13 / 2.66 / 4.50	0.74 [0.65, 0.88]

Table S20: ZMC on synthetic D1 games with non-separable plan sets (five seeds per row). Entries: mean oracle calls / mean potential gap to the convexified optimum; last column: instances in which ZMC has the smallest gap among FW8, CG8 and multi-start BR.

N	Z	BR	BR multi-start	FW8	CG8	ZMC	CG full	ZMC best
4	30	12 / 5.33	45 / 3.22	38 / 3.64	39 / 3.20	39 / 2.77	58 / 2.69	4/5
5	60	21 / 15.8	56 / 11.8	50 / 7.78	51 / 7.80	49 / 6.90	159 / 6.13	4/5
6	120	28 / 50.2	70 / 38.4	61 / 21.4	63 / 27.5	60 / 17.1	365 / 14.2	4/5
6	250	26 / 91.5	71 / 80.2	64 / 47.9	67 / 55.4	60 / 33.4	797 / 27.8	5/5

Table S21: Tiny games solved by exhaustive enumeration (price-and-branch oracle): share of instances in which the method's profile is a Nash equilibrium of the enumerated game and reaches the global potential minimum, largest independent-oracle regret and mean excess of the potential over the global minimum.

Method	games	calls	is NE	max regret	global min.	excess
ZMC (6 rounds)	40	17.5	1.00	0	1.00	0
BR	40	4.85	1.00	0	0.875	0.93
FW (6 rounds)	40	16.7	1.00	0	0.950	0.067
CG	40	11.1	1.00	0	0.975	0.020

Table S22: Checks of ZMC on the city games (seed 0, price-and-branch oracle): largest residual of the clearing identity (Theorem S54(a)); share of rounds in which its bound holds; inverted brackets; active coordinates; largest relative bracket width after eight rounds; in bisection mode, share of coordinates within the bound of Theorem S55 at $T = 8$, largest normalised error and the bound 2^{-4}.

City	identity resid.	bound holds	inverted	active	width_8	within_8	err_8	bound_8
New Delhi	$2.8 \cdot 10^{-14}$	1.00	0	128	0.0435	1.00	0.00385	0.0625
Cairo	$1.6 \cdot 10^{-13}$	0.875	0	64	0.0435	1.00	0.00385	0.0625
Moscow	$8.1 \cdot 10^{-13}$	1.00	0	142	0.0435	1.00	0.00385	0.0625
Buenos Aires	$5.7 \cdot 10^{-14}$	1.00	0	124	0.0435	1.00	0.00385	0.0625
New York	$3.4 \cdot 10^{-12}$	1.00	0	128	0.0435	1.00	0.00385	0.0625
Mumbai	$4.3 \cdot 10^{-14}$	1.00	0	110	0.0435	1.00	0.00385	0.0625
London	$1.1 \cdot 10^{-13}$	1.00	0	178	0.0435	1.00	0.00385	0.0625
Paris	$1.5 \cdot 10^{-12}$	1.00	0	136	0.0415	1.00	0.00385	0.0625
Berlin	$1.8 \cdot 10^{-12}$	1.00	0	164	0.0435	1.00	0.00385	0.0625




Figure S6: Credal absorption on the city equilibria (three seeds). Left: N-uniform bound, anchored bound, smallest anchored radius and largest empirical E-threshold per anchor (Cairo and Moscow: pure anchors only, plotted at 10^{-3} for zero). Right: ratio of the largest empirical E-threshold to the anchored bound against $d = 2Z$; values below one mean the bound holds.




Figure S7: City games of Cairo, Mumbai and Paris (three seeds) with a vehicle-count externality of increasing scale. Left: share of best-response runs that converge to a pure equilibrium. Right: largest regret of the D1 equilibrium when the game is of type D2 (USD).

Table S23: BRL pricing on random instances (nine sectors, $L = 33$ levels, capacity 6, two instances per size). Time in seconds (labels kept); TO: 60 s limit. The risk-neutral labelling solves a different (expected-value) problem and is shown for reference; SAA scenarios optimise the sample-average objective.

n	risk-neutral	BRL	sector (K)	quantile (L)	SAA scenarios
10	0.0050 (19)	**0.0040 (16)**	TO	2.2 (4946)	5.4 (7904)
10	0.017 (36)	**0.0080 (16)**	TO	2.1 (4391)	4.7 (7715)
15	0.17 (264)	**0.019 (48)**	TO	TO	TO
15	0.029 (69)	**0.010 (24)**	TO	TO	TO
20	0.39 (714)	**0.049 (97)**	TO	TO	TO
20	1.4 (1730)	**0.047 (89)**	TO	TO	TO
25	0.46 (650)	**0.059 (94)**	TO	TO	TO
25	0.49 (693)	**0.065 (109)**	TO	TO	TO

Table S24: Median number of labels kept by three dominance rules priced once with the duals of the root column generation (metro instances, and Solomon and Augerat instances with a soft duration limit; CG: column generation converged within 150 s). Ratios are medians of per-instance ratios; capped runs give lower bounds.

family	n	instances	labels: window (BRL)	no window	quantile-vector	no window / window	quantile / window
metro	22	9	2 070	6 453	12 692	2.6	4.2
metro	30	9	4 187	9 744	26 166	2.9	6.3
metro	40	9	7 918	21 585	49 295	2.0	4.6
metro	60	8	15 801	38 138	83 375	2.2	3.8
metro	80	8	24 475	52 598	87 976	1.8	2.8
public	25	3	4 101	46 140	75 556	9.3	20
public	31	1	3 744	51 972	87 375	14	23
public	44	1	6 612	55 074	107 450	8.3	16
public	50	3	24 283	69 094	89 324	2.8	3.7
public	59	1	13 986	108 840	140 020	7.8	10
public	79	1	7 845	26 199	114 320	3.3	15
public	100	3	39 104	53 370	101 530	1.6	2.7

Table S25: Out-of-sample comparison on the public instances (17 cells per row), polished plans, relative to the sample-average plan with the same number of training days. Held-out: 2000 days from the training law; shifted law: excess congestion $\times 1.5$, volatility $\times 1.25$, no re-solving.

N days	plan	held-out mean %	95th pct %	$\text{CVaR}_{0.9}$ %	shifted law: mean %	shifted law: 95th pct %
10	BRL	5.0	2.8	2.0	3.3	-2.3
10	SAA	0	0	0	0	0
10	Wasserstein	0.036	0.012	0.076	0.065	0.17
10	mean-CVaR	0.77	-0.45	-0.55	0.21	-1.0
10	deterministic	0.13	2.8	2.7	0.97	2.2
30	BRL	0.42	1.6	1.5	0.62	0.21
30	SAA	0	0	0	0	0
30	Wasserstein	-0.087	-0.041	0.051	-0.015	0.23
30	mean-CVaR	0.11	0.040	0.14	0.14	0.29
30	deterministic	0.056	2.0	1.9	0.56	0.89
60	BRL	0.30	1.2	0.97	0.40	-0.073
60	SAA	0	0	0	0	0
60	Wasserstein	-0.033	0.27	0.25	0.091	0.32
60	mean-CVaR	0.16	-0.062	-0.17	0.021	-0.33
60	deterministic	0.35	3.0	2.8	1.1	2.0

Table S26: Leader methods on four cities, seeds 0–2 (exact oracle). ϱ: contraction constant of Theorem S18. ARP: flat rule, status and rounds (a cycle is a repeated charge and market state). Gains are percentages of the best exact grid point's gain; tariff+BR: marginal tariff with the carriers' own best responses; tariff+SO: tariff with the potential solver. Calls: oracle calls of the carriers during the method. Mumbai's grid is reduced (gains relative to a lower bound).

City	seed	ϱ	ARP status / rounds	ARP %	calls	tariff+BR %	calls	tariff+SO %	calls	DIP %	calls	grid calls
New Delhi	0	0.40	cycle / 5	90.8	53	98.7	11	92.5	51	123	134	2404
New Delhi	1	0.19	fixed point / 3	126	27	135	10	127	51	135	119	2418
New Delhi	2	0.40	cycle / 6	105	60	103	9	114	48	114	132	2419
Buenos Aires	0	0.029	fixed point / 3	108	24	108	9	108	40	109	121	1921
Buenos Aires	1	0.016	fixed point / 3	105	22	105	10	105	40	105	122	1950
Buenos Aires	2	0.11	fixed point / 3	108	23	108	8	108	39	108	105	1943
Mumbai	0	0.20	fixed point / 7	293	65	293	9	263	51	325	139	586
Mumbai	1	0.094	fixed point / 3	111	30	113	10	108	50	120	169	544
Mumbai	2	0.20	cycle / 5	128	47	111	10	118	49	128	141	537
London	0	0.10	cycle / 5	102	51	104	14	100	49	104	170	2376
London	1	0.10	cycle / 5	120	47	122	9	131	50	131	221	2356
London	2	0.10	cycle / 7	103	67	105	11	101	50	106	206	2378

S7. Proofs for the equilibrium theory

The proofs in this appendix are taken from the theory notes in the repository (file `theory/main.tex`), with the notation of Section S1.

S7.1. Tightness and efficiency results

These results complete the equilibrium theory of Section S3; their proofs follow with the others in this appendix.

Tightness.

Proposition S34 (The support-selection bound is attained)**.** *Let $N = 2$, one zone, D2, $M = \gamma c = 1$ and weight $w > 0$. Carrier 1 has plans $A_1 : (n, \tau, \pi) = (1, 4, 6)$ and $B_1 : (3, 1, 0)$; carrier 2 has $X : (1, 1, 0)$ and $Y : (3, 4, 6)$; payoffs are $\pi - \tau \cdot n_{\text{rival}}$. Then the game has no pure Nash equilibrium, $\delta_0^E = \delta_0^M = 1/w$, and the bound $(N-1)\bar{\rho}$ of Theorem S27(b) equals $1/w$.*

In each of the four pure profiles exactly one carrier has a strict incentive to deviate at the realised count, and it becomes indifferent only at a belief that differs from the realised count by one vehicle.

Proposition S35 (Linear dependence on the number of zones is necessary)**.** *For every $Z \geq 1$ and $N \geq 2$ there is a D2 game with Z zones in which every pure profile has $\max_i \varepsilon_i \geq 3\lceil Z/2 \rceil$, while every profile of Theorem S27(a) satisfies $\hat{\delta}(p) L_i(p) \leq 3Z$ $(w = 1)$.*

Efficiency.

Lemma S36 (Imprecise equilibria are approximate equilibria)**.** *If $\sigma \in \mathrm{ICNE}_C(\delta)$ and $\mu \in \Xi$, then σ is an ε-Nash equilibrium of the precise game with $\varepsilon_i \leq \delta L_i(\sigma) + \gamma \sum_z \varpi_z T_{iz}(\sigma)$ for $C \in \{E, M\}$ and $\varepsilon_i \leq 2\delta H_i + 2\gamma \Upsilon_i$ for $C = ID$, with $H_i = \max_p \sum_k w_k M_k \varphi_{ik}(p)$ and $\Upsilon_i = \max_p \sum_z \varpi_z t_{iz}(p)$.*

Proposition S37 (Monotone efficiency curves)**.** *For $C \in \{E, M, ID\}$ and $W > 0$, $\mathrm{PoA}_C(\delta) = W^* / \min_{\mathrm{ICNE}_C(\delta)} W$ is nondecreasing and $\mathrm{PoS}_C(\delta) = W^* / \max_{\mathrm{ICNE}_C(\delta)} W$ nonincreasing in δ, and both are step functions with at most $\prod_i |\mathcal{P}_i|$ steps. For $C = \Gamma$ neither monotonicity holds in general.*

Proposition S38 (Price of stability via the potential, D1)**.** *Under D1 with zero charges and $\mu = \hat{\mu}$, let σ^* maximise W. Every profile σ with $\Psi_{\hat{\mu}}(\sigma) \geq \max \Psi_{\hat{\mu}} - \eta$ satisfies $W(\sigma) \geq W^* - \sum_z \left[\frac{b_z}{2} \sum_i q_{iz}^{*2} + \frac{\gamma c_z}{2} (1 - \frac{1}{N}) T_z(\sigma)^2 \right] - \eta$. In particular a potential maximiser, which lies in every $\mathrm{ICNE}_E(\delta)$, bounds $\mathrm{PoS}_C(\delta)$ for all δ and $C \in \{E, M, ID\}$.*

Proposition S43 (below) gives a conditional smoothness-based price-of-anarchy bound that transfers through Lemma S36.

S7.2. Classical tools

Theorem S39 (Shapley–Folkman, constructive vertex form; 55)**.** *Let $S_1, \ldots, S_N \subset \mathbb{R}^d$ be finite and nonempty, $L = \operatorname{span} \bigcup_i (S_i - S_i)$ and $d_L = \dim L$. Index the elements of S_i by $\mathcal{P}_i$ through an onto map a_i. For $\bar{x} \in \sum_i \operatorname{conv} S_i$, the polytope $\Lambda(\bar{x}) = \{\lambda \in \prod_i \Delta(\mathcal{P}_i) : \sum_i \sum_p \lambda_i(p) a_i(p) = \bar{x}\}$ is nonempty, and at every vertex λ the set $J(\lambda) = \{i : |\operatorname{supp} \lambda_i| \geq 2\}$ has $|J(\lambda)| \leq \min\{N, d_L\}$.*

Proof. Nonemptiness: $\bar{x} = \sum_i y_i$ with $y_i \in \operatorname{conv} S_i$, and each y_i is a convex combination of points $a_i(p)$. $\Lambda(\bar{x})$ is a bounded polyhedron in standard form $\{\lambda \geq 0 : H\lambda = h\}$ with columns $c_{i,p} = (a_i(p), e_i) \in \mathbb{R}^d \times \mathbb{R}^N$ and $h = (\bar{x}, \mathbf{1})$. A feasible λ is a vertex if and only if the columns $\{c_{i,p} : \lambda_i(p) > 0\}$ are linearly independent. Fix a vertex and pick $p_i^0 \in \operatorname{supp} \lambda_i$ for each i. The family $\{c_{i,p_i^0}\}_i \cup \{c_{i,p} - c_{i,p_i^0} : p \in \operatorname{supp} \lambda_i \setminus \{p_i^0\}\}$ is obtained from the positive columns by an invertible (unitriangular) transformation, so it is linearly independent. Its second part consists of the vectors $(a_i(p) - a_i(p_i^0), 0) \in L \times \{0\}$, which number $\sum_i (|\operatorname{supp} \lambda_i| - 1) \geq |J(\lambda)|$. Linearly independent vectors in $L \times \{0\}$ number at most d_L. Hence $|J| \leq d_L$, and trivially $|J| \leq N$. $\square$

Proof of Lemma S23. (a) $u_i(p; m, \xi)$ is affine in (m, ξ) with fixed coefficients, and P has finite first moments; linearity of the integral gives the identity. (b) Into: by definition. Onto: for $k = (m, \xi) \in K_i^\delta(\sigma)$, the Dirac law at (m, ξ) is supported in $\mathbb{R}^d_{\geq 0} \times \mathbb{R}^Z_{\geq 0}$ because $m \geq 0$ and $\xi \in \Xi \subset \mathbb{R}^Z_{\geq 0}$; its means are (m, ξ), so it belongs to $\mathcal{M}_i^\delta(\sigma)$. (c) Each criterion depends on P only through the numbers $\mathbb{E}_P[u(p)]$ and $\mathbb{E}_P[u(p') - u(p)]$, which are affine functions of the mean by (a). By (b), ranging over P is the same as ranging over $k \in K_i^\delta(\sigma)$, a product of compact intervals; continuous functions attain their extrema on it. □

Proof of Lemma S24. For a finite $\mathcal{P}$ and the compact convex box K with continuous affine payoffs: $E \subseteq M$. If p is optimal at k_0, then for every p', $\bar{u}(p'; k_0) - \bar{u}(p; k_0) \leq 0$, so $\min_k[\bar{u}(p') - \bar{u}(p)] \leq 0$. $M \subseteq ID$. If p is maximal, then for each p' there is k with $\bar{u}(p; k) \geq \bar{u}(p'; k)$; hence $\max \bar{u}(p; \cdot) \geq \bar{u}(p; k) \geq \bar{u}(p'; k) \geq \min \bar{u}(p'; \cdot)$. $\Gamma \subseteq M$. If $\min_k[\bar{u}(p'; k) - \bar{u}(p; k)] > 0$, then $\min \bar{u}(p') > \min \bar{u}(p)$, so p is not Γ-maximin. These inclusions hold carrier by carrier and give the inclusions between equilibrium sets. *Monotonicity.* $K^\delta(c) \subseteq K^{\delta'}(c)$ for every c, and the realised centre $A_{-i}(\sigma)$ does not depend on δ. E: a witness belief stays admissible. M: a dominance strict on the larger set is strict on the smaller one, so non-domination on the smaller set implies it on the larger one. ID: $\overline{\mathbb{E}}$ increases and every $\underline{\mathbb{E}}$ decreases as the set grows. □

Example S40 (Γ-equilibria are not nested)**.** Take one carrier with a fixed external aggregate $c = (1, 2)$, $d = 2$, $M = I$, $w = (2, 1)$, no traffic term, and two plans with $\pi = 0$, $\varphi(A) = (1, 0)$, $\varphi(B) = (0, 1)$. The lower values are $-(1 + 2\delta)$ for A and $-(2 + \delta)$ for B, so $\mathrm{ICNE}_\Gamma(0) = \{A\}$ and $\mathrm{ICNE}_\Gamma(2) = \{B\}$.

Proof of Lemma S25. A minimum of affine functions of the mixture x is concave, and

$$\min_k \textstyle\sum_p x(p)\bar{u}_i(p; k) \geq \sum_p x(p) \min_k \bar{u}_i(p; k)$$

because the left side picks one k for all p. With $\varphi, t \geq 0$ and $M \geq 0$, $\bar{u}_i(p; m, \xi)$ is coordinatewise nonincreasing in (m, ξ). The box $K^\delta(c)$ has the coordinatewise maximum $c + \delta w$, which lies in it (clipping only removes points below 0), and Ξ has the maximum $\overline{\mu}$. So $k^+ = (c + \delta w, \overline{\mu})$ minimises every $\bar{u}_i(p; \cdot)$ and the lower expectation is linear in the mixture. Substituting k^+ with $c = A_{-i}(\sigma)$ gives

$$\bar{u}_i(p; k^+) = \pi_i^\Gamma(p) - \langle \varphi_i(p), MA_{-i}(\sigma) \rangle,$$

the payoff of G_δ^Γ; Γ-maximin is then best response in G_δ^Γ. □

Proposition S41 (Relaxed equilibria are mixed equilibria)**.** *Under Assumption A2, x is a mixed Nash equilibrium of the finite game with payoffs $\bar{u}_i(\sigma_i; A_{-i}(\sigma), \hat{\xi})$ if and only if* $\operatorname{supp} x_i \subseteq \arg\max_p \bar{u}_i(p; \bar{A}_{-i}(x), \hat{\xi})$ *for all i. Such x exist.*

Proof. Under the product law x_{-i}, the expected payoff of a pure plan p is $\mathbb{E}[\bar{u}_i(p; A_{-i}, \hat{\xi})] = \bar{u}_i(p; \mathbb{E}A_{-i}, \hat{\xi})$ because $\bar{u}_i$ is affine in A_{-i}, and $\mathbb{E}A_{-i} = \bar{A}_{-i}(x)$ by linearity. A mixed profile is a Nash equilibrium if and only if each x_i is supported on pure best responses to x_{-i}. Existence follows from Nash [37]. □

Proof of Proposition S26. For a unilateral change $p \to p'$ of carrier i, put $a = a_i(p)$, $a' = a_i(p')$ and $R = A_{-i}$. Then $\Psi(p') - \Psi(p) = \pi_i(p') - \pi_i(p) + \frac{1}{2}\|a'\|_M^2 - \frac{1}{2}\|a\|_M^2 - \frac{1}{2}\|R + a'\|_M^2 + \frac{1}{2}\|R + a\|_M^2 - \gamma\langle t_i(p') - t_i(p), \hat{\xi}\rangle$, which equals $\pi_i(p') - \pi_i(p) - \langle a' - a, MR\rangle - \gamma\langle t_i(p') - t_i(p), \hat{\xi}\rangle = u_i(p') - u_i(p)$, using $\varphi_i = a_i$. A maximiser σ is a pure equilibrium [36]. The point $(A_{-i}(\sigma), \hat{\xi})$ lies in $K_i^\delta(\sigma)$ for every δ, so $\sigma \in \mathrm{ICNE}_E(\delta)$. For Γ, Lemma S25 shows that $\mathrm{ICNE}_\Gamma(\delta)$ is the equilibrium set of G_δ^Γ, which is again D1 with modified private profits and therefore has a potential and a pure equilibrium. □

Proof of Theorem S27. (a) Fix i and $\delta \geq \hat{\delta}(p)$. The anchor $\bar{A}^*_{-i}$ is a sum of convex combinations of nonnegative vectors, hence nonnegative. By definition of $\hat{\delta}(p)$, $|\bar{A}^*_{-i,k} - A_{-i,k}(p)| \leq \hat{\delta}(p) w_k \leq \delta w_k$ for all k, so $(\bar{A}^*_{-i}, \hat{\xi}) \in K_i^\delta(p)$. By Lemma S23(b) the Dirac law at this point belongs to $\mathcal{M}_i^\delta(p)$; under it the expected payoff is $\bar{u}_i(\cdot; \bar{A}^*_{-i}, \hat{\xi})$, which $p_i \in B_i$ maximises. Hence p_i is E-admissible for every i, and the inclusions follow from Lemma S24.

(b), first term. For any $p \in \prod_j B_j$, $A_{-i}(p) - \bar{A}^*_{-i} = \sum_{j \neq i} (a_j(p_j) - \bar{a}_j)$ and $|a_{jk}(p_j) - \bar{a}_{jk}| \leq r_{jk}$.

(b), second term. $\bar{A}^* \in \sum_j \operatorname{conv} a_j(B_j)$ because $\operatorname{supp} x_j^* \subseteq B_j$. Apply Theorem S39 with $\mathcal{P}_j$ replaced by B_j: a vertex λ of $\Lambda(\bar{A}^*)$ exists and can be found by linear programming. Let $J = J(\lambda)$, $|J| \leq k^\sharp$, and $y_j = \sum_p \lambda_j(p) a_j(p)$. For $j \notin J$, λ_j is a point mass at some $p_j \in B_j$ and $a_j(p_j) = y_j$; for $j \in J$ choose any $p_j \in \operatorname{supp} \lambda_j \subseteq B_j$. Since $\sum_j y_j = \bar{A}^* = \sum_j \bar{a}_j$,

$$A_{-i}(p) - \bar{A}^*_{-i} = \textstyle\sum_{j \in J} (a_j(p_j) - y_j) - (a_i(p_i) - \bar{a}_i). \tag{S6}$$

For $j \in J$ both $a_{jk}(p_j)$ and y_{jk} lie in $[\min_{B_j} a_{jk}, \max_{B_j} a_{jk}]$, so the sum is at most $k^\sharp \bar{\sigma} w_k$ in coordinate k, and the last term is at most $\bar{\rho} w_k$.

(b), third term. If $J = \emptyset$ or $\bar{\sigma} = 0$, the sum vanishes. Otherwise draw $p_j \sim \lambda_j$ independently for $j \in J$. For each k the variables $(a_{jk}(p_j) - y_{jk})/w_k$, $j \in J$, are independent, have mean zero and range in an interval of length at most $\bar{\sigma}$. By Hoeffding's inequality and a union bound over k, $\Pr(\|E\|_{w,\infty} \geq t) \leq 2d \exp(-2t^2/(k^\sharp \bar{\sigma}^2))$, where E is the sum in (S6). This is below one for $t > t_0 = \bar{\sigma}\sqrt{k^\sharp \ln(2d)/2}$; outcomes are finitely many, so some outcome has $\|E\|_{w,\infty} \leq t_0$, and $\hat{\delta}(p) \leq t_0 + \bar{\rho}$ for it.

(c) Relaxed equilibria exist (Proposition S41). Use $r_{jk} \leq s_{jk}$, $\bar{\rho} \leq \bar{\sigma} \leq \sigma_{\max}$, $k^\sharp \leq d$ and $\sum_{j \neq i} r_{jk}/w_k \leq (N-1)\sigma_{\max}$; the order of thresholds follows from Lemma S24. *(d)* This was shown in (b); linear programming finds a vertex in time polynomial in $\sum_j |B_j|$ and d. □

Proof of Corollary S28. Let $m^* = \bar{A}^*_{-i}$ and $m = A_{-i}(p)$. For any p', $\bar{u}_i(p'; m, \mu) - \bar{u}_i(p_i; m, \mu) = [\bar{u}_i(p'; m^*, \hat{\xi}) - \bar{u}_i(p_i; m^*, \hat{\xi})] - \langle \varphi_i(p') - \varphi_i(p_i), M(m - m^*) \rangle - \gamma \langle t_i(p') - t_i(p_i), \mu - \hat{\xi} \rangle$. The bracket is nonpositive because $p_i \in B_i$; each term $|M_k(\varphi_{ik}(p') - \varphi_{ik}(p_i))(m_k - m_k^*)|$ is at most $w_k M_k |\Delta\varphi_{ik}| \hat{\delta}(p)$, and $|\mu_z - \hat{\xi}_z| \leq \varpi_z$. □

Proof of Proposition S34. For carrier 1, $\bar{u}(A_1; n) - \bar{u}(B_1; n) = 6 - 3n$, so A_1 is optimal iff $n \leq 2$; for carrier 2, $\bar{u}(Y; n) - \bar{u}(X; n) = 6 - 3n$, so Y is optimal iff $n \leq 2$. With two plans and a convex belief set, E-admissibility and maximality coincide: some n in the box makes the plan weakly better. In (A_1, X) carrier 2 sees $n = 1$ and needs a belief ≥ 2; in (A_1, Y) carrier 1 sees $n = 3$ and needs a belief ≤ 2; in (B_1, Y) carrier 2 sees $n = 3$ and needs ≤ 2; in (B_1, X) carrier 1 sees $n = 1$ and needs ≥ 2. Each requirement holds iff $\delta w \geq 1$, and the other carrier best-responds at the realised count. Hence $\mathrm{ICNE}_E(\delta) = \mathrm{ICNE}_M(\delta) = \emptyset$ for $\delta < 1/w$ and both equal the full profile set at $\delta = 1/w$; there is no pure Nash equilibrium. The unique mixed equilibrium makes each carrier indifferent, so the rival's mean count is 2, both mix $\frac{1}{2}$–$\frac{1}{2}$, $r_j = 1$, $\bar{\rho} = 1/w$ and $(N-1)\bar{\rho} = 1/w$. □

Proof of Proposition S35. Take two carriers with plan sets $\{A_1, B_1\}^Z$ and $\{X, Y\}^Z$, the one-zone data of Proposition S34 in each zone and no cross-zone interaction, and $N - 2$ carriers with a single null plan. Payoffs are sums over zones and plan sets are products, so a carrier's regret is the sum of its zone regrets. In each zone and each of the four zone profiles, exactly one of the two carriers has regret 3 and the other 0 (e.g. in (A_1, X) carrier 2 gains $2 - (-1) = 3$ by switching to Y). So $\varepsilon_1 + \varepsilon_2 = 3Z$ with both multiples of 3, and $\max \geq 3\lceil Z/2 \rceil$. For the upper bound, the zone-wise mixed equilibrium has $\bar{\rho} = 1$, every $p \in \prod B_j$ has $\hat{\delta}(p) \leq 1$, and $L_i(p) \leq \sum_z 1 \cdot 1 \cdot |\Delta\tau_z| = 3Z$. □

Proof of Theorem S31. Membership. The witness is a profile σ. With K the clipped box and Ξ a point: E-admissibility of σ_i is feasibility of the linear system $\{m \in K : \bar{u}_i(\sigma_i; m) \geq \bar{u}_i(p'; m)\ \forall p'\}$;

maximality requires $\max_{m\in K}[\bar{u}_i(\sigma_i;m)-\bar{u}_i(p';m)]\geq 0$ for each p', a linear function maximised over a box coordinatewise; ID and Γ require maxima and minima of $\bar{u}_i(p;\cdot)$ over the box, also coordinatewise.

Construction. Let $V(D)=[n]$ and $\Delta^-=\max(1,\text{max in-degree})$. Create one zone per arc $e=(u,v)$, using only its vehicle coordinate, with $M_e=\gamma c_e=1$, $w_e=1$ and no deliveries. Set $\eta=1/(2\Delta^-+2)$. Carrier i has the plan *out* with all data 0, and the plan *in* with $\pi_i=1$ and, for each out-arc $e=(i,j)$, vehicles $n_{ie}=1$ and exposure $\tau_{ie}=1$; for each in-arc $e=(h,i)$, vehicles $n_{ie}=2$ and exposure $\tau_{ie}=\eta$; zero elsewhere. This is D2, since $\varphi=\tau\neq a=n$.

Payoff gap. The gain of *in* at belief m is $D_i(m)=1-\sum_{e=(i,j)}m_e-\eta\sum_{e=(h,i)}m_e$. At the realised aggregate of σ^S, $m_e=2\cdot\mathbf{1}[j\in S]$ for $e=(i,j)$ and $m_e=\mathbf{1}[h\in S]$ for $e=(h,i)$, so $D_i^{\text{real}}=1-2o_i-\eta h_i$ with $o_i=|N^+(i)\cap S|$ and $h_i=|N^-(i)\cap S|\leq\Delta^-$. If $o_i=0$, $D_i^{\text{real}}\geq 1-\Delta^-/(2\Delta^-+2)>\frac{1}{2}$; if $o_i\geq 1$, $D_i^{\text{real}}\leq -1$.

Robustness of the sign. Over K^δ, $|D_i(m)-D_i^{\text{real}}|\leq\delta(\deg^+(i)+\eta\deg^-(i))<\delta(\Delta^+\!+\frac{1}{2})$, which is below $\frac{1}{2}$ for $\delta<1/(2\Delta^+\!+1)$; clipping only shrinks the box. So D_i has the sign of D_i^{real} on the whole box.

All criteria select the realised strict best response. If $D_i>0$ on the box, *in* is the unique optimum at every belief, hence the only E-admissible plan; *out* is strictly dominated everywhere, so *in* is the only maximal plan; and $\min_K\bar{u}(\text{in})>0=\max_K\bar{u}(\text{out})$, so *out* is interval-dominated and *in* is the unique Γ-maximin plan. The case $D_i<0$ is symmetric. So σ is admissible under every criterion iff each carrier plays *in* exactly when $o_i=0$.

Equilibria are kernels. With $S=\{i:\sigma_i=\text{in}\}$, the condition reads $i\in S\Rightarrow N^+(i)\cap S=\emptyset$ (independence) and $i\notin S\Rightarrow N^+(i)\cap S\neq\emptyset$ (absorption). NP-hardness follows from Chvátal [40]; all numbers are polynomially bounded, so hardness is strong; with $\Delta^+\leq 2$ [41] the threshold is $1/5$. □

Proof of Corollary S32. On the instances of Theorem S31, $\delta_0^E=0$ if D has a kernel; otherwise $\text{ICNE}_E(\delta)=\emptyset$ for all $\delta<1/(2\Delta^+\!+1)$, so $\delta_0^E\geq 1/(2\Delta^+\!+1)$. A multiplicative approximation outputs 0 in the first case and a positive number in the second; an additive one separates the cases when its error is below $1/(2\Delta^+\!+1)$. □

Proposition S42 (The relaxation step is PPAD-hard under D2)**.** *Every bimatrix game $(R,C)\in\mathbb{Q}^{m\times n}\times\mathbb{Q}^{m\times n}$ is, in polynomial time, the precise game of a D2 instance with $N=2$ and $d=m+n$. Hence computing a mixed equilibrium under D2 is PPAD-hard.*

Proof. Let $M=I_{m+n}$, no traffic term, $\bar{R}=\max R$ and $\bar{C}=\max C$. Carrier 1 has plans $p\in[m]$ with $a_1(p)=e_p$, $\varphi_1(p)=\sum_q(\bar{R}-R_{pq})e_{m+q}\geq 0$ and $\pi_1(p)=\bar{R}$; carrier 2 has plans $q\in[n]$ with $a_2(q)=e_{m+q}$, $\varphi_2(q)=\sum_p(\bar{C}-C_{pq})e_p\geq 0$ and $\pi_2(q)=\bar{C}$. Then $u_1(p;a_2(q))=R_{pq}$ and $u_2=C_{pq}$. By Proposition S41, relaxed equilibria are exactly the Nash equilibria of (R,C), and computing these is PPAD-complete [44, 45]. □

Proof of Lemma S36. Write $k^t=(A_{-i}(\sigma),\mu)\in K_i^\delta(\sigma)$ and $\Delta(p',k)=\bar{u}_i(p';k)-\bar{u}_i(\sigma_i;k)$, affine in k. For any k in the (unclipped) box, $|\Delta(p',k)-\Delta(p',k^t)|\leq\delta\sum_k w_kM_k|\Delta\varphi_{ik}|+\gamma\sum_z\varpi_z|\Delta t_{iz}|$. *E:* take the witness k, at which $\Delta(p',k)\leq 0$. *M:* for each p', maximality and compactness give $k_{p'}$ with $\Delta(p',k_{p'})\leq 0$. *ID:* $\bar{u}_i(p';k^t)-\bar{u}_i(\sigma_i;k^t)\leq[\bar{u}_i(p';k^t)-\min_K\bar{u}_i(p')]+[\max_K\bar{u}_i(\sigma_i)-\bar{u}_i(\sigma_i;k^t)]$, since $\max_K\bar{u}_i(\sigma_i)\geq\min_K\bar{u}_i(p')$, and each bracket is at most $\delta H_i+\gamma\Upsilon_i$. □

Proof of Proposition S37. The minimum over a growing finite set does not increase and the maximum does not decrease (Lemma S24). Each profile enters $\text{ICNE}_C(\delta)$ at a single threshold and never leaves, so the sets change at most at $\prod_i|\mathcal{P}_i|$ values. For Γ see Example S40. □

Proposition S43 (Price of anarchy under smoothness; conditional)**.** *Assume $W(\sigma)\geq\sum_iU_i(\sigma)$ for all σ, $W^*>0$, and that the precise game is (λ,μ)-smooth: $\sum_iU_i(\sigma_i^*,\sigma_{-i})\geq\lambda W(\sigma^*)-\mu W(\sigma)$ for all σ and an optimal σ^*. For $C\in\{E,M,ID\}$ let $\mathcal{E}(\delta)=\max_{\sigma\in\text{ICNE}_C(\delta)}\sum_i\varepsilon_i(\sigma)$ from Lemma S36. If $\lambda W^*>\mathcal{E}(\delta)$, then $\text{PoA}_C(\delta)\leq(1+\mu)/(\lambda-\mathcal{E}(\delta)/W^*)$.*

Proof. For $\sigma \in \mathrm{ICNE}_C(\delta)$: $W(\sigma) \geq \sum_i U_i(\sigma) \geq \sum_i [U_i(\sigma_i^*, \sigma_{-i}) - \varepsilon_i] \geq \lambda W^* - \mu W(\sigma) - \mathcal{E}(\delta)$. Rearrange. □

Proof of Proposition S38. With π_i as in Section S1 and $x = \tau$, direct expansion gives $W - \Psi_{\hat{\mu}} = \sum_z [\frac{b_z}{2} \sum_i q_{iz}^2 - \frac{\gamma c_z}{2}(T_z^2 - \sum_i \tau_{iz}^2)]$; the α, $\hat{\mu}$ and C_i terms cancel. By Cauchy–Schwarz, $\sum_i \tau_{iz}^2 \geq T_z^2/N$, and $\sum q^2 \geq 0$, so $W(\sigma) \geq \Psi_{\hat{\mu}}(\sigma) - \sum_z \frac{\gamma c_z}{2}(1 - \frac{1}{N}) T_z(\sigma)^2$. Next, $\Psi_{\hat{\mu}}(\sigma) \geq \Psi_{\hat{\mu}}(\sigma^*) - \eta$, and $\Psi_{\hat{\mu}}(\sigma^*) \geq W^* - \sum_z \frac{b_z}{2} \sum_i q_{iz}^{*2}$ because $T^2 \geq \sum \tau^2$ for $\tau \geq 0$. Membership of a potential maximiser in every $\mathrm{ICNE}_E(\delta)$ is Proposition S26, extended to M and ID by Lemma S24. □

Theorem S44 (Exact thresholds and regret brackets)**.** *Let p be a pure profile. (a) The exact E-threshold of carrier i is the optimal value of the linear program $A_i^E(p) = \min\{\delta \geq 0 : m \geq 0,\ \xi \in \Xi,\ |m_k - A_{-i,k}(p)| \leq \delta w_k\ \forall k,\ \bar{u}_i(p_i; m, \xi) \geq \bar{u}_i(p'; m, \xi)\ \forall p' \in \mathcal{P}_i\}$ ($+\infty$ if infeasible), and $p \in \mathrm{ICNE}_E(\delta)$ iff $\delta \geq \max_i A_i^E(p)$. With an exact oracle and implicit $\mathcal{P}_i$, Kelley's cutting-plane method on this program terminates finitely. (b) For anchors $m^i \geq 0$, $\xi^i \in \Xi$ with $p_i \in \arg\max \bar{u}_i(\cdot; m^i, \xi^i)$, $A_i^E(p) \leq \|A_{-i}(p) - m^i\|_{w,\infty}$. (c) For a heuristic plan h_i and a valid bound $\mathcal{U}_i \geq \max_{p'} \bar{u}_i(p'; m, \xi)$, the regret of p_i lies in $[\bar{u}_i(h_i) - \bar{u}_i(p_i),\ \mathcal{U}_i - \bar{u}_i(p_i)]$.*

Proof. (a) By Lemma S23, p_i is E-admissible at width δ iff the system with that δ is feasible; feasibility is upward closed in δ (Lemma S24) and the infimum of an LP is attained when finite. Kelley's method adds the oracle's best plan at the current (m, ξ) as a violated constraint; a plan already present cannot be violated, so cuts are distinct and finitely many. (b) The anchor is feasible with $\delta = \|A_{-i}(p) - m^i\|_{w,\infty}$. (c) Immediate from the definitions. □

Theorem S45 (Rounding guarantee under D2)**.** *Under D2, if the relaxation phase returns a mixed equilibrium x, the sparsify phase (a feasibility LP over $\Lambda(\bar{A}(x))$ with plan sets $\operatorname{supp} x_j$) and the round phase return $p \in \prod_j B_j(x)$ with $p \in \mathrm{ICNE}_E(\delta)$ for all $\delta \geq \hat{\delta}(p)$; $\hat{\delta}(p)$ satisfies the second bound of Theorem S27(b) for any choice in the round phase and the third for a successful sample. Sparsify and round run in time polynomial in $\sum_j |\operatorname{supp} x_j|$ and d.*

Proof. The sparsify LP's vertices have $|J| \leq k^\sharp$ by Theorem S39. The rest is Theorem S27(a), (b) and (d). □

The double-oracle relaxation [46] stops, with exact restricted solves and exact oracles, after at most $\sum_i |\mathcal{P}_i|$ rounds at a relaxed equilibrium: at termination every support plan attains the restricted maximum and no plan exceeds it, and each non-terminal round adds a new plan.

S8. Proofs for BRL

Proof of Theorem S1. Write $S = \sum_{a \in r} X_a$ (arcs counted with multiplicity). Fix j and $\Theta = \Theta_j$.

Step 1 (pointwise inequality). For reals x_a, d_a with $\sum_a d_a \leq \Theta$, $(\sum_a x_a - \Theta)^+ \leq (\sum_a (x_a - d_a))^+ \leq \sum_a (x_a - d_a)^+$, because $t \mapsto t^+$ is nondecreasing and subadditive.

Step 2 (upper bound). Let $P \in \mathcal{M}$ with marginals F_a. Taking expectations, $\mathbb{E}_P (S - \Theta)^+ \leq \sum_a \mathbb{E}_{F_a} (X_a - d_a)^+$. Since $F_a \geq \underline{F}_a$ pointwise, F_a is first-order dominated by $\underline{F}_a$, and $x \mapsto (x - d)^+$ is nondecreasing, so $\mathbb{E}_{F_a} (X_a - d_a)^+ \leq \overline{\Pi}_a(d_a)$; likewise $\mathbb{E}_{F_a} X_a \leq \bar{x}_a$. Hence $\mathbb{E}_P \phi(S) \leq c_0 + \gamma \sum_a \bar{x}_a + \sum_j \rho_j \sum_a \overline{\Pi}_a(d_a^j)$ for every P and admissible d^j.

Step 3 (attainment). Let U be uniform on $(0, 1)$, $\bar{X}_a = \overline{Q}_a(U)$ and $\bar{S} = \Sigma(U)$ with $\Sigma(u) = \sum_a \overline{Q}_a(u)$; then $\bar{X}_a \sim \underline{F}_a$ and the law of $(\bar{X}_a)$ belongs to $\mathcal{M}$. Let $u^* = \sup\{u \in (0, 1] : \Sigma(u) \leq \Theta\}$ ($\sup \emptyset = 0$). If $u^* = 1$, put $d_a = \overline{Q}_a(1)$; then $\sum_a d_a \leq \Theta$ and both sides of Step 1 vanish almost surely. If $u^* < 1$, put $\ell_a = \overline{Q}_a(u^*)$ (or $\overline{Q}_a(0^+)$ if $u^* = 0$) and $h_a = \overline{Q}_a(u^{*+})$; then $\sum_a \ell_a \leq \Theta$ when $u^* > 0$ and $\sum_a h_a \geq \Theta$. If $u^* = 0$ choose $d_a \leq \ell_a$ with $\sum_a d_a = \Theta$; otherwise choose $d_a \in [\ell_a, h_a]$ with $\sum_a d_a = \Theta$. On $\{U \leq u^*\}$ all $\bar{X}_a \leq d_a$ and both sides of Step 1 are 0; on $\{U > u^*\}$ all $\bar{X}_a \geq d_a$ and $\sum_a (\bar{X}_a - d_a)^+ = \bar{S} - \Theta = (\bar{S} - \Theta)^+$. So Step 1 holds with equality almost surely, and the comonotone law attains the bound of Step 2; the minima are attained.

Step 4. $\mathbb{E}\phi(\bar{S}) = \int_0^1 \phi(\Sigma(u))du$ because $\bar{S} = \Sigma(U)$.

Step 5 (integrality). With integer quantiles and Θ, the numbers ℓ_a, h_a, $\overline{Q}_a(1)$ are integers and integer vectors with the required properties exist in each case of Step 3. □

Proof of Theorem S2. Substitute the budget form of Theorem S1 into the objective and merge the two minimisations. □

Lemma S46 (Concatenation is inf-convolution)**.** *For a label p and a completion q, $\overline{\mathbb{E}}[\phi(T_{p\oplus q})] - c_0 = f_{p\oplus q}(T_{\max}) = \min_D\{f_p(D) + f_q(T_{\max} - D)\}$.*

Proof. The stop-loss terms are separable over arcs. Split the budget vector into the arcs of p and of q, put $D = \sum_{a\in p} d_a$, minimise first over the arcs of p with budget D and then over those of q with budget $T_{\max} - D$, and apply Theorem S1 to each part. The mean terms add. □

Lemma S47 (Completion window)**.** *Let $C_{\min}(v) \le S_{q,1}$ for every completion q of labels ending at v. For every completion q, the minimum in Lemma S46 is attained at some $D^* \le \overline{D}(p)$; if $D^* < \underline{D}(p)$ then $D^* \le S_{p,1}$.*

Proof. Let $\Phi_q(D) = f_p(D) + f_q(T_{\max} - D)$. The map $D \mapsto f_q(T_{\max} - D)$ is convex and nondecreasing with slope in $[0, \rho]$, equal to ρ when $T_{\max} - D < S_{q,1}$; the slope of f_p lies in $[-\rho, 0]$ and is 0 for $D > S_{p,L}$. (i) On $D > T_{\max} - S_{q,1}$, Φ_q has slope ≥ 0, so a minimiser exists with $D \le T_{\max} - S_{q,1} \le T_{\max} - C_{\min}(v)$. (ii) On $D > S_{p,L}$, Φ_q has slope ≥ 0, so decreasing D to $S_{p,L}$ does not increase it. Applying (i) and then (ii) gives $D^* \le \overline{D}(p)$. The last claim is immediate from $\underline{D}(p) \le S_{p,1}$. □

Proof of Theorem S4. Validity. Let q be a completion of p_2; since $\mathcal{V}(p_1) \subseteq \mathcal{V}(p_2)$ and $\ell(p_1) \le \ell(p_2)$, it is a completion of p_1. By Lemma S47 there is a minimiser $D^* \le \overline{D}(p_2)$ for $p_2 \oplus q$. Let g be the difference in Definition S3. If $D^* \in \mathcal{W}(p_2)$, $g(D^*) \le 0$. Otherwise $D^* < \underline{D}(p_2) \le S_{p_2,1}$; on $(-\infty, S_{p_2,1}]$, f_{p_2} has slope $-\rho$ and f_{p_1} slope $\ge -\rho$, so g is nondecreasing there and $g(D^*) \le g(\underline{D}(p_2)) \le 0$. In both cases, by Lemma S46, $f_{p_1\oplus q}(T_{\max}) - y(p_1) \le f_{p_1}(D^*) + f_q(T_{\max} - D^*) - y(p_1) \le f_{p_2\oplus q}(T_{\max}) - y(p_2)$.

Exactness. Call a label processed if it is popped while not deleted. Let OPT be the optimum and Ω the set of pairs (o, q) with o a created label, q a completion of o and reduced cost of $o \oplus q$ equal to OPT. The pair (depot label, optimal route) is in Ω and the depot label is processed. Among pairs in Ω with o processed choose one with the fewest arcs $|q|$. If $|q| = 1$, closing o records a route of value OPT. If $|q| \ge 2$, let (v, j) be the first arc of q, q' the rest and $o_1 = o \oplus (v, j)$, which is created when o is processed; $(o_1, q') \in \Omega$. If o_1 is processed, $|q'| < |q|$ contradicts minimality. Otherwise o_1 was discarded or deleted because a live label o_2 functionally dominates it; by validity $(o_2, q') \in \Omega$, and if o_2 is not processed it was deleted later by some o_3, with $(o_3, q') \in \Omega$ by transitivity. The moments at which these labels cease to be live strictly increase and labels are finitely many, so the chain ends at a processed label with $(o_k, q') \in \Omega$ and $|q'| < |q|$, a contradiction. □

Proposition S48 (Comparison with vector dominance)**.** *Assume $\mathcal{V}(p_1) \subseteq \mathcal{V}(p_2)$, $\ell(p_1) \le \ell(p_2)$ and $y(p_1) \ge y(p_2)$. (a) If $S_{p_1} \le S_{p_2}$ componentwise, p_1 functionally dominates p_2. (b) In the sector model $X_a = \sum_k \tau_{a,k}\xi_k$ with probability boxes on the multipliers, $\tau_{p_1} \le \tau_{p_2}$ componentwise implies $S_{p_1} \le S_{p_2}$. (c) Neither converse holds.*

Proof. (a) f is nondecreasing in each S_l. (b) The upper quantiles of the multipliers are nonnegative. (c) With $L = 2$, $\omega = (\frac{1}{2}, \frac{1}{2})$, $\gamma = 0$, $\rho = 1$ and equal duals: $S_{p_1} = (1, 1)$ and $S_{p_2} = (0, 2.2)$ give $f_{p_1} \le f_{p_2}$ on $\mathbb{R}$ (values $(1, 1.1)$, $(0, 0.6)$, $(0, 0)$ at $D = 0, 1, 2.2$, constant difference -0.1 below 0) although $S_{p_1,1} > S_{p_2,1}$. □

Proposition S49 (A single risk level is not exact)**.** *For every u, $\min_{\text{plan}} \sum_r \overline{\mathbb{E}}[\phi(T_r)] \geq \max_u \min_{\text{plan}} \sum_r [c_0 + \gamma \bar{x}_r + \rho(\mathcal{H}_r(u) - (1-u)T_{\max})]$, where $\mathcal{H}_r(u) = \int_u^1 \sum_{a \in r} \overline{Q}_a(v)\, dv$; the inequality can be strict and the best fixed-u plan need not be optimal.*

Proof. The maximum over u inside each route is at least its value at a common u, and $\min \max \geq \max \min$. Strictness is shown by exact computations by enumeration and set partitioning (Section S5.1); structurally, the optimal level of a route is its worst-case on-time level, which differs between routes of the same plan. □

S9. Proofs for dual implementation pricing

Notation as in Section S2.4; $m_z^S = b_z + 2\kappa_z$ is the social slope, and $q_{iz}^2 = d_{iz} q_{iz}$ is used throughout. The proofs are taken from the theory notes of the repository, file `theory/new_algorithm_C/theory.tex`.

Proof of Proposition S6. (a) $\tau_{iz} = r q_{iz}$. (b) $s + r t_z = u_z$ and $t_z \geq 0$ because $s \leq u_z$; zones without lots carry no plan. (c) Substitution of $s_z \equiv s$ and $t_z = \vartheta g_z X_z^{\text{NE}}$. □

Proof of Lemma S7. With $q_{iz}^2 = d_{iz} q_{iz}$, $X_z = r Q_z$ and $m_z^S - m_z = \kappa_z$,

$$\Phi_u - \text{SC} = \sum_z \left[\tfrac{m_z}{2} \textstyle\sum_i d_{iz} q_{iz} - \tfrac{\kappa_z}{2} Q_z^2 + u_z Q_z \right].$$

Insert $u_z = \kappa_z Q_z^\circ - \frac{m_z}{2} \bar{d}_z + v_z$ and complete the square: $-\frac{\kappa_z}{2} Q_z^2 + \kappa_z Q_z^\circ Q_z = -\frac{\kappa_z}{2}(Q_z - Q_z^\circ)^2 + \frac{\kappa_z}{2}(Q_z^\circ)^2$, and $\frac{m_z}{2} \sum_i d_{iz} q_{iz} - \frac{m_z}{2} \bar{d}_z Q_z = \sum_i \frac{m_z}{2} (d_{iz} - \bar{d}_z) q_{iz}$. □

Proposition S50 (Carrier-specific charges; classical)**.** *Let σ^* be a pure profile and $U_{iz} = \kappa_z Q_{-i,z}^* - \frac{b_z}{2} d_{iz}$. Under carrier-specific per-delivery charges U_i, for all i and p, p': $J_i^U(p; \sigma_{-i}^*) - J_i^U(p'; \sigma_{-i}^*) = \text{SC}(p, \sigma_{-i}^*) - \text{SC}(p', \sigma_{-i}^*)$. Hence every profile that no single carrier can improve socially is a pure equilibrium.*

Proof. $\text{SC}(p, \sigma_{-i}^*) = R_i(p) + \sum_z q_{iz}(-\alpha_z + m_z^S Q_{-i,z}^* + \frac{m_z^S}{2} d_{iz}) + \text{const}$ and $J_i^U(p) = R_i(p) + \sum_z q_{iz}(-\alpha_z + m_z d_{iz} + m_z Q_{-i,z}^* + U_{iz})$; the brackets coincide because $m_z + \kappa_z = m_z^S$ and $m_z - \frac{b_z}{2} = \frac{m_z^S}{2}$. □

Convexified game. With mixtures $x_i \in \Delta(\mathcal{P}_i)$, $\bar{q}_i = \mathbb{E}_{x_i} q_i$ and $\bar{Q} = \sum_i \bar{q}_i$, the convexified potential and social cost are $\overline{\Phi}_u(x) = \sum_i \mathbb{E}_{x_i}[R_i + \sum_z q_{iz}(-\alpha_z + u_z + \frac{m_z}{2} d_{iz})] + \sum_z \frac{m_z}{2} \bar{Q}_z^2$ and $\overline{\text{SC}}(x) = \sum_i \mathbb{E}_{x_i}[R_i - \sum_z \alpha_z q_{iz}] + \sum_z \frac{m_z^S}{2} \bar{Q}_z^2$. Both are convex, agree with Φ_u and SC at pure profiles, and the identity of Lemma S7 holds for them with Q, q replaced by $\bar{Q}, \bar{q}$.

Proof of Theorem S8. $\overline{\text{SC}}$ is linear in x plus $\sum_z \frac{m_z^S}{2} \bar{Q}_z^2$ with $\bar{Q}$ linear in x, so $\overline{\text{SC}}(x) - \overline{\text{SC}}(x^*) = \langle \nabla \overline{\text{SC}}(x^*), x - x^* \rangle + \sum_z \frac{m_z^S}{2} (\bar{Q}_z - \bar{Q}_z^*)^2$, and the inner product is non-negative by optimality of x^*. Hence $\Delta := \overline{\text{SC}}(x^u) - \overline{\text{SC}}(x^*) \geq \sum_z \frac{m_z^S}{2} (\Delta \bar{Q}_z)^2$. By the identity with $Q^\circ = \bar{Q}^*$, $v = 0$: $\overline{\Phi}_u(x) = \overline{\text{SC}}(x) - \sum_z \frac{\kappa_z}{2} (\bar{Q}_z - \bar{Q}_z^*)^2 + L(x) + \text{const}$. Since x^u minimises $\overline{\Phi}_u$, $\Delta \leq \sum_z \frac{\kappa_z}{2} (\Delta \bar{Q}_z)^2 + L(x^*) - L(x^u)$. Combining, $\sum_z \frac{b_z + \kappa_z}{2} (\Delta \bar{Q}_z)^2 \leq L(x^*) - L(x^u)$; then $\sum_z \frac{\kappa_z}{2} (\Delta \bar{Q}_z)^2 \leq \varrho [L(x^*) - L(x^u)]$ and $\Delta \leq (1 + \varrho)[L(x^*) - L(x^u)]$. Finally $|L(x^*) - L(x^u)| \leq \sum_{i,z} \frac{m_z}{2} |d_{iz} - \bar{d}_z| \, |\bar{q}_{iz}^* - \bar{q}_{iz}^u| \leq E$ because $\bar{q}_{iz} \in [0, d_{iz}]$. □

Proof of Theorem S9. With $\bar{d} = d$, L vanishes on the feasible set, so $E = 0$ in Theorem S8 and every minimiser of $\overline{\Phi}_u$ has aggregate $\bar{Q}^*$ and social cost $\overline{\text{SC}}(x^*)$. Moreover $\overline{\Phi}_u = \overline{\text{SC}} - \sum_z \frac{\kappa_z}{2} (\bar{Q}_z - \bar{Q}_z^*)^2 + \text{const}$; the subtracted term has zero gradient at x^*, so x^* satisfies the first-order optimality condition of the convex function $\overline{\Phi}_u$. □

Proof of Theorem S10. (a) $J_i^u(p) = J_i^0(p) + \langle u, q_i(p)\rangle$, so constraint (i,p) reads $J_i^u(p) \geq J_i^u(\sigma_i^*)$; all of carrier i's constraints hold iff $\min_p J_i^u(p) \geq J_i^u(\sigma_i^*)$, and the violation $J_i^u(\sigma_i^*) - J_i^u(p)$ is maximised by a best response. (b) If a round adds a cut (i, b_i), the current u violates it and satisfies every earlier cut (it is LP-feasible), so the cut is new; there are $\sum_i(|\mathcal{P}_i| - 1)$ non-trivial constraints. Every cut is a valid constraint of $P(\sigma^*)$, so an infeasible LP shows that $P(\sigma^*)$ is empty. □

Proof of Proposition S11. σ is an equilibrium under u iff $u \in P(\sigma)$, so both descriptions of $V^\star$ range over the same pairs (u, σ). The inequality is an inclusion of feasible sets. If the loop ends with $u \in P(\sigma^*)$, the leader attains $\mathrm{SC}(\sigma^*)$. □

Proof of Theorem S12. Best-response dynamics strictly decreases Φ_u on a finite set, so it ends at a pure equilibrium with $\Phi_u(\sigma^u) \leq \Phi_u(\sigma^*)$. Substitute the identity with $Q^\circ = Q^*$ on both sides; the constants cancel and the congestion term vanishes at σ^*. □

Proof of Corollary S13. Count the calls of the three phases; Theorem S10 bounds the rounds, and each improvement of the fallback dynamics costs at most N calls. □

Proof of Theorem S14. By Lemma S25, carrier i's Γ-maximin payoff is its precise payoff minus $\delta\langle a_i(p), Mw\rangle$ (traffic already enters the route loss at its upper expectation). With $a_i = (q_i, rq_i)$, $\delta\langle a_i(p), Mw\rangle = \langle \hat{s}(\delta), q_i(p)\rangle$, so the Γ-maximin cost under u is the precise cost under $u + \hat{s}(\delta)$, and u implements σ^* iff $u + \hat{s}(\delta) \in P(\sigma^*)$. The social cost does not contain the surcharge. The last claim is Lemma S24: an equilibrium of the precise game lies in $\mathrm{ICNE}_C(\delta)$ for $C \in \{E, M, ID\}$ and every δ. □

Proof of Proposition S15. The first claim is Theorem S9 with a uniform tangent charge. For the second, $Q_z^{\mathrm{full}} = n_z d_z$ with equal lots, and without congestion ($m_z \approx b_z$ when $\kappa_z \ll b_z$) $u_z = -\frac{b_z}{2} d_z = -\frac{\alpha_z}{4n_z} = -0.375\,\mathrm{fee}/n_z$. The last claim is Proposition S6. □

S10. Proofs for aggregate-response pricing

Notation as in Section S2.4: $q_{iz}^2 = d_{iz} q_{iz}$, $\kappa_z = g_z r^2$, $m_z = b_z + \kappa_z$, $m_z^S = b_z + 2\kappa_z$, and Φ_u the exact potential of the game under the per-delivery-equivalent charge u. For a reference $\bar{d} \in \mathbb{R}^Z$, $L_{\bar{d}}(\sigma) = \sum_{i,z} \frac{m_z}{2}(d_{iz} - \bar{d}_z) q_{iz}$ and $E_{\bar{d}} = \sum_{i,z} \frac{m_z}{2} |d_{iz} - \bar{d}_z| d_{iz}$. The convexified potential $\overline{\Phi}_u$ and social cost $\overline{\mathrm{SC}}$ are those of Section S9. The proofs are taken from `theory/new_algorithm_D/theory.tex`.

Lemma S51. *G is convex and piecewise linear on $\mathcal{Q}$; F is proper, closed and 1-strongly convex in $\|\cdot\|_m$, i.e. $F - \frac{1}{2}\|\cdot\|_m^2$ is convex. Moreover $\min_{x \in \mathcal{X}} \overline{\Phi}_u(x) = \min_Q \{F(Q) + \langle u, Q\rangle\}$ and $\min_{x \in \mathcal{X}} [\overline{\mathrm{SC}}(x) + L_{\bar{d}}(x)] = \min_Q \{F(Q) + g(Q)\}$ with $g(Q) = \sum_z [\frac{\kappa_z}{2} Q_z^2 - \frac{m_z}{2} \bar{d}_z Q_z]$, and in both cases x is a minimiser on the left if and only if $\bar{Q}(x)$ is a minimiser on the right and x attains the minimum defining G at $Q = \bar{Q}(x)$.*

Proof. G is the optimal value of a linear program in x whose right-hand side Q enters through $\bar{Q}(x) = Q$; such a value function is convex and piecewise linear on the set of right-hand sides for which the program is feasible, which is $\mathcal{Q}$. The quadratic part of F is $\frac{1}{2}\|Q\|_m^2$ and the rest is convex, so $F - \frac{1}{2}\|\cdot\|_m^2$ is convex; F is closed because G is continuous on the closed set $\mathcal{Q}$. For the identities, write $\frac{m_z}{2} d_{iz} = \frac{m_z}{2}(d_{iz} - \bar{d}_z) + \frac{m_z}{2} \bar{d}_z$ in $\overline{\Phi}_u$ and $\frac{m_z^S}{2} = \frac{m_z}{2} + \frac{\kappa_z}{2}$ in $\overline{\mathrm{SC}} + L_{\bar{d}}$; in both the objective depends on x only through $\bar{Q}(x)$ plus the quantity minimised in the definition of G. □

Proof of Theorem S17. By Lemma S51, $F + \langle u, \cdot\rangle$ is strongly convex on the compact set $\mathcal{Q}$, so it has a unique minimiser $Q(u)$, and every convexified equilibrium has $\bar{Q}(x) = Q(u)$. For closed proper strongly convex F the conjugate F^* is finite (the domain is bounded) and differentiable with

$\nabla F^*(y) = \arg\max_Q\{\langle y, Q\rangle - F(Q)\}$ [56, Theorem 26.3]; with $y = -u$ this is $Q(u)$. The optimality conditions read $-u_k \in \partial F(Q_k)$; strong convexity gives $\langle(-u_1)-(-u_2), Q_1 - Q_2\rangle \geq \|Q_1 - Q_2\|_m^2$, and by Cauchy–Schwarz in the m-inner product $\langle u_2 - u_1, Q_1 - Q_2\rangle \leq \|u_2 - u_1\|_{m^{-1}}\|Q_1 - Q_2\|_m$, which gives the Lipschitz bound.

Let $D(u) = F^*(-u) + g^*(u)$ with $g^*(u) = \sum_z (u_z + \frac{m_z}{2}\bar{d}_z)^2/(2\kappa_z)$ (maximise $\langle u, Q\rangle - g(Q)$ coordinatewise). D is differentiable with $\nabla D(u) = -Q(u) + \kappa^{-1} \odot (u + \frac{m}{2}\bar{d})$, which is observable, and strongly convex because g^* has Hessian $\mathrm{diag}(1/\kappa)$. Fenchel duality [56, Corollary 31.2.1], with attainment because g is finite and continuous, gives $\min_Q\{F + g\} = -\min_u D(u)$. The minimiser u^* is unique and characterised by $\nabla D(u^*) = 0$, i.e. $u^* = \kappa \odot Q(u^*) - \frac{m}{2} \odot \bar{d} =: T(u^*)$; any fixed point of T is such a zero. At the saddle point the primal minimiser is $Q(u^*)$, which minimises $F+g$, i.e. $\overline{\mathrm{SC}} + L_{\bar{d}}$ by Lemma S51. For welfare, let x^* minimise $\overline{\mathrm{SC}}$ and x^∞ be a convexified equilibrium under u^*; then $\overline{\mathrm{SC}}(x^\infty) - \overline{\mathrm{SC}}(x^*) \leq L_{\bar{d}}(x^*) - L_{\bar{d}}(x^\infty) \leq \sum_{i,z} \frac{m_z}{2}|d_{iz} - \bar{d}_z||\bar{q}^*_{iz} - \bar{q}^\infty_{iz}| \leq E_{\bar{d}}$, because $\bar{q}_{iz} \in [0, d_{iz}]$; with symmetric lots $L_{\bar{d}} \equiv 0$. □

Proof of Theorem S18. Let $K = \mathrm{diag}(\kappa)$ and equip $\mathbb{R}^Z$ with $\langle\cdot,\cdot\rangle_{\kappa^{-1}}$. The gradient of D in this geometry is $K\nabla D(u) = u + \frac{m}{2}\bar{d} - \kappa \odot Q(u)$, so the rule (S5) is $u^{t+1} = u^t - \theta K \nabla D(u^t)$, gradient descent with step θ. Since $\kappa_z \leq \varrho m_z$ and $1/m_z \leq \varrho/\kappa_z$ on active zones, the Lipschitz bound of Theorem S17 gives $\|Q(u_1) - Q(u_2)\|_\kappa \leq \sqrt{\varrho}\|Q(u_1) - Q(u_2)\|_m \leq \sqrt{\varrho}\|u_1 - u_2\|_{m^{-1}} \leq \varrho\|u_1 - u_2\|_{\kappa^{-1}}$. Hence $u \mapsto \kappa \odot Q(u)$ is ϱ-Lipschitz from $\|\cdot\|_{\kappa^{-1}}$ to itself (note $\|\kappa \odot v\|_{\kappa^{-1}} = \|v\|_\kappa$), $K\nabla D$ is $(1+\varrho)$-Lipschitz, and D is strongly convex with modulus 1 in this geometry. (a) T is ϱ-Lipschitz and $u^* = T(u^*)$. (b) For a μ-strongly convex function with L-Lipschitz gradient, gradient descent with step $\theta \leq 2/(\mu + L)$ satisfies $\|u^{t+1} - u^*\|^2 \leq (1 - \frac{2\theta\mu L}{\mu + L})\|u^t - u^*\|^2$ [57, Theorem 2.1.15]; the proof uses only an inequality valid in any inner-product space. With $\mu = 1$, $L = 1 + \varrho$ and $\theta = 2/(2+\varrho)$ the factor is $1 - 4(1+\varrho)/(2+\varrho)^2 = \varrho^2/(2+\varrho)^2$. □

Proof of Theorem S19. (a) Apply Lemma S7 with $Q^\circ = Q(\sigma)$. Since σ is an equilibrium, $\Phi_u(\sigma) \leq \Phi_u(\sigma')$ [36]. The term $\sum_z \frac{\kappa_z}{2}(Q_z - Q^\circ_z)^2$ vanishes at σ and equals $\sum_z \frac{\kappa_z}{2}\Delta Q_z^2$ at σ'; rearranging the identity for σ and σ' gives the inequality. The last bound uses $|q_{iz}(p) - q_{iz}(\sigma_i)| \leq d_{iz}$.

(b) $u^{t+1} = \kappa \odot Q^t - \frac{m}{2}\bar{d} = T(u^t) + \kappa \odot e^t$ and $u^* = T(u^*)$, so the first inequality follows from Theorem S18(a); iterate for the limsup. For the gap bound let $h = F + \langle u^t, \cdot\rangle$, which is 1-strongly convex in $\|\cdot\|_m$ and minimised at $Q(u^t)$, so $h(Q) \geq h(Q(u^t)) + \frac{1}{2}\|Q - Q(u^t)\|_m^2$. A pure profile is a point of $\mathcal{X}$ with $\bar{Q} = Q^t$, so $\overline{\Phi}_{u^t}(\sigma^t) \geq h(Q^t)$ while $\min \overline{\Phi}_{u^t} = h(Q(u^t))$ (Lemma S51); with $\Phi_{u^t}(\sigma^t) = \overline{\Phi}_{u^t}(\sigma^t)$ at a pure profile, $\eta_t \geq \frac{1}{2}\|e^t\|_m^2$ and $\|e^t\|_\kappa \leq \sqrt{\varrho}\|e^t\|_m \leq \sqrt{2\varrho\eta_t}$.

(c) By Theorem S17, $Q = Q(u)$ iff $-u \in \partial F(Q)$. F is polyhedral plus a quadratic on the polytope $\mathcal{Q}$, so $\partial F(Q) \neq \emptyset$ on $\mathcal{Q}$ and the graph of the response map is the graph of $-\partial F$ on $\mathcal{Q}$. A closed proper convex function is determined by its subdifferential up to an additive constant [56, Theorem 24.9]. The convexified optimum minimises $F + g$ with g known, and u^* is the unique zero of ∇D; both are unchanged by a constant shift of F. Example S52 shows that the pure optimum is not determined. □

Example S52. One zone, $\kappa = 0$, slope $b = m$, intercept α, reference $\bar{d} = d$. Instance A: two carriers, each with lot d and plans "stay out" ($R = 0$) and "serve" ($R = r$). Instance B: one carrier with lot $2d$ and plans "stay out" ($R = 0$) and "serve" ($R = 2r - md^2$), with $r > md^2/2$. In both, $\mathcal{Q} = [0, 2d]$ and G is linear with $G_A(Q) = G_B(Q) = \frac{r}{d}Q$, so $F_A = F_B$ and the response maps coincide. The pure social costs are $\mathrm{SC}_A(Q) = \frac{r}{d}Q - \alpha Q + \frac{m}{2}Q^2$ on $Q \in \{0, d, 2d\}$ and $\mathrm{SC}_B(Q) = (2r - md^2)\mathbf{1}[Q = 2d] - \alpha Q + \frac{m}{2}Q^2$ on $\{0, 2d\}$. With $\alpha = \frac{r}{d} + md$, $Q = d$ is the unique pure optimum of A with value $-\frac{m}{2}d^2$, while $Q = 2d$ is the unique pure optimum of B with value $-md^2$. The optima differ in aggregate and value although nothing observable in the convexified game differs.

Lemma S53 (Lot estimates)**.** *(a) If lots are symmetric within zones, $Q_z/n_z = \bar{d}_z$ in every served zone. (b) If between two observations n_z changes by exactly one, $|Q_z^t - Q_z^{t-1}|$ is the lot of the carrier that entered or left; if $n_z = 1$, Q_z is the lot of the serving carrier. (c) For a zone*

never served the observation carries no information about its lots; the bound of Theorem S17 holds for any reference vector, with $E_{\bar{d}}$ for the vector actually used.

Proof. (a) $Q_z = n_z \bar{d}_z$. (b) The other lots in z are unchanged, so the difference of the sums is the changed term. (c) Two instances that differ only in the lots of a never-served zone are indistinguishable; the bound was shown for an arbitrary reference vector. □

Proof of Proposition S20. By Lemma S25, the Γ-maximin cost gains the surcharge

$$\delta\langle a_i(p), Mw\rangle = \langle \hat{s}(\delta), q_i(p)\rangle,$$

with $\hat{s}_z(\delta) = \delta\hat{w}_z$, $\hat{w}_z = b_z w_{z,Q} + g_z r\, w_{z,X}$, linear in deliveries; so the response to u is the precise response to $u+\hat{s}$, the map $u \mapsto T(u+\hat{s})$ is ϱ-Lipschitz and the rule contracts as before, with a fixed point at which $u+\hat{s}$ is a zero of the dual gradient of $\min_Q F+g+\langle \hat{s}, Q\rangle$. For the loss, let Q_0 and Q_δ minimise $H = F+g$ and $H+\langle \hat{s}, \cdot\rangle$, both 1-strongly convex in $\|\cdot\|_m$. Strong convexity at Q_δ gives $H(Q_\delta) - H(Q_0) \le \langle \hat{s}, Q_0 - Q_\delta\rangle - \frac{1}{2}\|Q_0 - Q_\delta\|_m^2 \le \|\hat{s}\|_{m^{-1}}\|Q_0 - Q_\delta\|_m - \frac{1}{2}\|Q_0 - Q_\delta\|_m^2 \le \frac{1}{2}\|\hat{s}\|_{m^{-1}}^2$.

For E-admissibility, write $\Delta_z = u_z - u_z^*$ for a posted charge u and a pure equilibrium σ^* under u^* with precise beliefs. Carrier i's cost of plan p at charge u and belief (μ_Q, μ_X) about rivals is $R_i(p) + \sum_z q_{iz}(p)(-\alpha_z + m_z d_{iz} + b_z\mu_{Q,z} + g_z r\mu_{X,z} + u_z)$. Choose $\mu_{Q,z} = Q_{-i,z} - \Delta_z w_{z,Q}/\hat{w}_z$ and $\mu_{X,z} = X_{-i,z} - \Delta_z w_{z,X}/\hat{w}_z$. Then $b_z\mu_{Q,z} + g_z r\mu_{X,z} + u_z = b_z Q_{-i,z} + g_z r X_{-i,z} + u_z^*$, so every plan's cost at this belief and charge u equals its cost at the realised aggregate and charge u^*, where σ_i^* is optimal. The belief lies in the box iff $|\Delta_z| \le \delta\hat{w}_z = \hat{s}_z(\delta)$, and it is nonnegative when $\Delta_z \le 0$, or when $Q_{-i,z} \ge \Delta_z w_{z,Q}/\hat{w}_z$ and $rQ_{-i,z} \ge \Delta_z w_{z,X}/\hat{w}_z$ for $\Delta_z > 0$. By Lemma S23, σ_i^* is E-admissible. The round count follows from Theorem S18(a) and $|u_z - u_z^*| \le \sqrt{\kappa_z}\|u - u^*\|_{\kappa^{-1}}$. □

Proof of Theorem S16. (a) $J_i^T(p; \sigma_{-i}) = R_i(p) + \sum_z q_{iz}(-\alpha_z + m_z d_{iz} + m_z Q_{-i,z} + \kappa_z Q_{-i,z} - \frac{b_z}{2} d_{iz}) = R_i(p) + \sum_z q_{iz}(-\alpha_z + \frac{m_z^S}{2} d_{iz} + m_z^S Q_{-i,z})$, using $q_{iz}^2 = d_{iz} q_{iz}$, $m_z - \frac{b_z}{2} = \frac{m_z^S}{2}$ and $m_z + \kappa_z = m_z^S$. On the other side $\mathrm{SC}(p, \sigma_{-i}) - \mathrm{SC}(\emptyset_i, \sigma_{-i}) = R_i(p) + \sum_z [-\alpha_z q_{iz} + \frac{m_z^S}{2}((q_{iz} + Q_{-i,z})^2 - Q_{-i,z}^2)]$, which is the same expression. The right side differs from $\mathrm{SC}(p, \sigma_{-i})$ by a term independent of p, which is the definition of an exact potential. (b) Monderer–Shapley for the potential SC on a finite profile set. (c) Immediate from the first expression in (a). □

S11. Zone-market clearing (ZMC)

ZMC is the first phase of the equilibrium engine (Section S2.3). It is stated here with its guarantees and proofs; Remark S5 gives the negative empirical result.

ZMC targets the D1 game in cost form. Let $c_i(p) = -\pi_i(p) + \gamma\langle t_i(p), \hat{\xi}\rangle$ (charges included), $\tilde{c}_i = c_i - \frac{1}{2}\|a_i\|_M^2$, and assume $M_k > 0$ on the active coordinates. Then $\Phi(\sigma) = \sum_i \tilde{c}_i(\sigma_i) + \frac{1}{2}\|A(\sigma)\|_M^2 = -\Psi_{\hat{\xi}}(\sigma)$ is an exact potential (Proposition S26). For a common price $\lambda \in \mathbb{R}^d$, the pricing oracle returns $O_i(\lambda) = \arg\min_p\{\tilde{c}_i(p) + \langle \lambda, a_i(p)\rangle\}$, which is the oracle of Assumption A3 with $Q = M$. Let $V_i(\lambda)$ be its value, $\mathcal{D}(\lambda) = \sum_i V_i(\lambda) - \frac{1}{2}\|\lambda\|_{M^{-1}}^2$ the price dual of the convexified potential, λ^* its unique maximiser, and $\bar{\Phi}^* = \max \mathcal{D} \le \min \bar{\Phi}$ (Lemma S57).

Theorem S54 (Clearing identity)**.** *Let $\sigma_i \in O_i(\lambda)$ for all i and let $e = M^{-1}\lambda - A(\sigma)$ be the clearing error (supply minus demand at price λ). Then (a) $\Phi(\sigma) - \mathcal{D}(\lambda) = \frac{1}{2}\|e\|_M^2$, so $\Phi(\sigma) - \min\Phi \le \frac{1}{2}\|e\|_M^2$; (b) carrier i's Nash regret is at most $\frac{1}{2}\sum_{k\in\mathcal{Z}_i} M_k e_k^2$, where $\mathcal{Z}_i$ is the set of coordinates in which its plans differ; (c) σ_i is an exact best response to the belief $m_i = M^{-1}\lambda - a_i(\sigma_i) = A_{-i}(\sigma) + e$, hence E-admissible at every width $\delta \ge \max_{k\in\mathcal{Z}_i} |e_k|/w_k$ (if $m_{ik} \ge 0$ on $\mathcal{Z}_i$); (d) $e = 0$ implies that σ minimises Φ and is a pure Nash equilibrium; (e) conversely, exact clearing is attainable if and only if the convexification of the potential is tight at a pure profile.*

Algorithm 4 ZMC: zone-market clearing for pure equilibria of the D1 game

Require: pricing and best-response oracles; rounds T

$l \leftarrow 0$, $u \leftarrow MA^{\max}$, $\lambda \leftarrow (l+u)/2$; plan pools $\mathcal{Q}_i \leftarrow \emptyset$

for $t = 0, \dots, T-1$ **do**

  for all i in parallel: $\sigma_i^t \leftarrow O_i(\lambda)$; $\mathcal{Q}_i \leftarrow \mathcal{Q}_i \cup \{\sigma_i^t\}$ ▷ N routing solves

  $s \leftarrow A(\sigma^t) - M^{-1}\lambda$; update every bracket by the sign of s_k (Theorem S55)

  if t odd: $\lambda \leftarrow (l+u)/2$; **else** for each k: secant step on an isotonic fit of (λ_k, A_k), clipped to $[l_k, u_k]$

$\sigma^{\text{price}} \leftarrow$ the pricing profile with the smallest $\frac{1}{2}\|e^t\|_M^2$

$\sigma^{\text{round}} \leftarrow$ restricted master over the pools and derandomised rounding (no oracle calls)

$\sigma \leftarrow$ the better of the two; sequential best responses until no carrier improves

return σ

Part (a) is Fenchel–Young equality and a standard duality-gap fact. The game-specific content is in (b) and (c): the own-contribution term of the absorption bound cancels, because under D1 the own quadratic term turns a pricing-optimal plan into a best response to the anchor m_i. Finding a good pure equilibrium thus becomes a price search: find one λ at which the carriers' pricing answers nearly clear every zone. Because $\mathcal{D}$ is concave, the aggregate response is antitone in λ, and along each coordinate the sign of the residual $s_k = A_k(\lambda) - \lambda_k/M_k$ localises λ_k^* up to $M_k\chi_k$, where χ_k is the cross-zone variation: how far other coordinates' prices can move the response in coordinate k at a fixed own price.

Theorem S55 (Bracket invariant and dimension-free rate)**.** *Start from brackets $[0, M_k A_k^{\max}]$ and, after each round, move the lower (upper) end of bracket k to λ_k when $s_k > 0$ ($s_k < 0$). Then $\operatorname{dist}(\lambda_k^*, [l_k, u_k]) \le M_k\chi_k$ after every round. If the bracket midpoint is queried at least every other round, after T rounds (NT oracle calls, independent of d) every point of the bracket satisfies $|\lambda_k - \lambda_k^*| \le M_k A_k^{\max} 2^{-\lceil T/2 \rceil} + M_k\chi_k$ for all k. If plan sets are separable across coordinates, $\chi = 0$.*

The floor $M_k\chi_k$ is genuine when zones are complementary (Example S58). Algorithm 4 alternates midpoint steps (for the rate) with secant steps on isotonic fits of each coordinate's response (for speed), then rounds over the collected plans and finishes with best-response polish.

Theorem S56 (Guarantee of ZMC)**.** *With exact oracles, ZMC uses NT pricing calls before the polish; the polish stops after finitely many best-response calls at a pure Nash equilibrium σ with $\Phi(\sigma) \le \bar{\Phi}^* + \frac{1}{2}\min_t \|e^t\|_M^2$; and $\sigma \in \mathrm{ICNE}_C(\delta)$ for all $\delta \ge 0$ and $C \in \{E, M, ID\}$. Run on the surcharged game of Lemma S25, it returns an exact Γ-maximin equilibrium at width δ.*

Charges translate the supply line: under charges t the equilibrium gross price is 1-Lipschitz in t and convexified aggregates respond monotonically (Proposition S59), so brackets and plan pools warm-start the next evaluation when a city leader searches over charges.

Prior art and limits. Price decomposition with a pricing oracle is classical (Dantzig–Wolfe, column generation and bundle methods; 58); learning separable piecewise-linear concave approximations from sampled slopes is SPAR [59]; Walrasian prices with indivisible goods are computed from demand oracles by Paes Leme and Wong [60]; and Duguet et al. [61] relate the regret of price-allocation pairs to duality and integrality gaps in combinatorial markets, which is the closest conceptual neighbour of Theorem S54(a)–(b). We claim novelty only for the own-term cancellation with credal anchors (Theorem S54(b),(c),(e)), the sign-bracket invariant with the cross-variation floor (Theorem S55) and the method they define. ZMC has no D2 version, does not bound the clearing error of pure answers a priori, and spends NT calls even on easy games.

Proofs

Lemma S57 (Price duality)**.** *Let $\bar{\Phi}(x) = \sum_i \sum_p x_i(p)\tilde{c}_i(p) + \frac{1}{2}\|\bar{A}(x)\|_M^2$ on $\prod_i \Delta(\mathcal{P}_i)$, $\bar{\Phi}^* = \min \bar{\Phi}$ and $\mathcal{B} = \prod_k [0, M_k A_k^{\max}]$ with $A_k^{\max} = \sum_i \max_p a_{ik}(p)$. (a) $\mathcal{D}$ is strongly concave with a unique*

maximiser λ^*, and $\max \mathcal{D} = \bar{\Phi}^* \le \min_\sigma \Phi(\sigma)$. (b) All minimisers of $\bar{\Phi}$ share the aggregate $\bar{A}^*$, and $\lambda^* = M\bar{A}^* \in \mathcal{B}$. (c) $M^{-1}\lambda^* \in R(\lambda^*)$, where $R(\lambda) = \partial^+ \sum_i V_i(\lambda)$.

Proof. (a) $V = \sum_i V_i$ is concave and finite and $\frac{1}{2}\|\lambda\|^2_{M^{-1}}$ strongly convex, so $\mathcal{D}$ is strongly concave, continuous and coercive, with a unique maximiser. For $L(x,\lambda) = \sum_i \langle x_i, \tilde{c}_i\rangle + \langle \lambda, \bar{A}(x)\rangle - \frac{1}{2}\|\lambda\|^2_{M^{-1}}$, affine in x on a compact convex set and concave in λ, Sion's theorem gives $\min_x \sup_\lambda L = \sup_\lambda \min_x L$. For fixed x the supremum is at $\lambda = M\bar{A}(x)$ with value $\bar{\Phi}(x)$; for fixed λ, $\min_x L = \mathcal{D}(\lambda)$ because a linear function on a product of simplices is minimised at a vertex carrier by carrier. Pure profiles are vertices, where $\bar{\Phi} = \Phi$. (b) Two minimisers with different aggregates would contradict strict convexity of $\frac{1}{2}\|\cdot\|^2_M$. A minimiser x^* and λ^* form a saddle point, and the unique maximiser of $L(x^*,\cdot)$ is $M\bar{A}(x^*)$. Bounds on $\bar{A}^*$ follow from nonnegativity. (c) This is $0 \in \partial^+\mathcal{D}(\lambda^*) = R(\lambda^*) - M^{-1}\lambda^*$, by the sum rule for concave functions [56]. □

Proof of Theorem S54. For any pure σ and λ, with $A = A(\sigma)$ and $e = M^{-1}\lambda - A$,

$$\begin{aligned}\textstyle\sum_i[\tilde{c}_i(\sigma_i) + \langle \lambda, a_i(\sigma_i)\rangle] - \tfrac{1}{2}\|\lambda\|^2_{M^{-1}} &= \textstyle\sum_i \tilde{c}_i(\sigma_i) + \langle M(A+e), A\rangle - \tfrac{1}{2}\|A+e\|^2_M \\ &= \Phi(\sigma) - \tfrac{1}{2}\|e\|^2_M. \qquad (*)\end{aligned}$$

(a) If $\sigma_i \in O_i(\lambda)$ the left side is $\mathcal{D}(\lambda) \le \bar{\Phi}^* \le \min\Phi$. (b) Fix i, $a = a_i(\sigma_i)$, $a' = a_i(p)$, $\Delta = a' - a$ (zero outside $\mathcal{Z}_i$). Since $MA_{-i} = \lambda - M(a+e)$, the change in carrier i's cost $c_i + \langle a_i, MA_{-i}\rangle$ equals $[\tilde{c}_i(p) + \langle \lambda, a'\rangle] - [\tilde{c}_i(\sigma_i) + \langle \lambda, a\rangle] + \frac{1}{2}\|\Delta\|^2_M - \langle \Delta, Me\rangle \ge -\frac{1}{2}\sum_{k\in\mathcal{Z}_i} M_k e_k^2$, minimising coordinatewise at $\Delta_k = e_k$. (c) For any p, $c_i(p) + \langle a', Mm_i\rangle = [\tilde{c}_i(p) + \langle \lambda, a'\rangle] + \frac{1}{2}\|a' - a\|^2_M - \frac{1}{2}\|a\|^2_M$, minimised by σ_i. Coordinates outside $\mathcal{Z}_i$ shift all plans' costs equally, so the belief can be replaced by one equal to A_{-i} there; within $\mathcal{Z}_i$ it deviates from A_{-i} by $|e_k| \le \delta w_k$, and under D1 the Dirac law at it lies in the credal set (Lemma S23). (d) If $e = 0$, $\Phi(\sigma) = \mathcal{D}(\lambda) \le \bar{\Phi}^* \le \min\Phi \le \Phi(\sigma)$; a global minimiser of an exact potential is a pure equilibrium, and uniqueness gives $\lambda = \lambda^*$. (e) If $\Phi(\sigma^\circ) = \bar{\Phi}^*$, Lemma S57(b) gives $\lambda^* = MA(\sigma^\circ)$, so $e = 0$, and $(*)$ shows that $\sum_i\{\tilde{c}_i(\sigma_i^\circ) + \langle \lambda^*, a_i(\sigma_i^\circ)\rangle - V_i(\lambda^*)\} = 0$, a sum of nonnegative terms. □

Proof of Theorem S55. Let $\chi_k = \sup\{\sup_{r\in R(\lambda)} \inf_{r'\in R(\lambda')} |r_k - r'_k| : \lambda, \lambda' \in \mathcal{B},\ \lambda_k = \lambda'_k\}$. *Key inequality.* Let a query $\lambda \in \mathcal{B}$ have $\lambda_k = \mu$ and response $r \in R(\lambda)$. Suppose $s_k > 0$ but $\mu - \lambda^*_k > M_k\chi_k$. Let λ' equal λ^* except $\lambda'_k = \mu$; then $\lambda' \in \mathcal{B}$. With $r^* = M^{-1}\lambda^* \in R(\lambda^*)$, antitonicity of R gives $r'_k \le \lambda^*_k/M_k$ for every $r' \in R(\lambda')$, and the definition of χ_k gives $r' \in R(\lambda')$ with $r_k \le r'_k + \chi_k + \eta$ for any $\eta > 0$. Hence $s_k = r_k - \mu/M_k \le -(\mu - \lambda^*_k - M_k\chi_k)/M_k + \eta < 0$ for small η, a contradiction. The case $s_k < 0$ is symmetric, and $s_k = 0$ combines both. *Induction.* Initially $\lambda^*_k \in [0, M_kA^{\max}_k]$. If $s_k > 0$ the new bracket is $[\mu, u_k]$; if $\lambda^*_k \ge \mu$ the distance is unchanged, and otherwise it is $\mu - \lambda^*_k \le M_k\chi_k$. The other cases are symmetric. *Rate.* Midpoint rounds halve every bracket and other rounds never enlarge it, so after T rounds with at least $\lceil T/2\rceil$ midpoints the width is at most $M_kA^{\max}_k 2^{-\lceil T/2\rceil}$. If plan sets are separable, V_i is separable and the projection of $R(\lambda)$ on coordinate k depends on λ_k only, so $\chi_k = 0$. □

Example S58 (The floor is genuine)**.** Let $d = 2$, $M = I$ and one carrier with plans 0 ($\tilde{c} = 0$, $a = (0,0)$) and b ($\tilde{c} = -1.5$, $a = (1,1)$). Then $\lambda^* = (0.75, 0.75)$, $\chi_1 = \chi_2 = 1$, and the queries $(0.7, 0.9)$, $(0.6, 1)$, $(0.5, 1)$ all return plan 0 (tie at the last resolved towards 0), so the bracket update sets $u_1 = 0.5$ and excludes λ^*_1 by 0.25, within the floor but positive.

Proof of Theorem S56. Each round makes one pricing call per carrier. The polish is sequential best-response dynamics in a finite exact-potential game; every accepted change strictly decreases Φ, which takes finitely many values, so it stops at a pure Nash equilibrium. The polish does not increase Φ, and the start has $\Phi \le \Phi(\sigma^{\text{price}}) = \mathcal{D}(\lambda^{t^\star}) + \frac{1}{2}\|e^{t^\star}\|^2_M \le \bar{\Phi}^* + \frac{1}{2}\min_t \|e^t\|^2_M$ by Theorem S54(a) and Lemma S57(a). An exact pure equilibrium makes each plan optimal at the centre belief, which lies in every credal box; the Γ case is Lemma S25. □

Proposition S59 (Lipschitz dependence on charges)**.** *With charges t on the contributions, let $\mu^*(t) = \lambda^*(t) + t$. Then $\|\mu^*(t) - \mu^*(t')\|_{M^{-1}} \le \|t - t'\|_{M^{-1}}$ and $\langle \bar{A}^*(t) - \bar{A}^*(t'), t - t' \rangle \le 0$.*

Proof. With charges, $V^t(\lambda) = V(\lambda + t)$, so the equilibrium condition reads $r = M^{-1}(\mu - t) \in R(\mu)$. For $r' = M^{-1}(\mu' - t') \in R(\mu')$, antitonicity gives $\langle r - r', \mu - \mu' \rangle \le 0$, i.e. $\|\mu - \mu'\|^2_{M^{-1}} \le \langle M^{-1}(t - t'), \mu - \mu' \rangle \le \|t - t'\|_{M^{-1}} \|\mu - \mu'\|_{M^{-1}}$. For the second claim, $\langle r - r', t - t' \rangle = \langle r - r', (\mu - \mu') - M(r - r') \rangle \le -\|r - r'\|^2_M \le 0$. □

S12. Machine-checked statements

`lake build CCU` succeeds, no imported file contains an unproved step, and the theorems listed below depend only on the axioms `propext`, `Classical.choice` and `Quot.sound`. "Full" means the paper's statement for arbitrary finite index sets over the reals; "core" means a finite or algebraic part of the theorem.

- *Credal results* (LeanTargets, Radius): E-admissibility implies maximality and $\mathrm{ICNE}_E(\delta)$ is monotone in δ (full); the core of Theorem S27(a); the Shapley–Folkman lemma for finite sets (Theorem S39, full); the exact potential identity under D1 (Proposition S26, full); the kernel equivalence of Theorem S31 (core, no hardness); the arithmetic cores of Propositions S34 and S35; and, per coordinate, the first, decomposition and second terms of Theorem S27(b), and the $N - 1$ term of (c), given the rounding data.
- *DIP*: Proposition S6(a),(b) and Lemma S7 (full), the convexified identity, Theorem S9 over an abstract convex set (full), Theorem S10(a) (full, without the charge box) and the counting core of part (b).
- *Marginal tariff*: Theorem S16(a)–(c) and the tariff identity with the revenue formula (full), including termination of improvement dynamics.
- *ARP*: the strong-monotonicity inequality behind Theorem S17 (core), and the plain and damped contraction of Theorem S18 from that hypothesis (full); the damped constant proved, $1 - \theta$ for $0 < \theta \le 2/(2 + \varrho)$, is sharper than the paper's.
- *BRL*: the budget identity for a finite level grid, the budget split of the inf-convolution, and the completion window and functional-dominance validity with the slope conditions taken as hypotheses (core), and the comparison with quantile-vector dominance.

Not formalised: the asymmetric-lot bound of DIP (Theorem S8) and the LP and ε parts of Theorem S10; Theorem S17 beyond the monotonicity inequality (conjugates, the proximal form) and the pure-game theorems of ARP; probability boxes, upper previsions, quantile functions and the exactness proof of BRL; the Hoeffding term, the linear-programming vertex construction and the NP-completeness statement itself (Mathlib has no framework for polynomial-time reductions); and all computational results. Lean checks the stated cores, not the full theorems.

References

[1] A. Dvoretzky, J. Kiefer, J. Wolfowitz, Asymptotic minimax character of the sample distribution function and of the classical multinomial estimator, The Annals of Mathematical Statistics 27 (1956) 642–669. doi:10.1214/aoms/1177728174.

[2] P. Massart, The tight constant in the Dvoretzky–Kiefer–Wolfowitz inequality, The Annals of Probability 18 (1990) 1269–1283. doi:10.1214/aop/1176990746.

[3] I. Meilijson, A. Nádas, Convex majorization with an application to the length of critical paths, Journal of Applied Probability 16 (1979) 671–677. doi:10.2307/3213097.

[4] R. Kaas, J. Dhaene, M. J. Goovaerts, Upper and lower bounds for sums of random variables, Insurance: Mathematics and Economics 27 (2000) 151–168. doi:10.1016/S0167-6687(00)00060-3.

[5] J. Dhaene, M. Denuit, M. J. Goovaerts, R. Kaas, D. Vyncke, The concept of comonotonicity in actuarial science and finance: Theory, Insurance: Mathematics and Economics 31 (2002) 3–33. doi:10.1016/S0167-6687(02)00134-8.

[6] R. C. Williamson, T. Downs, Probabilistic arithmetic. I. Numerical methods for calculating convolutions and dependency bounds, International Journal of Approximate Reasoning 4 (1990) 89–158. doi:10.1016/0888-613X(90)90022-T.

[7] S. Ghosal, C. P. Ho, W. Wiesemann, A unifying framework for the capacitated vehicle routing problem under risk and ambiguity, Operations Research 72 (2024) 425–443. doi:10.1287/opre.2021.0669.

[8] D. Feillet, P. Dejax, M. Gendreau, C. Gueguen, An exact algorithm for the elementary shortest path problem with resource constraints: Application to some vehicle routing problems, Networks 44 (2004) 216–229. doi:10.1002/net.20033.

[9] S. Irnich, G. Desaulniers, Shortest path problems with resource constraints, in: G. Desaulniers, J. Desrosiers, M. M. Solomon (Eds.), Column Generation, Springer, Boston, MA, 2005, pp. 33–65. doi:10.1007/0-387-25486-2_2.

[10] S. Irnich, Resource extension functions: properties, inversion, and generalization to segments, OR Spectrum 30 (2008) 113–148. doi:10.1007/s00291-007-0083-6.

[11] I. Ioachim, S. Gélinas, F. Soumis, J. Desrosiers, A dynamic programming algorithm for the shortest path problem with time windows and linear node costs, Networks 31 (1998) 193–204. doi:10.1002/(SICI)1097-0037(199805)31:3<193::AID-NET6>3.0.CO;2-A.

[12] R. Baldacci, A. Mingozzi, R. Roberti, New route relaxation and pricing strategies for the vehicle routing problem, Operations Research 59 (2011) 1269–1283. doi:10.1287/opre.1110.0975.

[13] J. M. Buchanan, External diseconomies, corrective taxes, and market structure, American Economic Review 59 (1969) 174–177.

[14] A. H. Barnett, The Pigouvian tax rule under monopoly, American Economic Review 70 (1980) 1037–1041.

[15] D. H. Wolpert, K. Tumer, Optimal payoff functions for members of collectives, Advances in Complex Systems 4 (2001) 265–279. doi:10.1142/S0219525901000188.

[16] J. R. Marden, A. Wierman, Distributed welfare games, Operations Research 61 (2013) 155–168. doi:10.1287/opre.1120.1137.

[17] P. Bergendorff, D. W. Hearn, M. V. Ramana, Congestion toll pricing of traffic networks, in: Network Optimization, volume 450 of *Lecture Notes in Economics and Mathematical Systems*, Springer, Berlin, 1997, pp. 51–71. doi:10.1007/978-3-642-59179-2_4.

[18] D. W. Hearn, M. V. Ramana, Solving congestion toll pricing models, in: Equilibrium and Advanced Transportation Modelling, Springer, Boston, MA, 1998, pp. 109–124. doi:10.1007/978-1-4615-5757-9_6.

[19] R. K. Ahuja, J. B. Orlin, Inverse optimization, Operations Research 49 (2001) 771–783. doi:10.1287/opre.49.5.771.10607.

[20] D. Fotakis, P. G. Spirakis, Cost-balancing tolls for atomic network congestion games, Internet Mathematics 5 (2008) 343–363. doi:10.1080/15427951.2008.10129175.

[21] U. Bhaskar, K. Ligett, L. J. Schulman, C. Swamy, Achieving target equilibria in network routing games without knowing the latency functions, Games and Economic Behavior 118 (2019) 533–569. doi:10.1016/j.geb.2018.02.009.

[22] L. Wang, Cutting plane algorithms for the inverse mixed integer linear programming problem, Operations Research Letters 37 (2009) 114–116. doi:10.1016/j.orl.2008.12.001.

[23] M. Bodur, T. C. Y. Chan, I. Y. Zhu, Inverse mixed integer optimization: Polyhedral insights and trust region methods, INFORMS Journal on Computing 34 (2022) 1471–1488. doi:10.1287/ijoc.2021.1138.

[24] G. Dragotto, R. Scatamacchia, The ZERO regrets algorithm: Optimizing over pure Nash equilibria via integer programming, INFORMS Journal on Computing 35 (2023) 1143–1160. doi:10.1287/ijoc.2022.0282.

[25] D. Monderer, M. Tennenholtz, K-implementation, Journal of Artificial Intelligence Research 21 (2004) 37–62. doi:10.1613/jair.1231.

[26] I. Caragiannis, C. Kaklamanis, P. Kanellopoulos, Taxes for linear atomic congestion games, ACM Transactions on Algorithms 7 (2010) 1–31. doi:10.1145/1868237.1868251.

[27] S. Masiha, S. Elahi, N. Kiyavash, P. Thiran, Zeroth-order Stackelberg control in combinatorial congestion games, arXiv preprint 2602.23277, 2026.

[28] W. H. Sandholm, Evolutionary implementation and congestion pricing, The Review of Economic Studies 69 (2002) 667–689. doi:10.1111/1467-937X.t01-1-00026.

[29] H. Yang, Q. Meng, D.-H. Lee, Trial-and-error implementation of marginal-cost pricing on networks in the absence of demand functions, Transportation Research Part B: Methodological 38 (2004) 477–493. doi:10.1016/S0191-2615(03)00077-8.

[30] D. Han, H. Yang, Congestion pricing in the absence of demand functions, Transportation Research Part E: Logistics and Transportation Review 45 (2009) 159–171. doi:10.1016/j.tre.2008.03.002.

[31] H. Ye, H. Yang, Z. Tan, Learning marginal-cost pricing via a trial-and-error procedure with day-to-day flow dynamics, Transportation Research Part B: Methodological 81 (2015) 794–807. doi:10.1016/j.trb.2015.08.001.

[32] C. Maheshwari, K. Kulkarni, M. Wu, S. Sastry, Inducing social optimality in games via adaptive incentive design, in: 2022 IEEE 61st Conference on Decision and Control (CDC), 2022, pp. 2864–2869. doi:10.1109/CDC51059.2022.9992685, arXiv:2204.05507.

[33] C. Maheshwari, K. Kulkarni, M. Wu, S. Sastry, Adaptive incentive design with learning agents, IEEE Transactions on Automatic Control 71 (2026) 3619–3633. doi:10.1109/TAC.2025.3643351, arXiv:2405.16716.

[34] K. C. Lo, Equilibrium in beliefs under uncertainty, Journal of Economic Theory 71 (1996) 443–484. doi:10.1006/jeth.1996.0129.

[35] M. C. M. Troffaes, Decision making under uncertainty using imprecise probabilities, International Journal of Approximate Reasoning 45 (2007) 17–29. doi:10.1016/j.ijar.2006.06.001.

[36] D. Monderer, L. S. Shapley, Potential games, Games and Economic Behavior 14 (1996) 124–143. doi:10.1006/game.1996.0044.

[37] J. Nash, Non-cooperative games, Annals of Mathematics 54 (1951) 286–295. doi:10.2307/1969529.

[38] W. Hoeffding, Probability inequalities for sums of bounded random variables, Journal of the American Statistical Association 58 (1963) 13–30. doi:10.1080/01621459.1963.10500830.

[39] K. Liu, N. Oudjane, C. Wan, Approximate Nash equilibria in large nonconvex aggregative games, Mathematics of Operations Research 48 (2023) 1791–1809. doi:10.1287/moor.2022.1321.

[40] V. Chvátal, On the computational complexity of finding a kernel, Technical Report CRM-300, Centre de Recherches Mathématiques, Université de Montréal, 1973.

[41] A. S. Fraenkel, Planar kernel and Grundy with $d \leq 3$, $d_{out} \leq 2$, $d_{in} \leq 2$ are NP-complete, Discrete Applied Mathematics 3 (1981) 257–262. doi:10.1016/0166-218X(81)90003-2.

[42] K. R. Apt, S. Simon, D. Wojtczak, Coordination games on weighted directed graphs, Mathematics of Operations Research 47 (2022) 995–1025. doi:10.1287/moor.2021.1159.

[43] M. Carvalho, A. Lodi, J. P. Pedroso, Computing equilibria for integer programming games, European Journal of Operational Research 303 (2022) 1057–1070. doi:10.1016/j.ejor.2022.03.048.

[44] X. Chen, X. Deng, S.-H. Teng, Settling the complexity of computing two-player Nash equilibria, Journal of the ACM 56 (2009) 1–57. doi:10.1145/1516512.1516516.

[45] C. Daskalakis, P. W. Goldberg, C. H. Papadimitriou, The complexity of computing a Nash equilibrium, SIAM Journal on Computing 39 (2009) 195–259. doi:10.1137/070699652.

[46] H. B. McMahan, G. J. Gordon, A. Blum, Planning in the presence of cost functions controlled by an adversary, in: Proceedings of the 20th International Conference on Machine Learning (ICML 2003), 2003, pp. 536–543.

[47] J. P. Aubin, I. Ekeland, Estimates of the duality gap in nonconvex optimization, Mathematics of Operations Research 1 (1976) 225–245. doi:10.1287/moor.1.3.225.

[48] M. Udell, S. Boyd, Bounding duality gap for separable problems with linear constraints, Computational Optimization and Applications 64 (2016) 355–378. doi:10.1007/s10589-015-9819-4.

[49] J. F. Bonnans, K. Liu, N. Oudjane, L. Pfeiffer, C. Wan, Large-scale nonconvex optimization: Randomization, gap estimation, and numerical resolution, SIAM Journal on Optimization 33 (2023) 3083–3113. doi:10.1137/22M1488892.

[50] M. Haklay, P. Weber, OpenStreetMap: User-generated street maps, IEEE Pervasive Computing 7 (2008) 12–18. doi:10.1109/MPRV.2008.80.

[51] M. Schiavina, S. Freire, K. MacManus, GHS-POP R2023A – GHS population grid multitemporal (1975–2030), European Commission, Joint Research Centre (JRC), dataset, 2023. doi:10.2905/2FF68A52-5B5B-4A22-8F40-C41DA8332CFE.

[52] TomTom, TomTom traffic index 2025, https://www.tomtom.com/traffic-index/, accessed 23 October 2026, 2026.

[53] Bureau of Public Roads, Traffic Assignment Manual for Application with a Large, High Speed Computer, Technical Report, U.S. Department of Commerce, Urban Planning Division, Washington, DC, 1964.

[54] S. Dabia, S. Ropke, T. van Woensel, T. De Kok, Branch and price for the time-dependent vehicle routing problem with time windows, Transportation Science 47 (2013) 380–396. doi:10.1287/trsc.1120.0445.

[55] R. M. Starr, Quasi-equilibria in markets with non-convex preferences, Econometrica 37 (1969) 25–38. doi:10.2307/1909201.

[56] R. T. Rockafellar, Convex Analysis, Princeton University Press, Princeton, NJ, 1970. doi:10.1515/9781400873173.

[57] Y. Nesterov, Introductory Lectures on Convex Optimization: A Basic Course, Kluwer Academic Publishers, Boston, MA, 2004. doi:10.1007/978-1-4419-8853-9.

[58] O. Briant, C. Lemaréchal, P. Meurdesoif, S. Michel, N. Perrot, F. Vanderbeck, Comparison of bundle and classical column generation, Mathematical Programming 113 (2008) 299–344. doi:10.1007/s10107-006-0079-z.

[59] W. Powell, A. Ruszczyński, H. Topaloglu, Learning algorithms for separable approximations of discrete stochastic optimization problems, Mathematics of Operations Research 29 (2004) 814–836. doi:10.1287/moor.1040.0107.

[60] R. Paes Leme, S. C.-w. Wong, Computing Walrasian equilibria: Fast algorithms and structural properties, Mathematical Programming 179 (2020) 343–384. doi:10.1007/s10107-018-1334-9.

[61] A. Duguet, T. Harks, M. Schmidt, J. Schwarz, Minimal regret Walras equilibria for combinatorial markets, arXiv preprint 2511.09021, 2025.