Highlights

TRACER-Med: Harmonizing Privacy, Fairness, and Uncertainty in Clinical Deep Learning

Quang-Vinh Dang

- Integrates DP, subgroup equity, uncertainty, and XAI in clinical AI.
- Subgroup-calibrated DP-SGD bounds sensitivity by $C_{\max}$ under Rényi DP.
- Private group-conditional conformal prediction provides coverage equity.
- Dual-channel Dirichlet attribution separates disease risk from clinical doubt.
- Evaluated across cardiovascular, metabolic, and acute inpatient cohorts.

TRACER-Med: Harmonizing Privacy, Fairness, and Uncertainty in Clinical Deep Learning

Quang-Vinh Dang[a,*]

[a]*British University Vietnam, Hung Yen, Vietnam*

Abstract

Background: Clinical deep learning deployment requires patient data privacy, demographic fairness, calibrated uncertainty, and faithful explainability. Standard Differentially Private Stochastic Gradient Descent (DP-SGD) often degrades minority performance and miscalibrates probabilities, inducing a Privacy–Fairness–Uncertainty Trilemma.
Methods: We propose **TRACER-Med**, an integrated architecture combining: (i) an Evidential Dirichlet network parameterizing second-order beliefs to separate predictive entropy from epistemic vacuity; (ii) Subgroup-Calibrated DP-SGD (SC-DP-SGD) with group-conditioned clipping and dynamic sensitivity reweighting calibrated to worst-case sensitivity $C_{\max}$ under Rényi Differential Privacy; (iii) Differentially Private Group-Conditional Conformal Prediction (DP-GCCP) with split-level privacy accounting; and (iv) Dual-Channel Dirichlet Attribution (DCDA) path-integrating diagnostic risk evidence and epistemic doubt.
Results: Evaluated across Statlog Heart ($N = 270$), Pima Diabetes ($N = 768$), and a Simulated Inpatient EHR cohort ($N = 10{,}000$), TRACER-Med reduces Equalized Odds disparity to 0.083 ± 0.02 (a 42% reduction vs DP-EDL) and Statistical Parity disparity to 0.232 ± 0.04 (a 27% reduction) on Statlog Heart, maintaining competitive discrimination (ROC-AUC 0.863 ± 0.05 on Heart, 0.826 ± 0.02 on Pima). Subgroup conformal coverage gaps are controlled at 0.045 ± 0.05 (Heart) and 0.058 ± 0.04 (Pima) at 90% target coverage, with empirical membership inference resistance (MIA AUC 0.50–0.55) under formal Differential Privacy accounting ($\epsilon = 15.36$ on Heart, 6.13 on Pima, and 2.49 on EHR at $\delta = 10^{-5}$).
Conclusions: TRACER-Med establishes an auditable blueprint for deploying equitable, well-calibrated, and privacy-preserving clinical AI.

Keywords: Trustworthy Machine Learning, Differential Privacy, Algorithmic Fairness, Evidential Deep Learning, Conformal Prediction, Explainable AI

1. Introduction

Artificial intelligence and deep neural networks have demonstrated diagnostic accuracy comparable to domain specialists across medical specialties, including acute mortality triage,

*Corresponding author.
Email address: vinh.dq4@buv.edu.vn (Quang-Vinh Dang)

cardiovascular risk scoring, digital pathology, and medical imaging [1, 2, 3]. Despite these achievements, translation into bedside clinical workflows remains limited. Clinical oversight committees, practitioners, and international regulatory authorities (including the U.S. Food and Drug Administration, HHS Office for Civil Rights, and European Union AI Act) recognize that predictive discrimination alone is insufficient for clinical adoption. Safe deployment requires adherence to four essential pillars of *Trustworthy Medical AI*:

1. **Provable Patient Privacy:** Models trained on confidential Electronic Health Records (EHRs) and patient diagnostic registries must provide rigorous bounds against re-identification and attribute reconstruction attacks, formalized via Differential Privacy (DP) [4, 5].
2. **Subgroup Algorithmic Fairness:** Prognostic and diagnostic models must not exacerbate healthcare disparities across demographic cohorts defined by biological sex, age, race, or socioeconomic indicators [6, 7, 8].
3. **Calibrated Uncertainty Quantification:** High-stakes medicine requires systems capable of expressing their own limitations. Algorithms must output well-calibrated probabilities, distinguish irreducible observational noise (aleatoric uncertainty) from missing diagnostic evidence (epistemic vacuity), and provide distribution-free finite-sample coverage guarantees [9, 10, 11].
4. **Faithful and Actionable Explainability:** Clinicians cannot safely act on uninterpretable decisions. Models must provide transparent feature attributions that explain not only why an alert was triggered, but also which physiological parameters are driving clinical hesitation [12, 13].

1.1. The Privacy–Fairness–Uncertainty Trilemma in Healthcare

Historically, these four dimensions have been investigated in separate subfields. When naively combined in deep neural networks, they exhibit significant friction, giving rise to what we characterize as the *Privacy–Fairness–Uncertainty Trilemma*:

- **Differential Privacy Disproportionately Impacts Minority Accuracy:** In standard Differentially Private Stochastic Gradient Descent (DP-SGD) [4], uniform per-sample gradient clipping and isotropic Gaussian noise addition produce an asymmetric equity penalty. Because underrepresented patient groups often exhibit distinct physiological presentations, their gradient norms and directions differ from the majority cohort. Uniform clipping disproportionately truncates minority gradients, while the shared noise floor degrades the effective Signal-to-Noise Ratio (SNR) of minority updates, widening diagnostic performance gaps [14, 15, 16, 17].

- **DP Perturbations Induce Subgroup Miscalibration:** Softmax-parameterized neural networks trained with DP-SGD often suffer from substantial probability distortion and miscalibration. Furthermore, this calibration degradation is unevenly distributed across demographic cohorts, leading to biased risk stratification in clinical triage [18, 19].

- **Standard Conformal Prediction Fails Conditional Subgroup Guarantees:** While split conformal prediction guarantees marginal coverage $1 - \alpha_{\text{conf}}$ over the aggregate population [20, 11], it does not guarantee subgroup-conditional validity. When conformal quantiles are estimated under differential privacy constraints, minority cohorts frequently experience coverage deficits, receiving prediction sets that fail to cover the true clinical condition at the nominal safety level [21, 22].

- **Gradient Perturbations Degrade Feature Attributions:** Post-hoc gradient attribution techniques (e.g., Integrated Gradients) depend on model input gradients. Under differential privacy, network weight constraints and regularization induce significant shifts in decision boundaries, often resulting in noisy, unstable attributions that fail faithfulness evaluations and obscure the true clinical rationale.

1.2. Our Contributions

To address these challenges within a cohesive architectural design, we propose **TRACER-Med** (**TR**ustworthy, **A**daptive, **C**onformalized, **E**vidential, and Differentially P**R**ivate Deep Learning for **Med**icine). Our primary contributions are:

1. **Systematic Dissection of Inter-Pillar Trade-offs:** We provide an analytical and empirical formulation of the Privacy–Fairness–Uncertainty Trilemma in clinical machine learning, examining why isolated privacy, fairness, and uncertainty interventions induce trade-offs when deployed on demographically imbalanced patient registries.
2. **Evidential Dirichlet Deep Learning Formulation:** We formulate a second-order Evidential Deep Learning (EDL) backbone that maps clinical patient representations to Dirichlet concentration parameters. This analytically separates aleatoric entropy from epistemic vacuity, providing calibrated uncertainty estimates under private optimization.
3. **Subgroup-Calibrated DP Optimization (SC-DP-SGD):** We derive an adaptive, group-conditioned clipping and dynamic sensitivity reweighting mechanism that bounds global ℓ_2 sensitivity by $C_{\max}$ across protected demographic groups. In Theorem 1, we prove that calibrating noise to $C_{\max}$ using fixed population demographic priors provides provable Rényi Differential Privacy guarantees, while SC-DP-SGD empirically reduces demographic disparity by 27% (Δ_{SP}) and 42% (Δ_{EO}) compared to DP-EDL on cardiovascular diagnostics.
4. **Differentially Private Group-Conditional Conformal Prediction (DP-GCCP):** We develop a private conformal calibration framework incorporating group-conditional non-conformity partitions, finite-sample concentration adjustments, and private quantile estimation on disjoint calibration data with rigorous split-level privacy accounting.
5. **Dual-Channel Dirichlet Attribution (DCDA):** Leveraging the closed-form structure of Dirichlet evidence, we introduce a dual-channel path-integrated attribution formulation. Channel A attributes input features driving positive diagnostic risk, while Channel B isolates features driving epistemic uncertainty and clinical doubt.
6. **Comprehensive Empirical Validation:** We evaluate TRACER-Med across three clinical benchmarks: Statlog Heart Disease ($N = 270$), Pima Indians Diabetes

($N = 768$), and a simulated 10,000-patient Inpatient EHR Cohort (parameterized from MIMIC-IV and eICU distributions). We demonstrate competitive diagnostic discrimination (ROC-AUC), calibrated probabilities (ECE), controlled membership inference risk (MIA AUC), equitable conformal coverage, and faithful dual-channel attributions.

2. Related Work

2.1. Differential Privacy in Biomedical Machine Learning

Differential Privacy (DP) [4] provides a formal framework for bounding the privacy risk of individual patient records during model training. In deep learning, DP-SGD restricts per-sample gradient ℓ_2 norms by clipping to a threshold C, adds Gaussian perturbation $\mathcal{N}(0, \sigma^2 C^2 \mathbf{I})$, and accumulates privacy loss via Rényi Differential Privacy (RDP) [5, 23] or Gaussian Differential Privacy (GDP) [24].

However, Bagdasaryan et al. [14] and Suriyakumar et al. [15] demonstrated that standard DP-SGD acts as an equity penalty: uniform gradient clipping truncates underrepresented patient examples more severely, widening demographic error gaps. Recent algorithmic proposals have explored fair DP in generic classification settings: Xu et al. [17] proposed DPSGD-F, adapting group clipping thresholds to equalize demographic parity; Tran et al. [16] formulated a Lagrangian dual optimization framework for fair private ERM; Esipova et al. [25] investigated gradient angle alignment; and Wang et al. [26] audited intersectional disparities in healthcare. Crucially, recent 2025–2026 investigations by Chen et al. [27] demonstrated that private clinical models exacerbate intersectional disparities across marginalized patient groups, while Tran et al. [28] developed fair private gradient descent routines for heterogeneous clinical cohorts. However, existing fair DP algorithms focus almost exclusively on point-prediction accuracy and parity metrics. They do not address epistemic uncertainty quantification under evidential priors, distribution-free conformal calibration, or dual-channel explanations. TRACER-Med bridges this divide by unifying subgroup-calibrated clipping with Dirichlet second-order beliefs and private conformal prediction.

2.2. Algorithmic Fairness in Clinical Healthcare

Algorithmic bias in biomedical models is well-documented. Obermeyer et al. [6] showed that commercial risk algorithms systematically prioritized healthier White patients over sicker Black patients for high-risk care management programs. Seyyed-Kalantari et al. [8] identified that convolutional networks for chest radiographs exhibited substantial underdiagnosis bias in female and underserved populations. Typical fairness interventions comprise pre-processing, in-processing (e.g., representation debiasing via optimal transport and Sliced Wasserstein Distance [29]), and post-processing (e.g., equalized odds thresholding [30]). However, the interaction of these interventions with differentially private gradient sanitization requires careful design to prevent noise from neutralizing fairness gains.

2.3. Evidential Uncertainty Quantification and Conformal Prediction

Standard deep neural networks with softmax parameterizations tend to output overconfident probabilities on out-of-distribution (OOD) or ambiguous clinical profiles [31]. Evidential Deep Learning (EDL) [10] parameterizes a Dirichlet distribution over categorical probability vectors, allowing analytical decomposition into aleatoric entropy and epistemic vacuity. While recent studies have examined theoretical nuances of loss-minimized evidential learning [32, 33], evidential parameterizations offer natural closed-form analytical properties suitable for path-integrated gradient explanations.

Conformal Prediction (CP) [20, 11] constructs distribution-free, finite-sample prediction sets $\mathcal{C}(X)$ that satisfy a marginal coverage guarantee $\mathbb{P}(Y \in \mathcal{C}(X)) \geq 1 - \alpha_{\text{conf}}$. To enforce equity across demographic subgroups, Mondrian conformal prediction [20] and group-conditional conformal prediction [21, 22] compute group-specific non-conformity quantiles. Recent clinical advances by Angelopoulos et al. [34] formulated conformal risk control for decision support under distribution shift, while Sesia et al. [35] established conformalized clinical risk assessments resilient to demographic and temporal shifts. Extending conformal prediction to differentially private settings requires private quantile mechanisms [36, 37] to avoid membership leakage while preserving bounded set size efficiency.

2.4. Explainable AI and Attribution Robustness

Attribution methods such as Integrated Gradients (IG) [12] provide axiomatic guarantees (including completeness and implementation invariance) for feature attribution. Most existing attribution frameworks are single-channel, explaining only the predicted disease class. However, in clinical decision support, identifying features that contribute to epistemic uncertainty is equally essential [38, 39], as underscored by Watson et al. [40] who introduced path-integrated attributions of predictive doubt, and Hüllermeier et al. [41] who formalized the clinical distinction between aleatoric and epistemic uncertainty. Furthermore, assessing explanation fidelity (e.g., via Infidelity metrics [42]) under differential privacy constraints is crucial to ensure that clinicians receive reliable diagnostic explanations.

3. Problem Formulation and Notation

3.1. Formal Setting and Notation

Let $\mathcal{D} = \{(\mathbf{x}_i, y_i, a_i)\}_{i=1}^{N}$ denote a clinical dataset where $\mathbf{x}_i \in \mathbb{R}^D$ represents patient clinical features, $y_i \in \{0, 1, \ldots, K-1\}$ denotes the clinical diagnosis or risk outcome, and $a_i \in \mathcal{A} = \{0, 1\}$ represents a binary protected demographic attribute. Let $p_a = \mathbb{P}(A = a)$ denote the known population prior demographic prevalence obtained from public health registries or census references, ensuring that p_a is strictly independent of the private training sample and introduces no auxiliary privacy leakage.

Differential Privacy: We adopt standard add/remove record adjacency [43]: two datasets $\mathcal{D}, \mathcal{D}'$ are neighboring ($\mathcal{D} \simeq \mathcal{D}'$) if one dataset is obtained by adding or removing a single patient record $z = (\mathbf{x}, y, a)$: $\mathcal{D}' = \mathcal{D} \cup \{z\}$ or $\mathcal{D}' = \mathcal{D} \setminus \{z\}$. A randomized mechanism

$\mathcal{M}$ satisfies (ϵ, δ)-Differential Privacy [4] if for all neighboring datasets $\mathcal{D} \simeq \mathcal{D}'$, and for all measurable sets of outputs $\mathcal{S} \subseteq \text{Range}(\mathcal{M})$:

$$\mathbb{P}(\mathcal{M}(\mathcal{D}) \in \mathcal{S}) \leq e^{\epsilon}\mathbb{P}(\mathcal{M}(\mathcal{D}') \in \mathcal{S}) + \delta \tag{1}$$

Under Rényi Differential Privacy (RDP) of order $\lambda_{\text{RDP}} > 1$ [5], the Rényi divergence between $\mathcal{M}(\mathcal{D})$ and $\mathcal{M}(\mathcal{D}')$ satisfies $D_{\lambda_{\text{RDP}}}(\mathcal{M}(\mathcal{D}) \parallel \mathcal{M}(\mathcal{D}')) \leq \epsilon(\lambda_{\text{RDP}})$.

Algorithmic Fairness Criteria: We evaluate two standard clinical equity metrics:

1. *Statistical Parity Disparity (Δ_{SP}):*

$$\Delta_{\text{SP}} = |\mathbb{P}(\hat{Y} = 1 \mid A = 0) - \mathbb{P}(\hat{Y} = 1 \mid A = 1)| \tag{2}$$

   We designate this demographic parity metric as Δ_{SP} throughout this paper to avoid notational confusion with Differential Privacy (DP).

2. *Equalized Odds Disparity (Δ_{EO}):*

$$\Delta_{\text{EO}} = \frac{1}{2}\left(|\text{FPR}_0 - \text{FPR}_1| + |\text{TPR}_0 - \text{TPR}_1|\right) \tag{3}$$

   where FPR_a and TPR_a denote the False Positive Rate and True Positive Rate within demographic cohort $a \in \{0, 1\}$. Note that Δ_{EO} represents the average absolute error gap across groups, distinguished from Hardt et al.'s maximum form [30].

In clinical medicine, differing baseline disease prevalence between demographic cohorts (e.g., coronary disease prevalence across biological sex) induces fundamental mathematical trade-offs: as established by Barocas, Hardt, and Narayanan [30], Statistical Parity (independence) and Equalized Odds (separation) are mutually exclusive under unequal base rates ($\mathbb{P}(Y = 1 \mid A = 0) \neq \mathbb{P}(Y = 1 \mid A = 1)$) unless predictions are perfectly accurate. Furthermore, Chouldechova [44] and Kleinberg et al. [45] demonstrated that predictive parity (within-group calibration) and error-rate balance cannot simultaneously hold under disparate prevalence. In high-stakes medicine, artificially forcing Statistical Parity ($\Delta_{\text{SP}} \to 0$) forces unequal false positive and false negative rates across groups, potentially subjecting low-prevalence cohorts to over-diagnosis or high-prevalence cohorts to under-treatment. Consequently, Equalized Odds (Δ_{EO}) serves as our primary clinical equity criterion, ensuring that conditional on true disease status, diagnostic sensitivity and specificity are equitable across demographic groups.

3.2. Identified Research Gaps

Our analysis identifies four core technical gaps:

- **Gap 1 (Disparate Impact from Uniform Gradient Clipping):** In demographically imbalanced clinical cohorts ($p_0 < p_1$), minority patients often present distinct clinical presentations. Under standard DP-SGD with uniform clipping threshold C, gradients from minority patients are clipped more severely, degrading their gradient representation relative to the majority cohort [14, 15].

- **Gap 2 (Uncertainty Conflation under Differential Privacy):** Cross-entropy classifiers parameterized by standard softmax output point estimates that conflate data variability with epistemic vacuity (deficiency of clinical evidence). Under DP gradient noise, probability estimates miscalibrate, yielding severe Expected Calibration Error (ECE) [18].
- **Gap 3 (Subgroup Coverage Deficits in Conformal Prediction):** Standard marginal conformal prediction guarantees aggregate population coverage $1 - \alpha_{\text{conf}}$, but permits substantial coverage gaps between demographic groups: $\mathbb{P}(Y \in \mathcal{C}(X) \mid A = 0) \ll 1 - \alpha_{\text{conf}}$. Estimating group quantiles privately without excessive set size inflation is essential for clinical equity [21, 36].
- **Gap 4 (Single-Channel Explanations):** Existing feature attribution methods provide a single attribution score for predicted classes, failing to inform clinicians which physiological features are driving model uncertainty and clinical doubt [38].

4. The TRACER-Med Architecture

TRACER-Med addresses these gaps through four integrated modules, summarized in Algorithm 1.

4.1. Evidential Dirichlet Deep Learning Formulation

Rather than predicting a single categorical probability vector via softmax, TRACER-Med parameterizes a second-order Dirichlet distribution over multinomial probabilities. The network encoder and non-negative evidence head map patient features $\mathbf{x} \in \mathbb{R}^D$ to evidence vector $\mathbf{e} = f_{\boldsymbol{\theta}}(\mathbf{x}) = \text{softplus}(\mathbf{z}) \geq \mathbf{0}$, defining Dirichlet concentration parameters:

$$\alpha_k = e_k + 1, \quad S = \sum_{k=1}^{K} \alpha_k \tag{4}$$

where S is the total Dirichlet evidence strength. The expected categorical probability for class k is given by:

$$\hat{p}_k = \frac{\alpha_k}{S} \tag{5}$$

Subjective Logic parameterizes the epistemic uncertainty (vacuity of evidence) as:

$$u = \frac{K}{S} \in (0, 1] \tag{6}$$

When the network observes novel or anomalous clinical features, $\mathbf{e} \to \mathbf{0}$, yielding $\alpha_k \to 1$, $S \to K$, and epistemic uncertainty $u \to 1$. Conversely, abundant clinical evidence drives $S \gg K$, yielding $u \to 0$. In parallel, total predictive uncertainty is captured by the Shannon entropy of the expected probabilities: $\mathcal{H}(\hat{\mathbf{p}}) = -\sum_{k=1}^{K} \hat{p}_k \ln \hat{p}_k$. We note that while Dirichlet parameterizations provide a closed-form analytical proxy for epistemic doubt, loss-minimized uncertainty estimates require careful validation [32, 33].

The Evidential loss combines the Bayesian expected cross-entropy loss under the Dirichlet distribution with a Kullback–Leibler (KL) divergence regularizer that penalizes misleading evidence for non-target classes:

$$\mathcal{L}_{\mathrm{EDL}}(\boldsymbol{\theta}) = \sum_{i=1}^{|\mathcal{B}|} \left[\sum_{k=1}^{K} y_{ik} \left(\psi(S_i) - \psi(\alpha_{ik}) \right) + \lambda_{\mathrm{reg}} \mathcal{D}_{\mathrm{KL}} \left(\mathrm{Dir}(\tilde{\boldsymbol{\alpha}}_i) \parallel \mathrm{Dir}(\mathbf{1}) \right) \right] \tag{7}$$

where $\psi(\cdot)$ denotes the digamma function, and $\tilde{\boldsymbol{\alpha}}_i = \mathbf{y}_i + (1 - \mathbf{y}_i) \odot \boldsymbol{\alpha}_i$.

4.2. Subgroup-Calibrated Differentially Private Optimization (SC-DP-SGD)

To prevent differential privacy from exacerbating demographic disparity, we introduce the *Subgroup-Calibrated DP-SGD* (SC-DP-SGD) optimizer.

Population-Calibrated Weights: To prevent batch demographic counts from introducing sensitivity variations, we compute scaling weights from population reference priors $p_a = \mathbb{P}(A = a)$:

$$w_a = \left(\frac{1}{2p_a} \right)^{\gamma}, \quad \gamma = 0.5 \tag{8}$$

normalized to satisfy $\sum_{b \in \mathcal{A}} p_b \tilde{w}_b^2 = 1$:

$$\tilde{w}_a = \frac{w_a}{\sqrt{\sum_{b \in \mathcal{A}} p_b w_b^2}} \tag{9}$$

For each patient example i belonging to demographic group $a_i \in \{0, 1\}$, the per-sample gradient $g_i = \nabla_{\boldsymbol{\theta}} \ell_i$ is clipped using the group-conditioned bound $C_{a_i} = C \cdot \tilde{w}_{a_i}$:

$$\bar{g}_i = g_i \cdot \min \left(1, \frac{C \cdot \tilde{w}_{a_i}}{\|g_i\|_2} \right) \tag{10}$$

Under add/remove record adjacency, adding or removing a record $z = (\mathbf{x}, y, a)$ changes the batch gradient sum by at most one clipped vector $\|\bar{g}_z\|_2 \leq C_a \leq C_{\max}$, where:

$$C_{\max} = C \max_{a \in \mathcal{A}} \tilde{w}_a \tag{11}$$

The aggregated gradient is sanitized by injecting Gaussian noise scaled to the worst-case sensitivity $C_{\max}$:

$$\tilde{G} = \frac{1}{|\mathcal{B}|} \left(\sum_{i \in \mathcal{B}} \bar{g}_i + \mathcal{N} \left(\mathbf{0}, \sigma^2 C_{\max}^2 \mathbf{I} \right) \right) \tag{12}$$

Theorem 1 (Privacy Guarantee of SC-DP-SGD)**.** *Let $\mathcal{M}$ denote the SC-DP-SGD algorithm operating on dataset $\mathcal{D}$ under add/remove record adjacency, with training sample size N_{tr}, Poisson subsampling ratio $q = |\mathcal{B}|/N_{tr}$, total training steps $T = E \cdot \lceil N_{tr}/|\mathcal{B}| \rceil$, base clipping bound C, group clipping thresholds $C_a = C\tilde{w}_a$, and noise standard deviation $\sigma C_{\max}$, where $C_{\max} = C \max_a \tilde{w}_a$.*

Then, $\mathcal{M}$ satisfies $(\lambda_{RDP}, \epsilon(\lambda_{RDP}))$-Rényi Differential Privacy for integer orders $\lambda_{RDP} \geq 2$ under the Poisson-subsampled Gaussian mechanism [23, 46]. Consequently, $\mathcal{M}$ satisfies standard (ϵ, δ)-Differential Privacy for target $\delta > 0$, with:

$$\epsilon = \min_{\lambda_{RDP} \geq 2} \left(\epsilon(\lambda_{RDP}) + \frac{\ln(1/\delta)}{\lambda_{RDP} - 1} \right) \tag{13}$$

Under our training configurations ($\sigma = 0.8, C = 1.0, \delta = 10^{-5}$), composition via the exact numerical RDP accountant yields:

- Statlog Heart: $N_{tr} = 180, |\mathcal{B}| = 32, q = 0.178, T = 113$ *steps ($E = 20$ epochs)* $\implies \epsilon = 15.36$.
- Pima Diabetes: $N_{tr} = 512, |\mathcal{B}| = 32, q = 0.0625, T = 320$ *steps ($E = 20$ epochs)* $\implies \epsilon = 6.13$.
- Simulated Inpatient EHR: $N_{tr} = 6{,}667, |\mathcal{B}| = 128, q = 0.0192, T = 625$ *steps ($E = 12$ epochs)* $\implies \epsilon = 2.49$.

Proof. Let $\mathcal{D}$ and $\mathcal{D}'$ be neighboring datasets differing by at most one patient record under add/remove adjacency: $\mathcal{D}' = \mathcal{D} \cup \{z\}$ or $\mathcal{D}' = \mathcal{D} \setminus \{z\}$ for some patient record $z = (\mathbf{x}, y, a)$. In SC-DP-SGD, adding or removing record z alters at most one clipped gradient vector $\bar{g}_z$ in the mini-batch sum. Since the clipping rule enforces $\|\bar{g}_z\|_2 \leq C_a \leq C_{\max}$, the global ℓ_2 sensitivity of the batch sum is strictly bounded by:

$$\Delta_2 = \sup_{\mathcal{D} \simeq \mathcal{D}'} \left\| \sum_{i \in \mathcal{B}} \bar{g}_i(\mathcal{D}) - \sum_{i \in \mathcal{B}'} \bar{g}_i(\mathcal{D}') \right\|_2 \leq C_{\max} \tag{14}$$

Because the injected Gaussian noise has covariance $\sigma^2 C_{\max}^2 \mathbf{I}$, the effective noise multiplier relative to the global ℓ_2 sensitivity is $\frac{\sigma C_{\max}}{\Delta_2} \geq \frac{\sigma C_{\max}}{C_{\max}} = \sigma$. Under Poisson subsampling with ratio $q = |\mathcal{B}|/N_{\text{tr}}$, the privacy loss per step is bounded by the exact Rényi divergence of the subsampled Gaussian mechanism $\epsilon(\lambda_{\text{RDP}})$ [23, 46]. By linear RDP composition over T steps, the cumulative divergence satisfies $T \cdot \epsilon(\lambda_{\text{RDP}})$. Conversion to standard (ϵ, δ)-DP follows directly by optimizing over integer orders $\lambda_{\text{RDP}} \in \{2, 3, \ldots, 64\}$. □

4.3. Latent Sliced Wasserstein Demographic Alignment

To regularize representation fairness during non-private training and feature extraction, TRACER-Med applies Sliced Wasserstein Distance (SWD) [29] alignment to penultimate latent representations $\mathbf{z} = h_{\boldsymbol{\theta}}(\mathbf{x}) \in \mathbb{R}^d$:

$$\mathcal{L}_{\text{fair}} = \mathcal{SW}(P(\mathbf{z} \mid A = 0), P(\mathbf{z} \mid A = 1)) = \frac{1}{M} \sum_{m=1}^{M} \int_0^1 \left| F^{-1}_{0,\boldsymbol{\omega}_m}(t) - F^{-1}_{1,\boldsymbol{\omega}_m}(t) \right| dt \tag{15}$$

where $\boldsymbol{\omega}_m \sim \text{Unif}(\mathbb{S}^{d-1})$ are $M = 50$ random unit projection vectors, and $F^{-1}_{a,\boldsymbol{\omega}_m}$ represents the 1D quantile function of group a's projected representations.

Decoupled DP Sensitivity Guarantee: Importantly, during private optimization with SC-DP-SGD, only the per-sample evidential Dirichlet loss $\mathcal{L}_{\text{EDL}}$ is backpropagated through the network. Because standard SWD couples samples across mini-batches—which would invalidate per-sample gradient independence—SWD is applied during representation pre-training on public/unlabelled reference data, or evaluated within non-private baselines (Fair-ERM). This architectural decoupling guarantees that no cross-sample gradient coupling occurs during private steps, ensuring that Theorem 1's per-sample sensitivity bound and privacy guarantee strictly govern all deployed network parameters.

4.4. Differentially Private Group-Conditional Conformal Prediction (DP-GCCP)

Following model convergence, TRACER-Med calibrates conformal prediction sets on a held-out calibration split $\mathcal{D}_{\text{cal}} = \{(\mathbf{x}_j, y_j, a_j)\}_{j=1}^{N_{\text{cal}}}$, disjoint from training data $\mathcal{D}_{\text{tr}}$. Non-conformity scores are computed as the residual probability of the true label: $s_j = 1 - \hat{p}_{y_j}(\mathbf{x}_j)$.

To establish distribution-free finite-sample coverage without leaking patient calibration scores, we deploy a private quantile mechanism on the empirical Cumulative Distribution Function (CDF) [36, 37]. For each protected demographic group $a \in \mathcal{A}$ with calibration subset $\mathcal{S}_a = \{s_j : a_j = a\}$, the empirical CDF $F_{\mathcal{S}_a}(t) = \frac{1}{|\mathcal{S}_a|} \sum_{s \in \mathcal{S}_a} \mathbb{I}(s \leq t)$ has global ℓ_1 sensitivity $\Delta_{\text{CDF}} = \frac{1}{|\mathcal{S}_a|}$ under add/remove record adjacency. The private threshold $\hat{q}_a$ is computed by perturbing the target quantile index with Laplace noise and a finite-sample concentration buffer τ_a:

$$\hat{q}_a = \inf \left\{ t \in [0,1] : F_{\mathcal{S}_a}(t) \geq 1 - \alpha_{\text{conf}} + \tau_a + \frac{\ln(4/\beta)}{|\mathcal{S}_a| \cdot \epsilon_{\text{conf}}} \right\} \tag{16}$$

where $\tau_a = \sqrt{\frac{\ln(4/\beta)}{2|\mathcal{S}_a|}}$ is the Dvoretzky–Kiefer–Wolfowitz (DKW) finite-sample concentration adjustment at confidence $1 - \beta/2$, and $\frac{\ln(4/\beta)}{|\mathcal{S}_a|\epsilon_{\text{conf}}}$ accounts for the Laplace perturbation tail bound.

Proposition 2 (Subgroup-Conditional Coverage Validity)**.** *Assuming exchangeability of calibration and test non-conformity scores within each demographic cohort $a \in \mathcal{A}$, the private prediction sets $\mathcal{C}(\mathbf{x}) = \{k : 1 - \hat{p}_k(\mathbf{x}) \leq \hat{q}_a\}$ satisfy finite-sample subgroup coverage:*

$$\mathbb{P}(Y \in \mathcal{C}(X) \mid A = a) \geq 1 - \alpha_{conf} \tag{17}$$

with probability at least $1 - \beta$ over the randomness of the calibration sample and the Laplace mechanism.

Proof. Let F_a denote the true population cumulative distribution function of non-conformity scores for demographic group a, and let $F_{\mathcal{S}_a}$ denote the empirical distribution on calibration split $\mathcal{S}_a$. By the Dvoretzky–Kiefer–Wolfowitz (DKW) inequality, the uniform empirical deviation satisfies:

$$\mathbb{P}\left(\sup_{t \in [0,1]} |F_{\mathcal{S}_a}(t) - F_a(t)| > \tau_a \right) \leq 2\exp(-2|\mathcal{S}_a|\tau_a^2) = \frac{\beta}{2} \tag{18}$$

for $\tau_a = \sqrt{\frac{\ln(4/\beta)}{2|\mathcal{S}_a|}}$. Furthermore, for the Laplace perturbation $\xi_a \sim \text{Lap}\left(\frac{1}{|\mathcal{S}_a|\epsilon_{\text{conf}}}\right)$, the one-sided tail probability satisfies $\mathbb{P}\left(\xi_a < -\frac{\ln(4/\beta)}{|\mathcal{S}_a|\epsilon_{\text{conf}}}\right) \leq \frac{1}{2}\exp(-\ln(4/\beta)) = \frac{\beta}{8} \leq \frac{\beta}{2}$. Applying a union bound over sampling concentration ($\beta/2$) and Laplace mechanism noise ($\beta/2$) ensures that with probability at least $1-\beta$, $F_a(\hat{q}_a) \geq 1 - \alpha_{\text{conf}}$, guaranteeing valid subgroup-conditional coverage $\mathbb{P}(Y \in \mathcal{C}(X) \mid A = a) \geq 1 - \alpha_{\text{conf}}$. □

Calibration Sample Size and Set Informative-ness: We note an important practical relationship between calibration sample size $|\mathcal{S}_a|$ and conformal set efficiency. In small minority cohorts (such as Statlog Heart with $|\mathcal{S}_0| \approx 15-20$), the conservative DKW buffer τ'_a pushes the threshold towards 1, yielding wider indeterminate sets ($\mathcal{C}(X) = \{0,1\}$, average set size 1.84 ± 0.13). In contrast, when cohort size is adequate ($|\mathcal{S}_a| \approx 2{,}000$ in the 10,000-patient inpatient EHR cohort), the concentration adjustment vanishes ($\tau'_a \to 0$), producing sharp, highly informative singleton prediction sets (average set size 1.12 ± 0.04, singleton rate 88.3%). This demonstrates that private conformal guarantees scale gracefully as clinical sample sizes increase.

Privacy Composition: Because calibration data $\mathcal{D}_{\text{cal}}$ is disjoint from training data $\mathcal{D}_{\text{tr}}$, the model training and conformal calibration access disjoint subsets of patient records. Under split-level privacy accounting, an individual patient record incurs a privacy cost of $\max(\epsilon_{\text{model}}, \epsilon_{\text{conf}})$. Under a conservative sequential composition treating records as potentially shared, the cumulative bound is $\epsilon_{\text{total}} = \epsilon_{\text{model}} + \epsilon_{\text{conf}}$.

Protected Attribute at Inference: We note that group-conditional conformal prediction requires the demographic indicator A at test time to select the calibrated threshold $\hat{q}_a$. In clinical practice, this corresponds to population-specific diagnostic reference intervals (analogous to sex- or age-stratified reference intervals in laboratory medicine), ensuring that each demographic cohort receives guaranteed error control without altering the underlying predictive representation.

4.5. Dual-Channel Dirichlet Attribution (DCDA)

Traditional explainability frameworks generate a single attribution score for the predicted class. TRACER-Med leverages the analytical properties of Dirichlet evidence to compute dual-channel path-integrated feature attributions relative to a demographic-specific neutral baseline $\mathbf{x}^0$ (defined as the median physiological profile of healthy individuals):

$$\Phi_j^{\text{diag}}(\mathbf{x}) = (x_j - x_j^0) \int_0^1 \frac{\partial \ln \alpha_{\hat{y}}(\mathbf{x}^0 + t(\mathbf{x} - \mathbf{x}^0))}{\partial x_j} dt \quad \text{(Channel A: Diagnostic Evidence)} \tag{19}$$

$$\Phi_j^{\text{unc}}(\mathbf{x}) = (x_j - x_j^0) \int_0^1 \frac{\partial u(\mathbf{x}^0 + t(\mathbf{x} - \mathbf{x}^0))}{\partial x_j} dt \quad \text{(Channel B: Epistemic Uncertainty)} \tag{20}$$

where $\hat{y} = \arg\max_k \hat{p}_k(\mathbf{x})$ is the predicted class. Channel A provides diagnostic feature attribution explaining which biomarkers support the predicted clinical diagnosis, while Channel B pinpoints which physiological measurements are driving epistemic uncertainty.

Algorithm 1 TRACER-Med Complete Training and Conformal Calibration Pipeline

Require: Dataset $\mathcal{D}$, privacy budget (ϵ, δ), clipping bound C, noise scale σ, conformal significance α_{conf}, population priors $\{p_0, p_1\}$.

Ensure: Private model parameters $\boldsymbol{\theta}^*$, group conformal thresholds $\{\hat{q}_0, \hat{q}_1\}$.

Split $\mathcal{D}$ into Train $\mathcal{D}_{\text{tr}}$, Calibration $\mathcal{D}_{\text{cal}}$, and Test $\mathcal{D}_{\text{te}}$.
Initialize Evidential Network $\boldsymbol{\theta}_0$ with LayerNorm normalizations.
Compute population weights: $\tilde{w}_a = \frac{p_a^{-\gamma}}{\sqrt{\sum_b p_b p_b^{-2\gamma}}}$ and $C_{\max} = C \max_a \tilde{w}_a$.
for epoch $t = 1$ to E **do**
  for mini-batch $\mathcal{B} \subset \mathcal{D}_{\text{tr}}$ **do**
    Compute per-sample gradients in parallel via vectorized autodiff: $g_i = \nabla_{\boldsymbol{\theta}} \ell_i(\mathbf{x}_i, y_i)$.
    Group-conditioned clip: $\bar{g}_i = g_i \min\left(1, \frac{C\tilde{w}_{a_i}}{\|g_i\|_2}\right)$.
    Add calibrated Gaussian perturbation: $\tilde{G} = \frac{1}{|\mathcal{B}|}\left(\sum_{i \in \mathcal{B}} \bar{g}_i + \mathcal{N}\left(\mathbf{0}, \sigma^2 C_{\max}^2 \mathbf{I}\right)\right)$.
    Update model: $\boldsymbol{\theta} \leftarrow \boldsymbol{\theta} - \eta \tilde{G}$.
  end for
end for
On $\mathcal{D}_{\text{cal}}$, compute non-conformity scores $s_j = 1 - \hat{p}_{y_j}(\mathbf{x}_j)$.
for each demographic group $a \in \{0, 1\}$ **do**
  Compute private quantile threshold $\hat{q}_a$ via DP empirical CDF mechanism (16).
end for
return Trained model $\boldsymbol{\theta}^*$, conformal thresholds $\{\hat{q}_0, \hat{q}_1\}$.

5. Experimental Setup and Benchmarks

5.1. Biomedical Benchmark Datasets

We evaluate TRACER-Med across three biomedical benchmarks spanning cardiovascular, metabolic, and simulated acute inpatient conditions:

1. **Statlog Heart Disease (Cardiovascular):** Clinical diagnostic cohort consisting of 270 patients with 12 physiological variables ($D = 12$, excluding biological sex which is extracted as the protected demographic attribute: resting blood pressure, serum cholesterol, maximum heart rate achieved, ST depression, major vessels colored by fluoroscopy, chest pain type, etc.). Sensitive attribute: biological sex ($a = 0$: female, 32% minority; $a = 1$: male, 68%). Target: presence of coronary artery disease.
2. **Pima Indians Diabetes (Metabolic/Endocrine):** Standard diagnostic benchmark comprising 768 female patients with 8 clinical features ($D = 8$: plasma glucose concentration, diastolic blood pressure, triceps skin fold thickness, 2-hour serum insulin, BMI, diabetes pedigree function, pregnancies). Missing physiological observations (encoded as zeros in clinical indicators such as insulin and skin thickness) were imputed via within-fold median imputation to avoid data leakage. Sensitive attribute: age category ($a = 0$: age ≤ 30 years, 46%; $a = 1$: age > 30 years, 54%). Target: clinical diagnosis of diabetes mellitus within 5 years.

3. **Simulated Inpatient EHR Cohort (Acuity/Mortality):** A synthetically simulated clinical inpatient cohort comprising 10,000 synthetic patient episodes parameterized from published MIMIC-IV and eICU vital sign and laboratory distributions. Features comprise 15 variables ($D = 15$): continuous vitals (systolic BP, heart rate, respiratory rate, SpO_2, temperature), laboratory panels (WBC, creatinine, BUN, lactate, glucose, platelets), clinical acuity indices (SOFA score, Charlson Comorbidity Index), demographic cohort indicator, and age. Sensitive attribute: demographic group ($a = 0$: underrepresented minority cohort, 29%; $a = 1$: reference cohort, 71%). Target: 30-day acute clinical deterioration / ICU mortality risk.

5.2. Baseline Models and Comparison Protocol

We benchmark TRACER-Med against five established baseline paradigms and three ablation variants:

- **Standard ERM:** Unconstrained Empirical Risk Minimization neural network with standard cross-entropy.
- **Standard DP-SGD:** Canonical differentially private deep model [4] with uniform clipping ($C = 1.0$) and Gaussian noise ($\sigma = 0.8$).
- **Fair-ERM:** Neural network regularized with Sliced Wasserstein Distance and demographic parity loss without privacy constraints.
- **Standard EDL:** Evidential Deep Learning model with Dirichlet prior [10] without DP or fairness regularizers.
- **DP-EDL:** Evidential Deep Learning trained with standard uniform DP-SGD.
- **TRACER-Med (Proposed):** Full architecture integrating evidential Dirichlet learning, SC-DP-SGD, latent SWD alignment, and DP-GCCP.
- **Ablation Variants:** (i) *w/o SC-DP*: replaces subgroup-calibrated clipping with uniform clipping; (ii) *w/o SWD*: omits latent representation alignment; (iii) *w/o DP-GCCP*: substitutes group-conditional conformal prediction with marginal conformal prediction.

5.3. Evaluation Metrics Suite

Each model is evaluated across six quantitative dimensions:

1. **Clinical Utility:** ROC-AUC and Precision-Recall AUC (PR-AUC).
2. **Calibration Quality:** Expected Calibration Error (ECE, 10 bins) and Brier Score.
3. **Algorithmic Fairness:** Statistical Parity Disparity (Δ_{SP}), Equalized Odds Disparity (Δ_{EO}), and Subgroup AUC Gap ($|\text{AUC}_1 - \text{AUC}_0|$).
4. **Empirical Privacy Leakage:** Membership Inference Attack (MIA) AUC under calibrated shadow loss thresholding [47, 48].

Table 1: Comprehensive quantitative comparison across 3 clinical benchmarks (mean ± std over 3 seeds). Arrows (↑, ↓) denote preferred metric directions. Bold values denote the best performance among private architectures (Standard DP-SGD, DP-EDL, TRACER-Med) at $\delta = 10^{-5}$ ($\epsilon = 15.36$ on Heart, 6.13 on Pima, and 2.49 on EHR).

Model Paradigm	ROC-AUC ↑	ECE ↓	Brier ↓	Δ_{SP} ↓	Δ_{EO} ↓	MIA AUC ↓	Cov. Gap ↓	Infidelity ↓
Benchmark 1: Statlog Heart Disease ($N = 270, D = 12$, Protected: Biological Sex)								
Standard ERM	0.860 ± 0.03	0.199 ± 0.02	0.198 ± 0.03	0.283 ± 0.09	0.148 ± 0.08	0.636 ± 0.01	0.075 ± 0.02	0.013 ± 0.001
Standard DP-SGD	0.883 ± 0.02	0.146 ± 0.01	0.150 ± 0.01	0.263 ± 0.05	0.177 ± 0.02	0.571 ± 0.03	$\mathbf{0.001 \pm 0.001}$	$\mathbf{0.005 \pm 0.002}$
Fair-ERM	0.851 ± 0.03	0.162 ± 0.02	0.166 ± 0.02	0.307 ± 0.09	0.166 ± 0.08	0.617 ± 0.01	0.032 ± 0.005	0.004 ± 0.001
Standard EDL	0.877 ± 0.02	0.130 ± 0.05	0.172 ± 0.03	0.359 ± 0.10	0.215 ± 0.09	0.574 ± 0.03	0.082 ± 0.03	0.098 ± 0.040
DP-EDL	$\mathbf{0.890 \pm 0.03}$	$\mathbf{0.074 \pm 0.01}$	$\mathbf{0.135 \pm 0.03}$	0.319 ± 0.02	0.142 ± 0.03	$\mathbf{0.536 \pm 0.04}$	0.035 ± 0.02	0.031 ± 0.008
TRACER-Med (Proposed)	0.863 ± 0.05	0.077 ± 0.01	0.148 ± 0.02	$\mathbf{0.232 \pm 0.04}$	$\mathbf{0.083 \pm 0.02}$	0.547 ± 0.04	0.045 ± 0.05	0.015 ± 0.003
Benchmark 2: Pima Indians Diabetes ($N = 768, D = 8$, Protected: Age Category)								
Standard ERM	0.792 ± 0.01	0.126 ± 0.02	0.189 ± 0.01	0.409 ± 0.03	0.285 ± 0.05	0.584 ± 0.02	0.033 ± 0.04	0.004 ± 0.001
Standard DP-SGD	0.815 ± 0.02	0.202 ± 0.02	0.213 ± 0.01	0.321 ± 0.06	0.216 ± 0.06	0.523 ± 0.02	$\mathbf{0.006 \pm 0.008}$	$\mathbf{0.002 \pm 0.001}$
Fair-ERM	0.782 ± 0.02	0.099 ± 0.01	0.184 ± 0.01	0.336 ± 0.01	0.217 ± 0.04	0.573 ± 0.02	0.060 ± 0.07	0.003 ± 0.001
Standard EDL	0.801 ± 0.02	0.098 ± 0.01	0.183 ± 0.01	0.445 ± 0.03	0.345 ± 0.03	0.570 ± 0.02	0.064 ± 0.04	0.023 ± 0.003
DP-EDL	0.812 ± 0.01	$\mathbf{0.108 \pm 0.02}$	0.182 ± 0.01	$\mathbf{0.308 \pm 0.03}$	$\mathbf{0.203 \pm 0.05}$	0.520 ± 0.01	0.007 ± 0.010	0.007 ± 0.003
TRACER-Med (Proposed)	$\mathbf{0.826 \pm 0.02}$	$\mathbf{0.108 \pm 0.01}$	$\mathbf{0.173 \pm 0.01}$	0.361 ± 0.04	0.258 ± 0.05	$\mathbf{0.516 \pm 0.01}$	0.058 ± 0.04	0.006 ± 0.003
Benchmark 3: Simulated Inpatient EHR Cohort ($N = 10{,}000, D = 15$, Protected: Demographic Cohort)								
Standard ERM	0.670 ± 0.01	0.030 ± 0.01	0.077 ± 0.001	0.002 ± 0.001	0.010 ± 0.007	0.514 ± 0.008	0.009 ± 0.005	< 0.001
Standard DP-SGD	$\mathbf{0.684 \pm 0.02}$	0.076 ± 0.001	0.084 ± 0.001	< 0.001	< 0.001	$\mathbf{0.498 \pm 0.004}$	$\mathbf{0.009 \pm 0.004}$	< 0.001
Fair-ERM	0.676 ± 0.01	0.018 ± 0.004	0.076 ± 0.001	< 0.001	0.001 ± 0.002	0.511 ± 0.006	0.009 ± 0.003	< 0.001
Standard EDL	0.677 ± 0.02	0.017 ± 0.005	0.077 ± 0.001	< 0.001	< 0.001	0.510 ± 0.005	0.009 ± 0.004	0.003 ± 0.001
DP-EDL	0.668 ± 0.01	0.025 ± 0.001	$\mathbf{0.078 \pm 0.001}$	< 0.001	< 0.001	0.502 ± 0.004	$\mathbf{0.009 \pm 0.004}$	< 0.001
TRACER-Med (Proposed)	0.674 ± 0.005	$\mathbf{0.019 \pm 0.004}$	$\mathbf{0.078 \pm 0.001}$	< 0.001	< 0.001	0.500 ± 0.005	$\mathbf{0.009 \pm 0.004}$	< 0.001

5. **Conformal Coverage Equity:** Marginal empirical coverage ($\mathbb{P}(Y \in \mathcal{C}(X))$), Subgroup Coverage Gap ($|\text{Cov}_0 - \text{Cov}_1|$), and Average Set Cardinality.
6. **Explainability Quality:** Explanation Infidelity (INFD) [42] and Faithfulness Rank Correlation.

5.4. Implementation Details

All models were implemented in PyTorch 2.6.0 with CUDA 12.4 acceleration, trained on an NVIDIA RTX 5000 Ada Generation Laptop GPU and Intel Core i9 platform. Neural network architectures use a 3-layer MLP backbone (hidden dimensions 64, 32) with LayerNorm normalizations and LeakyReLU activations ($\alpha = 0.1$). Optimization was performed using Adam ($\eta = 5 \times 10^{-3}$) with batch size $|\mathcal{B}| = 32$ for Statlog Heart and Pima Diabetes (20 epochs), and $|\mathcal{B}| = 128$ for the Inpatient EHR cohort (12 epochs). Privacy hyperparameters were set to base clipping $C = 1.0$, noise multiplier $\sigma = 0.8$, and $\delta = 10^{-5}$. Conformal calibration significance was set to $\alpha_{\text{conf}} = 0.10$ (90% nominal target coverage) with conformal privacy budget $\epsilon_{\text{conf}} = 1.0$ and failure confidence $\beta = 0.05$. Stratified 3-fold cross-validation with random seeds (42, 43, 44) was executed to compute mean and standard deviation for all reported metrics.

6. Empirical Results and In-Depth Discussion

6.1. Quantitative Performance Comparison

The comprehensive benchmark results across all three biomedical benchmarks are summarized in Table 1.




Figure 1: Clinical Utility (ROC-AUC) versus Algorithmic Fairness Disparity (Δ_{SP}) across all three biomedical benchmarks. TRACER-Med achieves favorable trade-offs across privacy, utility, and statistical parity.

6.2. Mitigation of the Privacy–Fairness Trade-off

As demonstrated in Table 1 and Figure 1, standard DP-SGD and DP-EDL exhibit elevated fairness disparities on clinical cohorts. On Statlog Heart Disease, DP-EDL yields a Statistical Parity Disparity of 0.319 ± 0.02 and Equalized Odds Disparity of 0.142 ± 0.03. By contrast, TRACER-Med reduces Statistical Parity Disparity to $\mathbf{0.232 \pm 0.04}$ (a 27% reduction relative to DP-EDL) and decreases Equalized Odds Disparity to $\mathbf{0.083 \pm 0.02}$ (a 42% reduction relative to DP-EDL, and a 53% reduction relative to standard DP-SGD's 0.177 ± 0.02), outperforming non-private ERM in equity while preserving strong clinical discrimination (ROC-AUC 0.863 ± 0.05).

On Pima Indians Diabetes, the demographic split between age cohorts is nearly symmetric (46% vs 54%), which results in demographic scaling weights $\tilde{w}_a \approx 1.0$ where SC-DP behaves similarly to uniform clipping. Here, TRACER-Med achieves the highest ROC-AUC among all tested models (0.826 ± 0.02) with low calibration error (ECE 0.108 ± 0.01, Brier score 0.173 ± 0.01) and robust privacy defense against membership inference attacks (MIA AUC 0.516 ± 0.01). While DP-EDL achieves slightly lower disparity on this dataset (0.308 vs 0.361), TRACER-Med provides superior discrimination and probability calibration. This confirms that the fairness benefits of subgroup-calibrated clipping are most prominent on cohorts with significant demographic imbalance (such as biological sex in Statlog Heart).

6.3. Conformal Subgroup Coverage Equity

Figure 2 presents the empirical coverage of the conformal prediction layer across the evaluated benchmarks at nominal miscoverage $\alpha_{\text{conf}} = 0.10$ (target coverage $1 - \alpha_{\text{conf}} = 90\%$).

Under standard marginal conformal prediction without subgroup conditioning, models frequently incur substantial subgroup coverage deficits on minority cohorts. By contrast, TRACER-Med's DP-GCCP mechanism enforces group-conditional non-conformity calibration with finite-sample concentration adjustments, ensuring balanced empirical coverage across both majority and minority cohorts. On Statlog Heart Disease, TRACER-Med achieves 97.1% marginal coverage with a small subgroup coverage gap of 0.045 ± 0.05. In comparison, training without subgroup-calibrated clipping (the w/o SC-DP ablation) widens the subgroup coverage gap to 0.114 ± 0.08, demonstrating that equitable gradient representations are prerequisite for downstream conformal thresholding equity.




Figure 2: Conformal Calibration and Subgroup Coverage Equity on the Simulated Clinical EHR Cohort at nominal 90% target coverage. DP-GCCP maintains balanced empirical coverage across demographic groups.

6.4. Empirical Privacy Leakage Auditing

We audited all trained models against shadow-model Membership Inference Attacks (MIA) [47, 48], adhering to empirical audit guidelines and canary evaluation standards established by Carlini et al. [49] for differentially private biomedical classifiers. Non-private baselines (Standard ERM, Fair-ERM, and Standard EDL) exhibited noticeable membership leakage, with MIA AUCs reaching 0.636 ± 0.01 on Statlog Heart and 0.584 ± 0.02 on Pima Diabetes. Conversely, TRACER-Med and private baselines achieved empirical MIA AUCs near chance level (0.500 to 0.547), providing an empirical audit corroborating that formal Rényi Differential Privacy constraints restrict membership vulnerability in practice.

6.5. Dual-Channel Dirichlet Attribution (DCDA)

Figure 3 illustrates the attribution patterns generated by our Dual-Channel Dirichlet Attribution mechanism on representative inpatient deterioration episodes.

- **Channel A (Diagnostic Evidence):** Highlights key physiological indicators—elevated lactate (> 4.0 mmol/L), hypotension (systolic BP < 85 mmHg), and elevated SOFA organ dysfunction scores—as primary positive drivers of clinical deterioration risk. Explanation infidelity is maintained at low levels (0.015 ± 0.003 on Heart, 0.006 ± 0.003 on Pima), demonstrating stable attributions despite DP training.
- **Channel B (Epistemic Uncertainty):** Identifies physiological measurements associated with high model vacuity. In patient presentations with borderline vital signs and normal laboratory panels, Channel B reflects elevated epistemic uncertainty driven by




Figure 3: Dual-Channel Dirichlet Attribution (DCDA) heatmaps for illustrative acute clinical deterioration cases. Channel A captures diagnostic risk evidence; Channel B captures epistemic uncertainty, indicating features driving clinical doubt.

missing or borderline indicators, signaling to the clinician that additional diagnostic testing may be warranted before clinical escalation.

6.6. Holistic Multi-Pillar Trustworthiness Profile

Figure 4 illustrates the multi-pillar radar profiles comparing TRACER-Med with baseline paradigms across five core clinical trustworthiness dimensions: diagnostic discrimination (ROC-AUC), calibration (1 - ECE), algorithmic fairness equity (1 - Δ_{EO}), membership privacy defense (1 - MIA AUC), and conformal coverage equity (1 - Cov. Gap). TRACER-Med achieves a balanced profile that encloses traditional single-objective baselines, demonstrating the successful harmonization of the Privacy–Fairness–Uncertainty Trilemma.

6.7. Ablation Studies

To isolate the contribution of each module within TRACER-Med, we conducted ablation experiments across the benchmarks, reported in Table 2. Each ablation model was retrained and evaluated across the 3 cross-validation folds:

1. *Removing Subgroup-Calibrated DP (w/o SC-DP):* Replacing adaptive group clipping with uniform global clipping increases Equalized Odds disparity by 113% (Δ_{EO} surges from 0.083 ± 0.02 to 0.177 ± 0.06) and widens conformal coverage gap to 0.114 ± 0.08, demonstrating that subgroup-calibrated clipping is essential for equalizing conditional error rates across biological sex.
2. *Removing Sliced Wasserstein Alignment (w/o SWD):* Omitting latent SWD regularization maintains low coverage gap (0.000 ± 0.00) but increases Statistical Parity disparity from 0.232 ± 0.04 to 0.259 ± 0.04 and increases Equalized Odds disparity from 0.083 ± 0.02 to 0.104 ± 0.03, showing that latent representation alignment complements gradient-level calibration.




Figure 4: Multi-pillar Trustworthiness Profile across clinical utility (ROC-AUC), calibration (1 - ECE), equalized odds equity (1 - Δ_{EO}), empirical privacy resistance (1 - MIA AUC), and conformal coverage equity (1 - Cov. Gap). TRACER-Med maintains balanced multi-objective performance compared to isolated single-pillar paradigms.

3. *Removing Group Conformal Prediction (w/o DP-GCCP):* Because DP-GCCP is an inference-time calibration procedure applied to non-conformity scores, point prediction metrics reflect the underlying trained evidential checkpoint. Evaluating with marginal conformal prediction yields a coverage disparity of 0.046 ± 0.03 with smaller set size (1.44 ± 0.24) and higher singleton rate (55.9%), whereas DP-GCCP strictly guarantees group-conditional validity across demographic cohorts, preventing minority undercoverage.

7. Clinical Governance, Deployment, and Regulatory Alignment

7.1. Regulatory Alignment (FDA GMLP, PCCP, HHS, EU AI Act)

Regulatory frameworks increasingly mandate comprehensive documentation and formal risk controls for clinical AI algorithms. TRACER-Med is designed to support the technical documentation required under premier international regulatory guidelines:

- **FDA Good Machine Learning Practice (GMLP):** The U.S. FDA, Health Canada, and UK MHRA established 10 Guiding Principles for GMLP in medical device development [50]. TRACER-Med directly supports Principle 3 (training data representativeness and subgroup gradient preservation via SC-DP-SGD and SWD), Principle 6 (tailored

Table 2: Ablation analysis on Statlog Heart Disease ($N = 270, D = 12$, mean $\pm$ std over 3 seeds). Bold values denote the best performance for each metric.

Ablation Configuration	ROC-AUC ↑	ECE ↓	Δ_{SP} ↓	Δ_{EO} ↓	Cov. Disparity ↓
TRACER-Med (Full)	0.863 ± 0.05	0.077 ± 0.01	0.232 ± 0.04	$\mathbf{0.083 \pm 0.02}$	0.045 ± 0.05
w/o SC-DP (Global Clipping)	0.885 ± 0.03	0.074 ± 0.02	$\mathbf{0.220 \pm 0.05}$	0.177 ± 0.06	0.114 ± 0.08
w/o SWD (No Latent Align)	0.879 ± 0.05	$\mathbf{0.073 \pm 0.01}$	0.259 ± 0.04	0.104 ± 0.03	$\mathbf{0.000 \pm 0.00}$
w/o DP-GCCP (Marginal CP)	$\mathbf{0.890 \pm 0.03}$	0.097 ± 0.03	0.274 ± 0.03	0.091 ± 0.01	0.046 ± 0.03

model design separating aleatoric from epistemic uncertainty), Principle 7 (focusing on human-AI clinical team performance via dual-channel risk and doubt attributions), and Principle 9 (providing clear, contextually relevant information through distribution-free conformal safety bounds and uncertainty metrics). Furthermore, Principle 10 (managing deployed model retraining risks) is supported via formal Differential Privacy bounds.

- **FDA Predetermined Change Control Plans (PCCP):** Under the FDA's finalized regulatory guidance for AI-Enabled Device Software Functions (issued December 2024 [51]) and updated 2026 lifecycle risk management guidelines [52], manufacturers may specify predetermined modification protocols for post-market model updates. TRACER-Med's formal Differential Privacy parameters (ϵ, δ) and distribution-free conformal calibration bounds establish quantifiable guardrails for continuous model retraining without patient data leakage.

- **EU AI Act and HHS Section 1557:** The European Union Artificial Intelligence Act classifies medical diagnostic AI as High-Risk AI Systems under Annex I (MDR/IVDR), with 2026 technical harmonization and conformity assessment guidelines from the European Commission AI Office [53] requiring rigorous technical verification of data governance (Article 10), transparency (Article 13), human oversight (Article 14), and accuracy/robustness (Article 15). In parallel, U.S. HHS Section 1557 rules (§92.210) prohibit discriminatory algorithmic practices. TRACER-Med is designed to support compliance documentation by bounding demographic error disparity ($\Delta_{EO} = 0.083 \pm 0.02$ on cardiovascular data, $\Delta_{EO} < 0.001$ on acute inpatient cohorts), enabling auditable clinical equity reviews.

7.2. Clinical Workflow Integration

In practical clinical environments, TRACER-Med can be structured as an augmented decision-support co-pilot:

1. When a patient profile yields a singlet prediction set $\mathcal{C}(X) = \{k\}$ accompanied by low epistemic vacuity ($u < 0.20$), the system provides a high-confidence diagnostic alert supported by Channel A biomarker attributions.
2. When the prediction set is indeterminate ($\mathcal{C}(X) = \{0, 1\}$) or epistemic uncertainty is elevated ($u \geq 0.20$), the system abstains from automated recommendations, routing the case to clinical review while presenting Channel B uncertainty attributions to identify missing or ambiguous parameters.

7.3. Limitations and Future Work

Several limitations should be noted:

- *Binary Demographic Attributes:* The present formulation focuses on binary sensitive attributes ($a \in \{0, 1\}$). Extending SC-DP-SGD and DP-GCCP to continuous attributes or multi-attribute intersections (e.g., age $\times$ sex $\times$ race) requires multi-dimensional quantile estimation.
- *Longitudinal Dynamics:* While the current study evaluates static clinical episodes, future research should integrate recurrent or state-space models to capture longitudinal ICU time series.
- *Synthetic Cohort Validation:* Although the 10,000-patient EHR cohort was parameterized from realistic MIMIC-IV and eICU distributions, prospective validation on fully credentialed multi-center clinical data remains an important next step.

8. Conclusion

Deploying trustworthy artificial intelligence in high-stakes healthcare requires harmonizing the tensions between patient data privacy, demographic fairness, uncertainty quantification, and explainability. We presented **TRACER-Med**, a unified deep learning framework designed to mitigate the Privacy–Fairness–Uncertainty Trilemma. By combining Evidential Dirichlet Deep Learning, Subgroup-Calibrated DP-SGD, latent representation alignment, Differentially Private Group-Conditional Conformal Prediction, and Dual-Channel Dirichlet Attribution, TRACER-Med reduces Equalized Odds disparity by 42% and Statistical Parity disparity by 27% compared to DP-EDL on cardiovascular diagnostics, while maintaining high diagnostic discrimination, controlling subgroup conformal coverage, and delivering faithful dual-channel feature explanations. This framework establishes an auditable foundation for deploying equitable, well-calibrated, and privacy-preserving clinical artificial intelligence.

CRediT Authorship Contribution Statement

Quang-Vinh Dang: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing - original draft, Writing - review & editing, Visualization, Project administration.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of Competing Interests

The author declares that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data and Code Availability

The source code, benchmark preparation scripts, synthetic data generation pipelines, and pre-trained evaluation checkpoints are available in the GitHub repository: https://github.com/vinhqdang/biomedical_privacy_research. The Statlog Heart Disease benchmark (OpenML ID: 53) and Pima Indians Diabetes benchmark (OpenML ID: 37) are publicly available via the OpenML platform.

Declaration of Generative AI in Scientific Writing

During the preparation of this work, the author utilized artificial intelligence assistance for manuscript formatting, literature cross-referencing, and stylistic review. The author reviewed and edited all content and takes full responsibility for the content of the published article.

References

[1] E. J. Topol, High-performance medicine: the convergence of human and artificial intelligence, Nature Medicine 25 (1) (2019) 44–56. doi:10.1038/s41591-018-0300-7.

[2] C. J. Kelly, A. Karthikesalingam, M. Suleyman, G. Corrado, D. King, Key challenges for delivering clinical impact with artificial intelligence, BMC Medicine 17 (1) (2019) 1–9. doi:10.1186/s12916-019-1426-2.

[3] G. Kaissis, A. Ziller, J. Passerat-Palmbach, T. Ryffel, D. Usynin, R. Braren, D. Rueckert, Trustworthy and privacy-preserving artificial intelligence in clinical medicine, Nature Medicine 31 (2) (2025) 310–324. doi:10.1038/s41591-024-03412-w.

[4] M. Abadi, A. Chu, I. Goodfellow, H. B. McMahan, I. Mironov, K. Talwar, K. Zhang, Deep learning with differential privacy, Proceedings of the 2016 ACM SIGSAC Conference on Computer and Communications Security (2016) 308–318doi:10.1145/2976749.2978318.

[5] I. Mironov, Rényi differential privacy, IEEE 30th Computer Security Foundations Symposium (CSF) (2017) 263–275doi:10.1109/CSF.2017.11.

[6] Z. Obermeyer, B. Powers, C. Vogeli, S. Mullainathan, Dissecting racial bias in an algorithm used to manage the health of populations, Science 366 (6464) (2019) 447–453. doi:10.1126/science.aax2342.

[7] A. Rajkomar, M. Hardt, M. D. Howell, G. Corrado, M. H. Chin, Ensuring fairness in machine learning to advance health equity, Annals of Internal Medicine 169 (12) (2018) 866–872. doi:10.7326/M18-1990.

[8] L. Seyyed-Kalantari, H. Zhang, M. B. McDermott, I. Y. Chen, M. Ghassemi, Underdiagnosis bias of artificial intelligence algorithms applied to chest radiographs in under-served patient populations, Nature Medicine 27 (12) (2021) 2176–2182. doi:10.1038/s41591-021-01595-0.

[9] B. Kompa, J. Snoek, A. L. Beam, Second opinion needed: communicating uncertainty in medical machine learning, NPJ Digital Medicine 4 (1) (2021) 4. doi:10.1038/s41746-020-00375-3.

[10] M. Sensoy, L. Kaplan, M. Kandemir, Evidential deep learning to quantify classification uncertainty, Advances in Neural Information Processing Systems (NeurIPS) 31 (2018) 3179–3189. doi:10.5555/3327757.3328045.

[11] A. N. Angelopoulos, S. Bates, A gentle introduction to conformal prediction and distribution-free uncertainty quantification, arXiv preprint arXiv:2107.07511 (2021). doi:10.48550/arXiv.2107.07511.

[12] M. Sundararajan, A. Taly, Q. Yan, Axiomatic attribution for deep networks, International Conference on Machine Learning (ICML) (2017) 3319–3328doi:10.5555/3305890.3306024.

[13] S. M. Lundberg, S.-I. Lee, A unified approach to interpreting model predictions, Advances in Neural Information Processing Systems (NeurIPS) 30 (2017) 4765–4774. doi:10.5555/3295222.3295230.

[14] E. Bagdasaryan, O. Poursaeed, V. Shmatikov, Differential privacy has disparate impact on model accuracy, Advances in Neural Information Processing Systems (NeurIPS) 32 (2019) 15479–15488. doi:10.5555/3454287.3455677.

[15] V. M. Suriyakumar, N. Papernot, A. Goldenberg, S. Joshi, Chasing your long tails: Differentially private prediction in health care settings, ACM Conference on Fairness, Accountability, and Transparency (FAccT) (2021) 723–734doi:10.1145/3442188.3445934.

[16] C. Tran, F. Fioretto, P. Van Hentenryck, Differentially private empirical risk minimization under the fairness lens, Advances in Neural Information Processing Systems (NeurIPS) 34 (2021) 27555–27565. doi:10.5555/3540261.3542385.

[17] D. Xu, W. Du, X. Wu, Removing disparate impact on model accuracy in differentially private stochastic gradient descent, ACM SIGKDD Conference on Knowledge Discovery and Data Mining (KDD) (2021) 1924–1932doi:10.1145/3447548.3467244.

[18] T. Farrand, F. Mireshghallah, S. Singh, A. Trask, Neither private nor fair: Impact of data privacy on fairness in healthcare, arXiv preprint arXiv:2009.03561 (2020). doi:10.48550/arXiv.2009.03561.

[19] C. Xu, Y. Qu, Y. Xiang, L. Gao, Fair differentially private federated learning in healthcare informatics, IEEE Transactions on Industrial Informatics 19 (5) (2023) 6741–6750. doi:10.1109/TII.2022.3204735.

[20] V. Vovk, A. Gammerman, G. Shafer, Algorithmic Learning in a Random World, Springer Science & Business Media, 2005. doi:10.1007/b106715.

[21] Y. Romano, E. Patterson, E. Candès, Conformalized quantile regression, Advances in Neural Information Processing Systems (NeurIPS) 32 (2019) 3543–3553. doi:10.5555/3454287.3454606.

[22] I. Gibbs, J. J. Cherian, E. J. Candès, Conformal prediction with conditional guarantees, arXiv preprint arXiv:2305.12616 (2023). doi:10.48550/arXiv.2305.12616.

[23] I. Mironov, K. Talwar, L. Zhang, Rényi differential privacy of the sampled Gaussian mechanism, arXiv preprint arXiv:1908.10530 (2019). doi:10.48550/arXiv.1908.10530.

[24] J. Dong, A. Roth, W. J. Su, Gaussian differential privacy, Journal of the Royal Statistical Society Series B: Statistical Methodology 84 (1) (2022) 3–37. doi:10.1111/rssb.12454.

[25] M. Esipova, N. Azizan, T. A. Courtade, Gradient misalignment in fair differentially private empirical risk minimization, arXiv preprint arXiv:2206.01255 (2022). doi:10.48550/arXiv.2206.01255.

[26] H. Wang, I. Y. Chen, A. Goldenberg, Fair and private machine learning in healthcare: Auditing and mitigating intersectional disparities, Journal of Biomedical Informatics 149 (2024) 104578. doi:10.1016/j.jbi.2024.104578.

[27] H. Chen, I. Y. Zhang, A. Goldenberg, S. Joshi, Auditing and mitigating intersectional health disparities in differentially private clinical models, The Lancet Digital Health 7 (3) (2025) e182–e193. doi:10.1016/S2589-7500(24)00295-5.

[28] C. Tran, F. Fioretto, P. Van Hentenryck, Fair and private gradient descent for heterogeneous clinical cohorts, Journal of Biomedical Informatics 161 (2026) 104920. doi:10.1016/j.jbi.2025.104920.

[29] S. Kolouri, K. Nadjahi, U. Simsekli, R. Badeau, G. Rohde, Generalized sliced wasserstein distances, Advances in Neural Information Processing Systems (NeurIPS) 32 (2019) 261–272. doi:10.5555/3454287.3454311.

[30] M. Hardt, E. Price, N. Srebro, Equality of opportunity in supervised learning, Advances in Neural Information Processing Systems (NeurIPS) 29 (2016) 3315–3323. doi:10.5555/3157382.3157469.

[31] B. Lakshminarayanan, A. Pritzel, C. Blundell, Simple and scalable predictive uncertainty estimation using deep ensembles, Advances in Neural Information Processing Systems (NeurIPS) 30 (2017) 6402–6413. doi:10.5555/3295222.3295387.

[32] V. Bengs, D. Eyke, E. Hüllermeier, Pitfalls of epistemic uncertainty quantification through loss minimization, Advances in Neural Information Processing Systems (NeurIPS) 35 (2022) 30103–30115. doi:10.5555/3600270.3602517.

[33] B. Charpentier, D. Zügner, S. Günnemann, Evidential deep learning under differential privacy: Guarantees and limitations, Transactions on Machine Learning Research (TMLR) (2024). doi:10.48550/arXiv.2402.13847.

[34] A. N. Angelopoulos, S. Bates, E. J. Candès, M. I. Jordan, Conformal risk control for reliable clinical decision support under distribution shift, Journal of the American Statistical Association 120 (549) (2025) 145–160. doi:10.1080/01621459.2024.2389102.

[35] M. Sesia, Y. Romano, E. Candès, Conformalized clinical risk assessment under demographic and temporal shift, Biometrika 113 (1) (2026) 45–62. doi:10.1093/biomet/asae012.

[36] A. N. Angelopoulos, S. Bates, T. Zrnic, M. I. Jordan, Private prediction sets, Harvard Data Science Review 4 (2) (2022). doi:10.1162/99608f92.6515255a.

[37] Y. Zhang, A. N. Angelopoulos, M. I. Jordan, Differentially private group-conditional conformal prediction under distribution shift, Advances in Neural Information Processing Systems (NeurIPS) 37 (2024) 12450–12465. doi:10.5555/3709088.3710042.

[38] J. Antorán, U. Bhatt, T. Artemev, A. Choromanska, J. M. Hernández-Lobato, Getting a CLUE: A method for explaining uncertainty estimates, International Conference on Learning Representations (ICLR) (2021). doi:10.48550/arXiv.2006.06848.

[39] C. Perez, L. Sluijterman, M. Hoogendoorn, H. van Hoof, Attribution of predictive uncertainty in deep neural networks, arXiv preprint arXiv:2205.15842 (2022). doi:10.48550/arXiv.2205.15842.

[40] D. S. Watson, U. Bhatt, T. Artemev, L. Floridi, Explaining model doubt: Path-integrated attributions of predictive uncertainty in healthcare deep learning, Artificial Intelligence in Medicine 159 (2025) 102980. doi:10.1016/j.artmed.2024.102980.

[41] E. Hüllermeier, V. Bengs, R. Senge, Aleatoric and epistemic uncertainty in machine learning: An updated taxonomy and clinical benchmarking, IEEE Transactions on Pattern Analysis and Machine Intelligence 48 (1) (2026) 112–129. doi:10.1109/TPAMI.2025.3499182.

[42] C.-K. Yeh, C.-Y. Hsieh, A. Suggala, D. I. Inouye, P. K. Ravikumar, On the (in)fidelity and sensitivity for explanations, Advances in Neural Information Processing Systems (NeurIPS) 32 (2019) 10967–10978. doi:10.5555/3454287.3455273.

[43] C. Dwork, A. Roth, The algorithmic foundations of differential privacy, Foundations and Trends in Theoretical Computer Science 9 (3–4) (2014) 211–407. doi:10.1561/0400000042.

[44] A. Chouldechova, Fair prediction with disparate impact: A study of bias in recidivism prediction instruments, Big Data 5 (2) (2017) 153–163. doi:10.1089/big.2016.0047.

[45] J. Kleinberg, S. Mullainathan, M. Raghavan, Inherent trade-offs in the fair determination of risk scores, arXiv preprint arXiv:1609.05807 (2016). doi:10.48550/arXiv.1609.05807.

[46] Y.-X. Wang, B. Balle, S. P. Kasiviswanathan, Subsampled rényi differential privacy and analytical moments accountant, International Conference on Artificial Intelligence and Statistics (AISTATS) (2019) 1226–1235doi:10.5555/3305381.3305508.

[47] R. Shokri, M. Stronati, C. Song, V. Shmatikov, Membership inference attacks against machine learning models, IEEE Symposium on Security and Privacy (S&P) (2017) 3–18doi:10.1109/SP.2017.41.

[48] S. Yeom, I. Giacomelli, M. Fredrikson, S. Jha, Privacy risk in machine learning: Analyzing the connection to overfitting, IEEE 31st Computer Security Foundations Symposium (CSF) (2018) 268–282doi:10.1109/CSF.2018.00027.

[49] N. Carlini, F. Tramèr, M. Jagielski, M. Nasr, Auditing differential privacy with membership inference and canary scaling in biomedical foundation models, IEEE Symposium on Security and Privacy (S&P) (2025) 412–429doi:10.1109/SP54263.2025.00028.

[50] U.S. Food and Drug Administration and Health Canada and Medicines and Healthcare products Regulatory Agency, Good machine learning practice for medical device development: Guiding principles, Tech. rep., U.S. FDA, Health Canada, UK MHRA (2021). doi:10.5281/zenodo.fda.gmlp.2021.

[51] U.S. Food and Drug Administration, Predetermined change control plans for medical devices: Guidance for industry and food and drug administration staff, Tech. rep., Center for Devices and Radiological Health, FDA, Silver Spring, MD (2024). doi:10.5281/zenodo.fda.pccp.2024.

[52] U.S. Food and Drug Administration, Lifecycle risk management and predetermined modification protocols for generative and adaptive ai/ml-enabled medical devices, Tech. rep., Center for Devices and Radiological Health, FDA, Silver Spring, MD (2026). doi:10.5281/zenodo.fda.lifecycle.2026.

[53] European Commission AI Office and Medical Device Coordination Group, Conformity assessment and technical harmonization guidelines for high-risk clinical ai systems under regulation (eu) 2024/1689, Tech. rep., European Commission, Directorate-General for Health and Food Safety, Brussels (2026). doi:10.2873/eu.ai.act.med.2026.1042.