Cluster Editing above a P_3 packing: the treewidth of the intersection graph

Quang-Vinh Dang
British University Vietnam, Hung Yen, Vietnam
vinh.dq4@buv.edu.vn

Abstract

Cluster Editing asks for the fewest edge insertions and deletions that turn a graph into a disjoint union of cliques. Li, Pilipczuk and Sorge parameterize it by the excess ℓ of the solution size over a modification-disjoint packing H of induced P_3's, and leave open for which packings the problem is fixed-parameter tractable. We use the treewidth w of the intersection graph $I(H)$ of the packing as the structural measure. For $\ell = 0$ the problem is a constraint satisfaction problem with domain size 3 whose constraint scopes are cliques of $I(H)$; for general ℓ, guessing at most ℓ edits gives such a problem with domain size 7. This yields algorithms in time $3^{w+1}n^{O(1)}$ and $7^{w+1}n^{2\ell+O(1)}$, and $n^{\ell} f(\ell, w)\, 7^{w+\ell+1} n^{O(1)}$ after a locality argument. Under ETH there is no $2^{o(w)}n^{O(1)}$ algorithm even for $\ell = 0$, and neither w nor ℓ alone suffices. If the components of $I(H)$ have bounded size, the problem is fixed-parameter tractable in ℓ and has a kernel. For pair-maximal packings we give a normal form of optimal solutions, a dichotomy of inserted pairs into local and non-local ones, and an instance family in which an optimum must delete an edge between members at arbitrary distance in $I(H)$. Whether the problem is fixed-parameter tractable in $\ell + w$ is open.

1 Introduction

Cluster Editing asks for the fewest edge insertions and deletions that turn a graph G into a disjoint union of cliques. Every induced P_3 must be hit by an edit, so a packing H of induced P_3's that pairwise share no vertex pair (a *modification-disjoint* packing) gives the lower bound $|H|$ on the solution size. Li, Pilipczuk and Sorge [14] parameterize by the *excess* ℓ over this bound (problem CEaMP): is there a solution of size at most $|H| + \ell$? They show NP-hardness already for $\ell = 0$, with every vertex in at most 23 packed P_3's, and an $n^{2\ell+O(1)}$ algorithm when every vertex lies in at most two packed P_3's; whether the latter case is fixed-parameter tractable is open [14]. For vertex-disjoint packings the problem is fixed-parameter tractable in ℓ [15].

The question of this paper is which measurable *interaction structure* of H makes the local repair choices (one edit per packed P_3, in three ways) combine into a global solution in $f(\ell, w)\, n^{O(1)}$ time. We take $w = \mathrm{tw}(I(H))$, where the intersection graph $I(H)$ has the packed P_3's as nodes and joins two of them when they share a vertex.

1.1 Results

Our first theorem collects the results of Sections 3 and 4.

Theorem 1.1 (Main theorem)**.** *Let (G, H, ℓ) be an instance of CEaMP, $w = \mathrm{tw}(I(H))$, and assume a tree decomposition of $I(H)$ of width w is given.*

(a) CSP representation. $(G, H, 0)$ *is a YES instance iff a constraint satisfaction problem with one variable of domain* $\{1, 2, 3\}$ *per packed* P_3 *is satisfiable; every constraint has at most three variables, and its scope is a clique of* $I(H)$*. For* $\ell > 0$*, for each of the at most* $n^{2\ell}$ *sets* U *of at most* ℓ *hard pairs there is such a problem with domain size* 7 *and unary costs, and* (G, H, ℓ) *is a YES instance iff for some* U *its minimum cost is at most* $|H| + \ell - |U|$*.*

(b) Algorithms. *CEaMP is solvable in time* $3^{w+1}n^{O(1)}$ *for* $\ell = 0$ *and in time* $7^{w+1}n^{2\ell+O(1)}$ *for general* ℓ*. The latter is XP in* ℓ *with single-exponential dependence on* w*; it is not a fixed-parameter algorithm.*

(c) ETH lower bound. *Unless ETH fails, there is no* $2^{o(w)}n^{O(1)}$ *algorithm for* $\ell = 0$*, even if every vertex lies in at most* 23 *packed* P_3*'s.*

(d) Neither parameter suffices alone. *CEaMP is NP-hard when* $w = 0$ *(with* ℓ *part of the input) and, by [14], when* $\ell = 0$*.*

Parts (a) and (b) are Theorems 3.4 and 3.5; (c) and (d) are Theorem 4.1 and Proposition 4.3. The CSP view and the dynamic program are standard once the interaction structure is identified (Observation 3.2); (c) is a consequence of the reduction of [14].

The remaining results concern the structure of optimal solutions and the question whether (ℓ, w) suffices.

- *Bounded components* (Section 5). If s is the maximum number of vertices spanned by a connected component of $I(H)$, CEaMP is solvable in time $1.62^{(\binom{s}{2}+1)\ell}n^{O(1)} + n3^s$ and has a kernel with at most $(s^2 - s + 2)\ell$ vertices; in particular $I(H)$ a perfect matching is fixed-parameter tractable in ℓ (Theorem 5.1). This is the reduction rule of van Bevern, Froese and Komusiewicz [15] applied to components of $I(H)$; we verify its hypotheses and the counting at the level of components, and cite the soundness of the rule [15, Lemma 6.3], the algorithm of Böcker [2] and the $2k$ kernel of Chen and Meng [4]. A variant for components of arbitrary size leaves at most 2ℓ components (Theorem 5.10).

- *Normal form and an* n^ℓ *algorithm* (Sections 6 and 7). For a pair-maximal packing (no induced P_3 has all three pairs hard), every cluster that contains an end of a hard edit is connected, of size at most $2D + \ell + 1$ and of bounded hard degree, and one end of a hard edit determines at most Z^2 candidates for the other, where $Z = O(w + \ell)$ (Corollary 6.7). This gives an algorithm in time $(n+1)^\ell(6w+9\ell+7)^{2\ell}7^{w+\ell+1}n^{O(1)}$ for every packing (Theorem 7.2), halving the exponent of Theorem 3.5, and a fixed-parameter algorithm for a stronger parameter ω (Theorem 7.3). The parameter ω is not bounded by any function of (ℓ, w) (Proposition 7.4).

- *Locality of inserted pairs* (Section 8). Every inserted hard pair of an optimal solution of a pair-maximal packing has a middle vertex and is either of type T1 (two soft steps in distinct packed P_3's; local in a decomposition of $I(H)$) or of type T2 (one kept hard step). Merges of two pieces joined by no soft pair are of type T1 only (Corollary 8.3); merges joined by a soft pair can be of type T2 (examples with $n = 7$ and $n = 8$, Proposition 8.7). For every $m \geq 3$ there is an instance with $w = 1$, $c = 2$, $\ell = 1$ in which every optimum deletes a hard edge whose ends lie in packed P_3's at distance $m - 1$ in $I(H)$ (Proposition 8.8).

Appendices A–G contain further proved statements that are not used in the main text: twins, the structure of the vertices outside the packing and a shrink rule, counting lemmas, the dynamic program for ω, conflict weights and a pruned variant, a family on which reduction on unions of packed P_3's fails, and tunnel gadgets.

What is open. **OPEN:** whether CEaMP is fixed-parameter tractable in $\ell + \mathrm{tw}(I(H))$, or W[1]-hard. This is open already for $w = 1$ with tree components of unbounded size. Our algorithms are XP in ℓ, Theorem 5.1 needs bounded components, and no hardness is known. The normal form of Section 6 applies to a pair-maximal extension of the given packing, whose intersection graph may have larger treewidth. Section 9 lists the open problems and summarises, in one table, which result holds for the input packing and which for a pair-maximal extension.

1.2 Related work

van Bevern, Froese and Komusiewicz [15] parameterize F-free editing above packings; for Cluster Editing they obtain fixed-parameter algorithms for vertex-disjoint P_3 packings and leave open the modification-disjoint case, which Li, Pilipczuk and Sorge [14] settle negatively for $\ell = 0$ and study for $1/c$-integral packings. Modification-disjoint P_3 packings are also used as a lower-bound tool for cluster vertex splitting [8]; for broader context see the survey [5]. To our knowledge, the treewidth of the intersection graph of the packing has not been used as a parameter before. Komusiewicz and Uhlmann [12] show that Cluster Editing is NP-hard on graphs of maximum degree 6 and has no $2^{o(k)}n^{O(1)}$ algorithm under ETH. The critical-clique (twin) property used in Appendix A goes back to [10, 7]; the half-degree property of Lemma 6.2 is the same single-vertex-move argument. Cubic expanders of linear treewidth are cited from Grohe and Marx [9].

2 Preliminaries

Graphs and solutions. Graphs are simple and undirected, $n = |V(G)|$. A *pair* is a 2-subset of $V(G)$; for a set S of pairs, $G \triangle S$ flips the adjacency of each pair in S. A graph is a *cluster graph* iff it has no induced P_3, i.e. no vertex triple with exactly two edges. A *solution* is a set S of pairs with $G \triangle S$ a cluster graph; $\mathrm{OPT}(G)$ is the minimum size of a solution. Solutions correspond to partitions of V into *clusters* (the cliques of $G \triangle S$); the *cost* of a partition is the number of pairs ab with $[ab \in E] \neq [a, b \text{ in one cluster}]$, and an *optimum* is a partition of minimum cost. We write $\alpha(\cdot)$ for the independence number and $\nu(\cdot)$ for the matching number.

Packings, soft and hard pairs. A *packing* $H = \{P_1, \ldots, P_p\}$ is a set of induced P_3's $P_i = (a_i, b_i, c_i)$ with centre b_i ($a_ib_i, b_ic_i \in E$, $a_ic_i \notin E$); its elements are the *members* or *packed P_3's*. Write $\mathrm{pairs}(P_i)$ for the three pairs inside $V(P_i)$. H is *modification-disjoint* if the sets $\mathrm{pairs}(P_i)$ are pairwise disjoint; this is equivalent to: any two members share at most one vertex. All packings below are modification-disjoint. A pair is *soft* if it lies in $\bigcup_i \mathrm{pairs}(P_i)$ (its *owner* is then unique) and *hard* otherwise; a *hard edge* is a hard pair in E and a *hard non-edge* a hard pair outside E. For a vertex v, $Q(v)$ is the set of soft partners of v, $N_h(v)$ the set of hard neighbours and $d_h(v) = |N_h(v)|$. G_{soft} is the *soft graph* on V whose edges are the soft pairs.

Problem CEaMP. Given G, a modification-disjoint packing H and $\ell \geq 0$, decide whether there is a solution S with $|S| \leq |H| + \ell$ (the convention of [14]; ℓ is the *excess*). We call (G, H, ℓ) a *YES instance* if $\mathrm{OPT}(G) \leq |H| + \ell$.

The intersection graph. The *intersection graph* $I(H)$ has node set H and an edge PP' iff $V(P) \cap V(P') \neq \emptyset$. The members through one vertex form a clique of $I(H)$; hence (Helly property of tree decompositions) they lie in a common bag. Put $c = c(H) = \max_v |\{P \ni v\}|$ and $w = \mathrm{tw}(I(H))$ (with $w = 0$ if $I(H)$ has no edge), so $c \leq w + 1$. Tree decompositions of width w are assumed *given*

in the algorithmic statements; one of width w is computed in time $2^{O(w^3)}n$ [3, 6], and one of width at most $2w+1$ in time $2^{O(w)}n$ [13], which would change the bases below (e.g. 7^{w+1} to 7^{2w+2}).

Lemma 2.1 (Degrees). *For every vertex v, $|Q(v)| = 2\,|\{P \in H : v \in V(P)\}|$. Hence $D(H) := \max_v |Q(v)| = 2c \le 2(w+1)$.*

Proof. Each member $P \ni v$ contributes the two soft pairs of P at v, i.e. two partners. Two members through v cannot have a common partner $x \neq v$, since they would then share the two vertices v, x, hence a pair. So the partner sets are disjoint and $|Q(v)| = 2|\{P \ni v\}|$. □

Below, D denotes a *number* with $D \ge \max_v |Q(v)|$ (for instance $D = D(H) = 2c$), and $D(H)$ the function value.

Lemma 2.2 (Solutions decompose). *Let S be any set of pairs such that $G \triangle S$ is a cluster graph, $x_i = S \cap \mathrm{pairs}(P_i)$ and $U = S \setminus \bigcup_i \mathrm{pairs}(P_i)$ (the* hard edits*). Then every $x_i \neq \emptyset$, $|S| = \sum_i |x_i| + |U|$, and $|U| \le |S| - |H|$.*

Proof. If $x_i = \emptyset$ the triple $V(P_i)$ keeps its adjacencies in $G \triangle S$ and is an induced P_3. The pair sets are disjoint, so $|S| = \sum |x_i| + |U| \ge |H| + |U|$. □

Consequently the *excess* of a solution is $\sum_i (|x_i| - 1) + |U|$, and every solution of size at most $|H| + \ell$ has $|U| \le \ell$.

2.1 Maximalisation and pair-maximal packings

A family Φ of induced P_3's is *hard over H* if the sets $\mathrm{pairs}(P)$, $P \in \Phi$, are pairwise disjoint and disjoint from the soft pairs of H; then $H \cup \Phi$ is a packing with $|H| + |\Phi|$ members. H is *pair-maximal* if no induced P_3 of G has all three pairs hard, i.e. if H is maximal among modification-disjoint packings. This is strictly stronger than maximality among vertex-disjoint packings (Proposition 4.5).

Lemma 2.3 (Hard families). *Let (G, H, ℓ) be a YES instance and Φ a family hard over H. Then $|\Phi| \le \ell$. In general* $\mathrm{OPT} \ge |H| + |\Phi|$.

Proof. $H \cup \Phi$ is a modification-disjoint packing of $|H| + |\Phi|$ induced P_3's. Lemma 2.2, applied to $H \cup \Phi$ and an optimum solution, gives $\mathrm{OPT} \ge |H| + |\Phi|$. In a YES instance $\mathrm{OPT} \le |H| + \ell$. □

Lemma 2.4 (Maximalisation). *Let H be any packing and M a maximal family hard over H (inclusion-maximal; found greedily in polynomial time). Put $H' = H \cup M$, $\ell' = \ell - |M|$. Then:*

(a) $\mathrm{OPT} \ge |H'|$. *In particular $|M| > \ell$ implies that (G, H, ℓ) is a NO instance.*

(b) $\mathrm{OPT} \le |H| + \ell$ *iff* ($|M| \le \ell$ *and* $\mathrm{OPT} \le |H'| + \ell'$). *So (G, H, ℓ) and (G, H', ℓ') are equivalent whenever $\ell' \ge 0$.*

(c) H' is a pair-maximal packing ("hard" now means hard with respect to H'; the pairs of M have become soft).

(d) $\mathrm{tw}(I(H')) \le \mathrm{tw}(I(H)) + |M|$, *hence $w' \le w + |M| \le w + \ell$.*

(e) $c' \le \min\{c + |M|,\, w' + 1\}$ and $D(H') = 2c' \le D(H) + 2|M|$.

A tree decomposition of $I(H')$ of width at most $w + |M|$ is obtained in polynomial time from one of $I(H)$.

Proof. (a) is Lemma 2.3 (second sentence: OPT $\geq |H| + |M|$). (b) If OPT $\leq |H| + \ell$ then $|M| \leq \ell$ by (a), and $|H'| + \ell' = |H| + \ell$; the converse is the same identity.

(c) The members of M have pairwise disjoint pair sets, disjoint from those of H, so H' is a packing. Suppose an induced P_3 Q has all its pairs hard with respect to H'. Then $M \cup \{Q\}$ is a family hard over H ($Q \notin M$ because the pairs of Q are not in pairs(M)), contradicting maximality of M.

(d) Let $(\mathcal{T}, \{\beta_t\})$ be a tree decomposition of $I(H)$ of width w (if $H = \emptyset$ take a single empty bag). $I(H')[H] = I(H)$. Put $\beta'_t = \beta_t \cup M$. Every node of H still occupies a connected subtree. Every node of M occupies all of $\mathcal{T}$. An edge of $I(H')$ with both ends in H is covered as before; an edge with an end in M is covered by any bag containing its other end (if both ends are in M every bag covers it). Bags have at most $w + 1 + |M|$ nodes.

(e) At most $|M|$ members of M pass through a vertex, so $c' \leq c + |M|$; $c' \leq w' + 1$ is the inequality $c \leq w + 1$ for H'. By Lemma 2.1, $D(H') = 2c' \leq D(H) + 2|M|$. □

Example 2.5 (Tightness of Lemma 2.4(d))**.** Let G be the star with centre v and leaves $p, q, a_1, b_1, \ldots, a_m, b_m$, $H = \{(p, v, q)\}$ and $M = \{(a_i, v, b_i) : 1 \leq i \leq m\}$. M is a maximal family hard over H: every induced P_3 of a star is (x, v, y) with leaves x, y, and every pair xv is already soft. $I(H)$ is one node, tw $= 0$; $I(H \cup M) = K_{m+1}$, tw $= m = 0 + |M|$.

Convention: pair-maximal packings and primed parameters. Sections 6–8 and the appendices state their results for a *pair-maximal* H; where a result needs a YES instance or an optimal solution, it says so. For an arbitrary input packing H let M be as in Lemma 2.4 and read the statements for the *primed instance* (G, H', ℓ') with parameters $w' \leq w + |M|$, $c' \leq c + |M|$, $D' \leq D + 2|M|$, $\ell' = \ell - |M|$. The family M is not unique (it depends on the greedy order), so quantities defined from H' (such as ω in Section 7) are parameters of the primed instance, not functions of the input (G, H, ℓ). Only Theorems 3.4, 3.5, 5.1, 5.10 and 7.2 are statements about the input H in terms of w, ℓ and n; the table in Section 9 lists the scope of every result.

Notation. $P_i = (a_i, b_i, c_i)$ with centre b_i; c is the maximum number of members through a vertex (unsubscripted), and $\kappa_i \geq 1$ the number of edited pairs among pairs(P_i). $U = U^- \cup U^+$ is the set of hard edits, U^- the deleted hard edges and U^+ the inserted hard non-edges, $u = |U|$. $\ell' = \ell - |M|$ is the primed excess and $\bar{\ell}$ the excess left after component accounting (Section 5). $\mathcal{C}_j$ are the components of $I(H)$ and W_j the vertex sets they span. $R = V \setminus V(H)$ (Appendix A). $B = 2D + 3\ell$, $Z = B + D + 1$ and BIG are defined in Section 6. Hard components K and the cliques $\mathcal{K}_v$ of the constructions in Section 7 are distinct objects.

3 The interaction CSP and its dynamic programs

Observation 3.1. *Let S be a solution with $|S| \leq |H|$. Then S has exactly one pair in each* pairs(P_i) *and no other pair, and $G \triangle S$ is a cluster graph iff no vertex triple is an induced P_3 of $G \triangle S$. Consequently $(G, H, 0)$ is a YES instance iff the following CSP is satisfiable: variables $x_i \in \{1, 2, 3\}$ (which pair of P_i is flipped), and for every vertex triple t the constraint "t is not an induced P_3 after flipping", whose truth depends on the owners' variables of the soft pairs of t only (hard pairs keep their status in G).*

Proof. By Lemma 2.2, $|x_i| \geq 1$ for all i and $|S| = \sum |x_i| + |U| \leq |H|$ force $|x_i| = 1$ and $U = \emptyset$. The rest is the definition of a cluster graph. □

The CSP has $\Theta(n^3)$ constraints, not $O(|H|)$.

Observation 3.2. *The set of owners of the soft pairs of any triple is a clique of $I(H)$ (at most three nodes). Hence the primal graph of the CSP is a subgraph of $I(H)$ (not every edge of $I(H)$ is witnessed by a triple), and a tree decomposition of $I(H)$ is one of the primal graph.*

Proof. Two pairs of one triple share a vertex v; their owners both contain v, so they coincide or are adjacent in $I(H)$. □

The natural parameter of the CSP is therefore the treewidth of its primal graph, which is at most w; the ETH bound of Section 4 is stated for $I(H)$.

Lemma 3.3 (Dynamic programming over $I(H)$)**.** *Let there be one variable x_i per member, with a domain $\mathcal{D}_i$ of size at most d and a unary cost $\gamma_i : \mathcal{D}_i \to \mathbb{N}$, and let $\mathcal{C}$ be a set of constraints, each given by a test of its satisfaction on $\prod \mathcal{D}_i$ in polynomial time, whose scope is a clique of $I(H)$. Given a tree decomposition of $I(H)$ of width w, a satisfying assignment of minimum total cost is found in time $d^{w+1}(|H| + |\mathcal{C}|)^{O(1)}$.*

Proof. Let $(\mathcal{T}, \{\beta_s\})$ be the tree decomposition. By the Helly property each clique lies in a bag; assign each constraint to one such bag. For a node s and an assignment α of the variables of β_s, let $f_s(\alpha)$ be the minimum cost over the variables in the subtree of s, among assignments extending α and satisfying all constraints assigned in the subtree. The standard recurrences (introduce, forget, join; see [6]) compute all f_s; each table has at most d^{w+1} entries. □

Theorem 3.4. *Given (G, H) and a tree decomposition of $I(H)$ of width w, CEaMP with $\ell = 0$ is solvable in $3^{w+1}n^{O(1)}$ time.*

Proof. Observations 3.1 and 3.2 give a CSP with domains of size 3, all unary costs 1 and $O(n^3)$ constraints with clique scopes; apply Lemma 3.3. □

Theorem 3.5. *Given (G, H, ℓ) and a tree decomposition of $I(H)$ of width w, CEaMP is solvable in $7^{w+1}n^{2\ell+O(1)}$ time.*

Proof. Let $\mathcal{D}_i$ be the 7 non-empty subsets of pairs(P_i). By Lemma 2.2, a solution of size $\leq |H| + \ell$ has the form $S = U \cup \bigcup x_i$ with $x_i \in \mathcal{D}_i$ and U a set of at most ℓ hard pairs; there are at most $\sum_{j \leq \ell} \binom{n^2/2}{j} \leq n^{2\ell}$ candidates for U.

Fix U and put $G_U = G \triangle U$. Since U avoids soft pairs, every P_i is still an induced P_3 of G_U with the same pairs, so H is still a modification-disjoint packing. For a triple t, the status of each hard pair of t in $G_U \triangle x$ is fixed, and each soft pair is determined by G_U and the variable of its owner. Hence t gives a constraint C_t ("t is not an induced P_3 in $G_U \triangle x$") whose scope is the owner set of t, a clique of $I(H)$ (the proof of Observation 3.2 does not use G itself). Triples with empty scope are tested directly, in $O(n^3)$ time. Seek $x \in \prod \mathcal{D}_i$ satisfying all C_t and minimizing $\sum_i |x_i|$; by Lemma 3.3 this takes $7^{w+1}n^{O(1)}$ time. Accept iff for some U the minimum m_U satisfies $m_U + |U| \leq |H| + \ell$. Soundness: any feasible (U, x) gives the solution $U \cup \bigcup x_i$. Completeness: Lemma 2.2. □

Theorem 3.4 is the case $\ell = 0$ of Theorem 3.5 with the smaller base 3. Together with Observations 3.1 and 3.2 and Lemma 3.3 these are parts (a) and (b) of Theorem 1.1. The algorithm of Theorem 3.5 is not fixed-parameter tractable: ℓ is part of the input and appears in the exponent of n, and Proposition 4.3 shows that w alone does not suffice.

Remark 3.6 (Domain $\{1,2,3\}$ does not suffice for $\ell > 0$). In the encoding above, a variable ranges over the non-empty subsets of pairs(P_i). For $\ell > 0$ it cannot be restricted to the three single-pair values: an edited packed P_3 can receive two or three edits in every optimum. For two: $n = 5$, $E = \{02, 04, 12, 23, 24\}$, $H = \{(1,2,3)\}$, OPT $= 2$, unique optimum $\{12, 23\}$. For three: $n = 6$, $E = \{01, 02, 03, 04, 12, 14, 15, 24, 25, 35, 45\}$, $H = \{(0,3,5)\}$, OPT $= 3$, unique optimum $\{03, 05, 35\}$ (both checked by enumerating all set partitions). We do not claim that 7 is the smallest possible domain size in other encodings. For $c = 2$, $I(H)$ is subcubic but may have unbounded treewidth, so Theorem 3.5 is a different tradeoff from the $n^{2\ell}$ algorithm of [14], not a recovery of it.

4 Limits of the parameterization

4.1 An ETH lower bound in the exponent

Theorem 4.1. *Assume ETH. There is no algorithm that decides CEaMP with $\ell = 0$ in time $2^{o(\mathrm{tw}(I(H)))}n^{O(1)}$. This holds even when every vertex lies in at most 23 packed P_3's (so $I(H)$ has maximum degree at most 66), and whether or not a tree decomposition is supplied.*

The theorem is a corollary of the reduction of Li, Pilipczuk and Sorge.

Proof. By the construction behind [14, Thm. 1] there is a polynomial reduction from 3-SAT to CEaMP with $\ell = 0$ in which every vertex lies in at most 23 packed P_3's and the graph has $|V| = O(N+m)$ vertices for a formula with N variables and m clauses (each variable occurrence and each clause is replaced by a gadget of constant size). By the Sparsification Lemma [11] (see also [6]) we may assume $m = O(N)$ at the cost of $2^{o(N)}$ extra time, so $|V| = O(N)$.

Every member has three vertices and every vertex lies in at most 23 members, hence $3|H| = \sum_v |\{P \ni v\}| \le 23|V|$, i.e. $|H| \le 23|V|/3 = O(N)$. A single bag containing all members is a tree decomposition of $I(H)$, so $\mathrm{tw}(I(H)) \le |H| - 1 = O(N)$, and such a decomposition can be supplied. An algorithm with running time $2^{o(\mathrm{tw}(I(H)))}n^{O(1)}$, which may compute its own decomposition, would therefore decide 3-SAT in time $2^{o(N)}$, contradicting ETH. The degree bound: a member meets at most $3 \cdot 22 = 66$ others. □

Remark 4.2. (1) The bound is on the exponent only: it does not show that the base 3 (or 7) is tight, and $2^{o(w)}$ excludes $3^{o(w)}$ and $7^{o(w)}$ alike. A SETH-type bound $(3-\varepsilon)^w$ is open. (2) Since $\mathrm{tw}(I(H)) \le |H| - 1 \le n$, a $2^{o(\mathrm{tw}(I(H)))}n^{O(1)}$ algorithm is in particular a $2^{o(n)}$ algorithm, so the theorem is no stronger than the statement that CEaMP at $\ell = 0$ has no $2^{o(n)}$ algorithm under ETH; it says that the exponent of Theorem 3.4 cannot be improved to $o(w)$ on instances with $|H| = \Theta(n)$.

4.2 Other parameters do not suffice

Proposition 4.3. *CEaMP is NP-hard when $I(H)$ has no edge ($w = 0$, vertex-disjoint packing), with ℓ part of the input. Hence w alone is not enough, and (by Theorem 1 of [14]) neither is $\ell = 0$ alone.*

Proof. Cluster Editing is NP-hard (e.g. already on graphs of maximum degree 6 [12]). Given (G,k) compute greedily a maximal set H of vertex-disjoint induced P_3's (if none, G is a cluster graph). Vertex-disjoint implies modification-disjoint and $I(H)$ is edgeless. Since OPT $\ge |H|$, (G,k) is a YES instance iff $(G,H,k-|H|)$ is (a NO instance if $k < |H|$). □

Proposition 4.4. *(a) For the star $G = K_{1,2m}$ ($m \ge 1$) every packing $H \ne \emptyset$ has $I(H) = K_{|H|}$ and every YES instance (G,H,ℓ) satisfies $\ell + \mathrm{tw}(I(H)) \ge 2m - 2$, while $\mathrm{tw}(G) = 1$. (b) For the $k \times k$ grid G and a family H of vertex-disjoint induced P_3's along rows, $\mathrm{tw}(I(H)) = 0$ and $\mathrm{tw}(G) = k$.*

Proof. (a) Leaves have degree one, so the centre of any induced P_3 is the hub v; all packed P_3's are (a, v, b) with distinct leaves (the pairs va are disjoint), pairwise sharing v, so $I(H) = K_p$, $p = |H| \le m$, $\mathrm{tw} = p - 1$. If the hub's cluster contains j leaves, the cost is $(2m - j) + \binom{j}{2}$ (the other leaves are best left as singletons), which is minimized, with value $2m - 1$, at $j \in \{1, 2\}$; so $\mathrm{OPT} = 2m - 1$ and a YES instance has $\ell \ge 2m - 1 - p$, i.e. $\ell + \mathrm{tw}(I(H)) \ge 2m - 2$. (b) Rows of a grid are induced P_3's (no triangles) and $\mathrm{tw}(k \times k \text{ grid}) = k$ is standard. □

Proposition 4.4 concerns the numbers $\mathrm{tw}(G)$ and $\mathrm{tw}(I(H))$ only, which are incomparable. Part (a) is the only part that bears on the pair (ℓ, w): bounded $\mathrm{tw}(G)$ does not make $\ell + w$ small, because on a star small $\mathrm{tw}(I(H))$ forces large ℓ. Part (b) says nothing about ℓ.

4.3 Weaker maximality hypotheses fail

The following two instances show that the structure theory of Appendix A and Section 6 needs pair-maximality and that edits touching vertices outside $V(H)$ cannot be excluded.

Proposition 4.5 (Pair-maximality cannot be replaced by vertex-maximality)**.** *Let G have vertices $0, \dots, 4$ and edges $03, 13, 14, 24, 34$, and $H = \{(0, 3, 1)\}$ (pairs $03, 13, 01$). Then H is maximal among vertex-disjoint packings (only the two vertices $2, 4$ remain),* $\mathrm{OPT} = 2$ *and the unique optimum is $\{03, 24\}$; it contains the edit 24 with both endpoints outside $V(H)$. Here $\{3, 4, 2\}$ is an induced P_3 all of whose pairs are hard, so H is not pair-maximal.*

Proof. The partition $\{\{0\}, \{1, 3, 4\}, \{2\}\}$ has the two edits $03, 24$. Conversely, $0\,3\,1$ and $3\,4\,2$ are induced P_3's with disjoint pair sets $\{03, 13, 01\}$ and $\{34, 24, 23\}$, so $\mathrm{OPT} \ge 2$. That this is the only optimum is checked by enumerating all 52 partitions of the five vertices. □

A second example: for the path $0\,1\,2\,3\,4$ and $H = \{(0, 1, 2)\}$ the remaining vertices $3, 4$ cannot form a P_3, so H is maximal among vertex-disjoint packings, but $(2, 3, 4)$ is an induced P_3 whose pairs $23, 34, 24$ are disjoint from $\mathrm{pairs}(H)$, so H is not pair-maximal; the pair-maximal extension is no longer vertex-disjoint.

Proposition 4.6 (Strong locality fails)**.** *Let $G = C_4$ with edges $01, 02, 13, 23$ and $H = \{(0, 1, 3)\}$ (pairs $01, 13, 03$; vertex $2 \notin V(H)$). Then H is pair-maximal,* $\mathrm{OPT} = 2 = |H| + 1$*, so $(G, H, 0)$ is a NO instance, and there are exactly three optima $\{03, 12\}$, $\{01, 23\}$, $\{02, 13\}$, each containing a hard pair at vertex 2.*

Proof. Every induced P_3 of C_4 contains one of the soft pairs $01, 13, 03$, so H is pair-maximal. $\mathrm{OPT} \ge |H| = 1$, and a single edit leaves an induced P_3 (checked on the six pairs), so $\mathrm{OPT} = 2$; the three optima are found by enumerating all 15 partitions. □

This refutes the claim that all hard edits of some optimum have both endpoints in $V(H)$; what remains true is that no edit has both endpoints outside $V(H)$ (Theorem A.5 in Appendix A).

5 Components of $I(H)$

Let $\mathcal{C}_1, \dots, \mathcal{C}_r$ be the connected components of $I(H)$, $p_j = |\mathcal{C}_j|$ the number of members in $\mathcal{C}_j$, and $W_j = \bigcup_{P \in \mathcal{C}_j} V(P)$. Put $s = \max_j |W_j|$, the largest number of *vertices* spanned by a component. If $s_P = \max_j p_j$ counts members instead, then $s \le 2s_P + 1$: order the members of a component so that each shares a vertex with an earlier one; the first spans 3 vertices and each later one adds at most 2.

Theorem 5.1 (Component theorem). *CEaMP can be solved in time* $1.62^{(\binom{s}{2}+1)\ell}\, n^{O(1)} + n\, 3^s$ *and admits a kernel with at most* $(s^2 - s + 2)\,\ell$ *vertices, where* s *is the maximum number of vertices spanned by a connected component of* $I(H)$*. No maximality hypothesis is used. The kernel is an instance of CEaMP with empty packing. The proof uses the soundness of Reduction Rule 6.2 of [15] (their Lemma 6.3), the algorithm of Böcker [2] and the* $2k$*-vertex kernel of Chen and Meng [4].*

Corollary 5.2. *If* $I(H)$ *is a perfect matching (so* $w = 1$*,* $c = 2$*), CEaMP is fixed-parameter tractable in* ℓ*: here* $s = 5$*, the time is* $1.62^{11\ell} n^{O(1)}$ *and the kernel has at most* 22ℓ *vertices. More generally components with at most* s_P *members give Theorem 5.1 with* $s = 2s_P + 1$*.*

Proof. Two adjacent nodes of $I(H)$ share exactly one vertex (H is modification-disjoint), so $|W_j| = 3 + 3 - 1 = 5$; $s^2 - s + 2 = 22$ and $\binom{5}{2} + 1 = 11$. □

What is cited and what is proved. We use the problem of van Bevern, Froese and Komusiewicz [15]: an instance $(G, \mathcal{G}, k)$ consists of a graph, a set $\mathcal{G}$ of *vertex-disjoint induced subgraphs* G_j with $1 \le \tau(G_j) \le t$, where τ is the cluster-editing optimum, and a budget k; its parameters are t and $k - \sum_j \tau(G_j)$ (Definition 1.2 and Problem 1.3 of [15]; connectivity of the G_j is not required, and their size enters only through t). We prove below that the components of $I(H)$ define such an instance with excess at most ℓ (Lemma 5.3), that the reduction rule can be applied at component level without disturbing the other components (Lemma 5.4), and the bound $r \le 2\bar{\ell}$ on the number of irreducible components (Lemma 5.6); the last re-derives, for components, the counting argument of [15, Lemmas 3.2 and 6.5]. The following are *cited, not reproved*: (i) soundness of their Reduction Rule 6.2 for an arbitrary induced subgraph $G[W]$ (their Lemma 6.3, whose proof uses only G, W and the optimal solution S of $G[W]$); (ii) the time to test and apply the rule (their Lemma 6.4); (iii) the algorithm of Böcker [2] running in $O(1.62^k + n + m)$ and the kernel with at most $2k$ vertices for Cluster Editing [4], both as used in [15].

5.1 Components as a packing in the sense of van Bevern et al.

Lemma 5.3 (Components form a packing in the sense of [15]). *Let* $G_j = G[W_j]$*,* $\tau_j = \mathrm{OPT}(G_j)$*,* $k = |H| + \ell$ *and* $\bar{\ell} = k - \sum_j \tau_j$*.*

1. *The sets* W_j *are pairwise disjoint and each* G_j *is an induced subgraph.*
2. $p_j \le \tau_j \le \lfloor \binom{s}{2}/2 \rfloor =: t_s$ *and* $\tau_j \ge 1$*.*
3. $\bar{\ell} = \ell - \sum_j (\tau_j - p_j) \le \ell$*.*
4. *For every solution* S *of* G *and every* j*,* $S_j := S \cap \binom{W_j}{2}$ *is a solution of* G_j*, so* $|S_j| \ge \tau_j$*; the sets* $\binom{W_j}{2}$ *are pairwise disjoint, so* $|S| = \sum_j |S_j| + |S_{\mathrm{ext}}|$ *where* S_{ext} *are the pairs of* S *inside no* W_j*. In particular* $\mathrm{OPT}(G) \ge \sum_j \tau_j$*, and if* $\bar{\ell} < 0$ *then* (G, H, ℓ) *is a NO instance.*

Proof. (1) A vertex in $W_i \cap W_j$, $i \ne j$, lies in a member of $\mathcal{C}_i$ and in a member of $\mathcal{C}_j$; these are adjacent in $I(H)$, contradicting that $\mathcal{C}_i, \mathcal{C}_j$ are different components. Induced by definition.

(2) The members of $\mathcal{C}_j$ are induced P_3's of G_j with pairwise disjoint pair sets inside $\binom{W_j}{2}$. A solution of G_j must edit a pair of each of them (Lemma 2.2); hence $\tau_j \ge p_j \ge 1$. For the upper bound, let m_j be the number of edges of G_j; deleting all edges, or inserting all non-edges, yields a cluster graph, so $\tau_j \le \min(m_j, \binom{|W_j|}{2} - m_j) \le \binom{|W_j|}{2}/2 \le \binom{s}{2}/2$.

(3) $|H| = \sum_j p_j$ because the components partition H.

(4) $(G \triangle S)[W_j] = G_j \triangle S_j$ is an induced subgraph of a cluster graph, hence a cluster graph. The rest is counting with disjoint pair sets. □

Thus (G, H, ℓ) is a YES instance iff $(G, \{G_j\}, k)$ is a YES instance of the problem of [15] with $t \le t_s$ and excess $\bar{\ell} \le \ell$ (answer NO if $\bar{\ell} < 0$). Components, not single members, are needed: a pair uv with $u \in W_j$ and $v \notin W_j$ lies in no W_i at all when v is outside $V(H)$ or in another component, hence is not counted in $\sum_j \tau_j$ and is an excess edit (Lemma 5.3(4)); for a packing whose members share vertices, the same pair may be a soft pair of another member and is then free, which is the obstruction of [14].

5.2 Reducible sets and the counting lemma

A set $\mathcal{F}$ of members is *pair-closed* if no member outside $\mathcal{F}$ has two vertices in $W_{\mathcal{F}} := \bigcup_{P \in \mathcal{F}} V(P)$; every component of $I(H)$ is pair-closed, and then no outside member meets $W_{\mathcal{F}}$ at all. An optimal solution $S_{\mathcal{F}}$ of $G[W_{\mathcal{F}}]$ satisfies the *boundary condition* if for every $y \notin W_{\mathcal{F}}$ the set $N(y) \cap W_{\mathcal{F}}$ is empty or exactly one cluster of $G[W_{\mathcal{F}}] \triangle S_{\mathcal{F}}$. $\mathcal{F}$ is *reducible* if some optimal $S_{\mathcal{F}}$ satisfies it, and *irreducible* otherwise. This is the condition of Reduction Rule 6.2 of [15] for the induced subgraph $G[W_{\mathcal{F}}]$ (stated there as two clauses: vertices in one cluster have the same neighbours outside $W_{\mathcal{F}}$; vertices in different clusters have no common neighbour outside $W_{\mathcal{F}}$; the two forms are equivalent). A component $\mathcal{C}_j$ is *tight* if $\tau_j = p_j$.

Lemma 5.4 (Rule 6.2 at component level)**.** *If component $\mathcal{C}_j$ is reducible with S_j, then $(G, \{G_i\}, k)$ is a YES instance iff $(G \triangle S_j, \{G_i\}_{i \neq j}, k - \tau_j)$ is. In the new instance the graphs G_i $(i \neq j)$ are unchanged, still vertex-disjoint and induced, with the same τ_i; the excess $\bar{\ell}$ is unchanged; and for every $i \neq j$ the boundary condition, hence reducibility, is unchanged. So one pass over the components, in any order, reaches an instance in which no component is reducible.*

Proof. The equivalence is Lemma 6.3 of [15] (cited; sound for every induced subgraph). Both k and $\sum \tau$ drop by τ_j. The set S_j consists of pairs with both ends in W_j, so no pair incident to a vertex of W_i, $i \neq j$, changes: $G[W_i]$ and all neighbourhoods $N(u)$, $u \in W_i$, are the same before and after, and the boundary condition for i refers only to these. □

Lemma 5.5 (Boundary violations force an edited boundary pair)**.** *Let S be any solution, $\mathcal{F}$ pair-closed, $S_{\mathcal{F}} = S \cap \binom{W_{\mathcal{F}}}{2}$ and $y \notin W_{\mathcal{F}}$. If $N(y) \cap W_{\mathcal{F}}$ is neither empty nor a cluster of $G[W_{\mathcal{F}}] \triangle S_{\mathcal{F}}$, then S contains a pair $\{y, u\}$ with $u \in W_{\mathcal{F}}$.*

Proof. Suppose not. Then $N_{G \triangle S}(y) \cap W_{\mathcal{F}} = N_G(y) \cap W_{\mathcal{F}}$. In the cluster graph $G \triangle S$ this set is $\mathrm{cluster}(y) \cap W_{\mathcal{F}}$ (as $y \notin W_{\mathcal{F}}$), which is empty or a cluster of the induced subgraph $(G \triangle S)[W_{\mathcal{F}}] = G[W_{\mathcal{F}}] \triangle S_{\mathcal{F}}$. □

Lemma 5.6 (Counting irreducible components)**.** *Let S be a solution of G and J a set of irreducible components of $I(H)$. Then $|S| \ge \sum_j \tau_j + |J|/2$. In particular, if $|S| \le k = \sum_j \tau_j + \bar{\ell}$, then $|J| \le 2\bar{\ell}$.*

Proof. Claim: for every $j \in J$, (a) $|S_j| \ge \tau_j + 1$, or (b) S contains a pair with exactly one end in W_j (an *external* pair of j). Suppose (a) fails; then $|S_j| = \tau_j$ by Lemma 5.3(4), so S_j is an optimal solution of G_j, and since $\mathcal{C}_j$ is irreducible, S_j violates the boundary condition at some $y \notin W_j$. A component is pair-closed, so Lemma 5.5 gives a pair $\{y, u\} \in S$ with $u \in W_j$.

Let J_a be the indices of J satisfying (a) and J_b the others, which satisfy (b). An external pair of j has exactly one end in W_j; it lies in no $\binom{W_i}{2}$, so it belongs to S_{ext}, and it is external for at most two indices (the components containing its ends). Hence $|S_{\mathrm{ext}}| \ge |J_b|/2$. Since $|S| = \sum_j |S_j| + |S_{\mathrm{ext}}|$ and $|S_j| \ge \tau_j$ for all j, we get $|S| \ge \sum_j \tau_j + |J_a| + |J_b|/2 \ge \sum_j \tau_j + |J|/2$. □

The bound concerns irreducible components, not components with positive internal excess $\tau_j > p_j$: a reducible component may have large internal excess, and it is removed by Lemma 5.4 at no cost in $\bar{\ell}$.

Proof of Theorem 5.1. Algorithm. (1) Compute the components of $I(H)$, the optima τ_j by the subset dynamic programme over W_j ($O(3^{|W_j|})$), and $\bar{\ell}$; if $\bar{\ell} < 0$ answer NO (Lemma 5.3(4)). (2) For each component test reducibility and, if reducible, apply the rule (Lemma 5.4); the test of [15, Lemma 6.4] needs a constant number of exact cluster-editing computations on subgraphs of G_j plus polynomial time, which the subset programme provides in $O(3^{|W_j|})$ each. (3) Let r^* be the number of remaining components. If $r^* > 2\bar{\ell}$, answer NO (Lemma 5.6 with J the set of all remaining components). Otherwise the current instance has budget $k^* = \sum_{\text{remaining}} \tau_j + \bar{\ell} \le r^* t_s + \bar{\ell} \le (2t_s + 1)\bar{\ell}$. Forget the packing and solve plain Cluster Editing with budget k^* by Böcker's algorithm [2] in $O(1.62^{k^*} + n + m)$.

Correctness follows from Lemma 5.3 (the instance is of the required form and $\bar{\ell}$ is the excess), Lemma 5.4 (steps preserve the answer), and Lemma 5.6.

Time. The components have at least 3 vertices and are disjoint, so there are at most $n/3$ of them and steps (1), (2) take $O(n3^s) + n^{O(1)}$. Step (3) takes $1.62^{k^*} n^{O(1)}$ with $k^* \le (2t_s+1)\bar{\ell} \le (\binom{s}{2}+1)\ell$, since $2t_s \le \binom{s}{2}$ and $\bar{\ell} \le \ell$.

Kernel. After step (2) and the checks of step (3), apply the kernelization of Cluster Editing with at most $2k^*$ vertices [4] to (G, k^*) and add the empty packing: an equivalent CEaMP instance with empty packing, budget $\le k^*$ and at most $2k^* \le (4t_s+2)\ell \le (s^2 - s + 2)\ell$ vertices. □

Remark 5.7 (Scope). Theorem 5.1 is fixed-parameter tractable in (ℓ, s); it does not answer the main question, since s is not bounded by a function of w (a path or tree $I(H)$ has $w \le 1$ and arbitrarily large components), and the exponent $\binom{s}{2}\ell$ is of no practical interest. The argument uses neither pair-maximality nor Appendix A. For a perfect matching it covers the smallest non-trivial instance of the case $c = 2$ left open in [14] for fixed-parameter tractability in ℓ.

5.3 Components of arbitrary size

For components too large for the subset programme, tightness and reducibility can still be decided when w is small.

Proposition 5.8. *Given a tree decomposition of $I(H)$ of width w and a component $\mathcal{C}$ of $I(H)$, one decides in time $3^{w+1}n^{O(1)}$ whether $\mathcal{C}$ is tight; if it is tight, one decides in the same time whether it is reducible and finds a witnessing $S_{\mathcal{C}}$.*

Proof. A tight solution of $G[W_{\mathcal{C}}]$ consists of one edited pair per member and nothing else (each member needs at least one edited pair and the pair sets are disjoint), and every such solution is optimal since $\tau \ge p$; so for a tight $\mathcal{C}$ the optimal solutions are exactly the tight ones. Variables: the three choices of each member. Constraint: $G[W_{\mathcal{C}}] \triangle S$ has no induced P_3, i.e. every triple of $W_{\mathcal{C}}$ is fine. A triple with a soft pair involves at most three members (one per soft pair, pair sets being disjoint); each owns two of the three vertices of the triple, so any two of them share a vertex, they form a clique of $I(H)$ and lie in a common bag. A triple without soft pair is checked statically. So $\mathcal{C}$ is tight iff this CSP is satisfiable, and Lemma 3.3 decides this in $3^{w+1}n^{O(1)}$ time. The boundary condition at $y \notin W_{\mathcal{C}}$ with $A_y = N(y) \cap W_{\mathcal{C}} \neq \emptyset$ says: (i) every pair inside A_y is an edge of $G \triangle S$ and (ii) every pair between A_y and $W_{\mathcal{C}} \setminus A_y$ is a non-edge of $G \triangle S$ (given that $G[W_{\mathcal{C}}] \triangle S$ is a cluster graph this says A_y is a cluster). Each such pair is hard (static) or owned by one member (a unary constraint). Reducibility is therefore the same CSP with additional unary constraints. □

(For a non-tight component the optimal $S_{\mathcal{C}}$ has more than p edits and is not a 3^p assignment; the proposition says nothing about it.)

Lemma 5.9 (Reduction preserves pair-maximality)**.** *Let H be pair-maximal, $\mathcal{F} \subseteq H$ pair-closed and tight with $S_{\mathcal{F}}$ an optimal solution of $G[W_{\mathcal{F}}]$ satisfying the boundary condition, $G^\circ = G \triangle S_{\mathcal{F}}$ and $H^\circ = H \setminus \mathcal{F}$. Then H° is a modification-disjoint packing of G°, (G, H, ℓ) and (G°, H°, ℓ) are equivalent, H° is pair-maximal in G°, and $I(H^\circ)$ is an induced subgraph of $I(H)$, so* $\mathrm{tw}(I(H^\circ)) \leq \mathrm{tw}(I(H))$.

Proof. Members outside $\mathcal{F}$ have at most one vertex in $W_{\mathcal{F}}$, so none of their pairs lies in $W_{\mathcal{F}}$; $S_{\mathcal{F}}$ does not touch them, so they remain induced P_3's with the same pairs, and H° is a modification-disjoint packing of G°. Since $\mathcal{F}$ is tight, $|S_{\mathcal{F}}| = |\mathcal{F}|$, and Rule 6.2 (Lemma 6.3 of [15]) gives $\mathrm{OPT}(G) = \mathrm{OPT}(G^\circ) + |\mathcal{F}|$, hence the equivalence with unchanged ℓ.

Pair-maximality. Let Q be an induced P_3 of G°; we show that it has a soft pair with respect to H°. Let $q = |V(Q) \cap W_{\mathcal{F}}|$. If $q = 3$, Q is an induced P_3 of $G^\circ[W_{\mathcal{F}}] = G[W_{\mathcal{F}}] \triangle S_{\mathcal{F}}$, a cluster graph: impossible. If $q = 2$, $V(Q) = \{p, p', y\}$ with $p, p' \in W_{\mathcal{F}}$, $y \notin W_{\mathcal{F}}$. The pairs yp, yp' are unchanged, and by the boundary condition $N_{G^\circ}(y) \cap W_{\mathcal{F}}$ is empty or one cluster K of the cluster graph $G^\circ[W_{\mathcal{F}}]$. If it is empty, Q has at most one edge. If both $p, p' \in K$, they are adjacent and Q is a triangle. If exactly one is in K, the other is a non-neighbour of y and of the first (different clusters), so Q has one edge. If neither is in K, Q has at most one edge. In no case is Q an induced P_3. If $q \leq 1$, every pair of Q has at most one end in $W_{\mathcal{F}}$, so Q is an induced P_3 of G with the same pairs and, H being pair-maximal, has a pair in pairs(H); this pair is not in pairs$(\mathcal{F})$ (those lie inside $W_{\mathcal{F}}$), so it is soft with respect to H°. □

Theorem 5.10 (Component reduction)**.** *Let H be* any *packing, $\bar{\ell} = \ell - \sum_j (\tau_j - p_j)$, and apply Rule 6.2 to every reducible tight component (found by Proposition 5.8). Then, modulo the cited soundness of Rule 6.2 (Lemma 6.3 of [15]), the reduced instance is equivalent to (G, H, ℓ), and if it is a YES instance then $\bar{\ell} \geq 0$ and the number r of components of I of the remaining packing satisfies $r \leq \ell + \bar{\ell} \leq 2\ell$. In particular* if more than 2ℓ components remain, the answer is NO*. No τ_j of a non-tight component is needed by the algorithm.*

Proof. The rule. The set S_j used for a reducible tight component $\mathcal{C}_j$ consists of pairs inside W_j. So $G[W_i]$ and the sets $N(y) \cap W_i$ $(i \neq j)$ do not change, τ_i, tightness and (ir)reducibility of the other components are unchanged, ℓ and $\bar{\ell}$ are unchanged (Lemma 5.4), and one pass reaches an instance with no reducible tight component.

Counting. Work in the reduced instance. Let $\mathcal{A}$ be the set of tight (irreducible) components and $\mathcal{B}$ the set of non-tight ones. Since $\sum_j \tau_j = |H| + \sum_j (\tau_j - p_j)$, we have $|H| + \ell = \sum_j \tau_j + \bar{\ell}$. If $\bar{\ell} < 0$ the instance is a NO instance (Lemma 5.3). Otherwise let S be an optimal solution with $|S| \leq \sum_j \tau_j + \bar{\ell}$. By Lemma 5.6 with $J = \mathcal{A}$, $|\mathcal{A}| \leq 2\bar{\ell}$. Each $j \in \mathcal{B}$ has $\tau_j - p_j \geq 1$, so $|\mathcal{B}| \leq \sum_j (\tau_j - p_j) = \ell - \bar{\ell}$. Therefore $r \leq 2\bar{\ell} + \ell - \bar{\ell} = \ell + \bar{\ell} \leq 2\ell$ (as $\bar{\ell} \leq \ell$). □

The theorem is phrased through the number of remaining components so that it does not need the τ_j of non-tight components, which are not computable in polynomial time for a large component.

Corollary 5.11. *(a) For $\ell = 0$ a YES instance is reduced to the empty packing; this agrees with Theorem 3.4. (b) For an arbitrary packing H, compute M and $H' = H \cup M$ (Lemma 2.4) and then reduce tight reducible components repeatedly. This ends, in polynomial time, with a pair-maximal packing H^*, a budget $\ell^* = \ell - |M| \leq \ell$, $\mathrm{tw}(I(H^*)) \leq w + |M| \leq w + \ell$, and at most $2\ell^*$ components of the intersection graph, or with a correct NO answer.*

Proof. (a) $r \leq 2\ell = 0$. (b) After the single maximalisation H' is pair-maximal and $\mathrm{tw}(I(H')) \leq w + |M|$ (Lemma 2.4). Each reduction step removes the members of a tight reducible component, keeps the budget (Lemma 5.9), keeps the packing pair-maximal and does not increase the treewidth

of the intersection graph (it is an induced subgraph of the previous one). The loop ends after at most $|H'|$ steps, when no tight component is reducible; Theorem 5.10 bounds the number of components by $2\ell^*$. □

What remains after the reduction is the case of *few but arbitrarily large* components of $I(H)$ (for $w = 1$, a single tree of members), which is Open Problem 4.

6 Optimal solutions of pair-maximal packings

Setting. Throughout this section H is pair-maximal (Section 2; for an arbitrary packing read the primed instance), $D \geq \max_v |Q(v)|$ is a number, and S is a solution with $|S| \leq |H| + \ell$, so that its set U of hard edits has $u = |U| \leq \ell$ (Lemma 2.2); U^- and U^+ are the deleted hard edges (hard edges across clusters) and the inserted hard non-edges (hard non-edges inside clusters), and $\mathrm{del}_x, \mathrm{ins}_x$ count the elements of U^- and U^+ at the vertex x. Solutions and partitions correspond (Section 2). A *hard component* is a connected component of the graph $(V, \text{hard edges of } G)$; it is *clean* if all its non-adjacent pairs are soft.

Lemma 6.1 (Cost formula)**.** *Let π be a partition of V and $f(C) = \binom{|C|}{2} - 2e(C)$, where $e(C)$ is the number of edges of G inside C. Then* $\mathrm{cost}(\pi) = |E| + \sum_{C \in \pi} f(C)$.

Proof. The cost counts non-edges inside blocks, $\sum_C [\binom{|C|}{2} - e(C)]$, plus edges across blocks, $|E| - \sum_C e(C)$. □

6.1 Clusters of optimal solutions: half-degree and diameter

Lemma 6.2 (Half-degree and diameter)**.** *Let S be an optimal solution, C a cluster of $G \triangle S$ and $v \in C$. Put $d_C(v) = |N_G(v) \cap C|$.*

(a) $d_C(v) \geq (|C| - 1)/2$.

(b) If $a, b \in C$ and $ab \notin E$, then some $z \in C \setminus \{a, b\}$ has $az, zb \in E$.

This is the single-vertex-move argument behind the critical-clique property of Cluster Editing [10, 7]; we include the short proof.

Proof. (a) Let $S' = S \triangle \{vx : x \in C \setminus \{v\}\}$. Then $G \triangle S'$ is $G \triangle S$ with v cut off from the rest of C, a cluster graph. For $x \in C \setminus \{v\}$ with $vx \in E$ the pair $vx \notin S$ and $vx \in S'$: this adds $d_C(v)$ pairs. For $x \in C \setminus \{v\}$ with $vx \notin E$ the pair is in S and not in S': this removes $|C| - 1 - d_C(v)$ pairs. All other pairs are unchanged. So $|S'| = |S| + 2d_C(v) - (|C| - 1)$, and $|S'| \geq |S|$ by optimality.

(b) $N_G(a) \cap C$ and $N_G(b) \cap C$ are subsets of $C \setminus \{a, b\}$ (as $a \notin N(a)$ and $ab \notin E$), a set with $|C| - 2$ elements, while $d_C(a) + d_C(b) \geq |C| - 1$ by (a). So the two sets intersect. □

Lemma 6.2 uses neither H nor ℓ; only local optimality under single-vertex moves.

6.2 Hard pairs

Lemma 6.3 (HE and H2)**.** *Let H be pair-maximal. (a) If ab, bc are hard edges (a, b, c distinct), then $ac \in E$ or ac is soft. (b) If uv is a hard non-edge, then $N_h(u) \cap N_h(v) = \emptyset$.*

Proof. (a) If $ac \notin E$ and ac were hard, then $a\,b\,c$ would be an induced P_3 with all three pairs hard, contradicting pair-maximality. (b) If $z \in N_h(u) \cap N_h(v)$, then $u\,z\,v$ is an induced P_3 ($uz, zv \in E$, $uv \notin E$) all of whose pairs are hard. □

No YES hypothesis is needed. Without pair-maximality the lemma fails ($H = \emptyset$ and any induced P_3).

Lemma 6.4 (EX: bounds at a hard edit)**.** *Let H be pair-maximal, S a solution with hard edits U, $u = |U|$, let $xy \in U$ and let C be the cluster of x. Then:*

(a) if xy is deleted, $d_h(x) \le D + \mathrm{del}_x + \mathrm{del}_y - 1$; if xy is inserted, $d_h(x) \le D + \mathrm{ins}_y - 1 + \mathrm{del}_x$. In both cases $d_h(x) \le D + u$;

(b) $|C| \le 1 + D + d_h(x) + \mathrm{ins}_x$, hence $|C| \le 1 + 2D + u$;

(c) every $z \in C$ has $d_h(z) \le |C| - 1 + \mathrm{del}_z \le 2D + 2u$.

Proof. (a) *Deleted:* $x \in C$, $y \notin C$. A hard neighbour z of x with xz not deleted ("kept") lies in C, and $z \neq y$. Apply Lemma 6.3(a) with centre x to the hard edges xz, xy: $zy \in E$ or zy is soft. If zy is a hard edge, it joins $z \in C$ to $y \notin C$, so it is a deleted hard edge at y other than xy (there are $\mathrm{del}_y - 1$ of these); otherwise $z \in Q(y)$, at most D vertices. So at most $D + \mathrm{del}_y - 1$ hard neighbours are kept, and $d_h(x) \le D + \mathrm{del}_y - 1 + \mathrm{del}_x$. *Inserted:* $x, y \in C$. A kept hard neighbour z of x lies in C and $z \neq y$ (as $xy \notin E$); zy is a pair inside C, hence an edge of $G \triangle S$. If zy were a hard edge, $z \in N_h(x) \cap N_h(y)$, contradicting Lemma 6.3. So zy is soft ($z \in Q(y)$, at most D choices) or a hard non-edge, inserted at y and different from xy ($\mathrm{ins}_y - 1$ choices). So $d_h(x) \le D + \mathrm{ins}_y - 1 + \mathrm{del}_x$. In the deleted case the pairs counted by $\mathrm{del}_x + \mathrm{del}_y - 1$ are distinct pairs of U, so the sum is $\le u$; in the inserted case $\mathrm{ins}_y - 1 + \mathrm{del}_x \le u - 1$ (an inserted pair at y other than xy and a deleted pair at x are distinct elements of $U \setminus \{xy\}$).

(b) For $z \in C \setminus \{x\}$ the pair xz is an edge of $G \triangle S$. Either xz is soft (at most D such z), or a kept hard edge (at most $d_h(x)$), or a hard non-edge, inserted at x (at most ins_x). With (a): in the deleted case $|C| \le 1 + D + (D + \mathrm{del}_x + \mathrm{del}_y - 1) + \mathrm{ins}_x \le 1 + 2D + u$ because the deleted pairs at x or y and the inserted pairs at x are distinct elements of U; in the inserted case $\mathrm{ins}_x + \mathrm{ins}_y - 1$ counts the inserted pairs at x or y and del_x is disjoint from them, so the total is again $\le u$.

(c) Hard neighbours of z inside C are at most $|C| - 1$; hard neighbours outside C are joined by deleted edges, at most $\mathrm{del}_z \le u$. Use (b). □

Lemma 6.5 (Connected clusters)**.** *In an optimal solution every cluster C induces a connected subgraph $G[C]$.*

Proof. If $C = C_1 \cup C_2$ with $C_1, C_2 \neq \emptyset$ and no edge of G between them, splitting C into C_1, C_2 turns the $|C_1||C_2| > 0$ inserted pairs between them into non-edges outside S and creates no deletion. □

6.3 Normal form and candidates

Fix ℓ and put

$$B = 2D + 3\ell, \qquad L_B = \{v : \ d_h(v) \le B\}, \qquad Z = B + D + 1 = 3D + 3\ell + 1.$$

In an optimal solution of size at most $|H| + \ell$ call a cluster *exceptional* if it contains an endpoint of a pair of U. Let BIG be the union of the hard components that contain a vertex v with $d_h(v) > B$.

Lemma 6.6 (NF: normal form)**.** *Let H be pair-maximal and S an optimal solution with $|S| \le |H| + \ell$, so $u \le \ell$.*

(α) There are at most $2u \le 2\ell$ exceptional clusters. Each exceptional cluster C has $|C| \le 1 + 2D + u \le 2D + \ell + 1$, lies in L_B, and $G[C]$ is connected.

(β) Let C be a non-exceptional cluster. Then no hard edge leaves C and no hard non-edge lies in C; so C is a union of whole hard components, each clean, and all pairs between different components inside C are soft. If C contains at least two components, $|C| \le 2D$.

(γ) Every hard component K of BIG *is a whole cluster of S, and this cluster is non-exceptional. Consequently every exceptional cluster is disjoint from* BIG*, and every component of* BIG *is clean.*

Proof. (α) Each pair of U has at most two clusters at its ends, which gives the count. The bound on $|C|$ is Lemma 6.4(b) (for an endpoint $x \in C$ of a pair in U), the inclusion $C \subseteq L_B$ is Lemma 6.4(c) together with $2D + 2u \le B$, and connectivity is Lemma 6.5.

(β) A hard edge leaving C would be deleted, with an endpoint in C; a hard non-edge inside C would be inserted. Both would make C exceptional. If K is a hard component meeting C, hard edges at a vertex of $K \cap C$ do not leave C, so $K \subseteq C$ by connectivity of K; K is clean since C contains no hard non-edge. Two vertices of different components inside C are not joined by a hard edge, and are not a hard non-edge, so their pair is soft. If C contains components $K_1 \neq K_2$, a vertex $x \in K_1$ has all vertices of $C \setminus K_1$ as soft partners, so $|C \setminus K_1| \le D$; likewise $|C \setminus K_2| \le D$; since $K_1 \subseteq C \setminus K_2$, $|C| = |K_1| + |C \setminus K_1| \le 2D$.

(γ) Let $d_h(v) > B$ and let $K = K(v)$ be its hard component. By Lemma 6.4(a) no endpoint of a pair of U has hard degree above $D + u \le B$, and by (α) no vertex of an exceptional cluster has hard degree above B. So the cluster C of v is non-exceptional, and by (β) $K \subseteq C$ and C is a union of components. Now $|K| \ge d_h(v) + 1 > B + 1 > 2D$, so by ($\beta$) C cannot contain two components; hence $C = K$. Every other vertex of K lies in the same cluster. Cleanness is (β). The disjointness statement follows since a component of BIG is a non-exceptional cluster. □

Corollary 6.7 (CAND: bounded candidate partners)**.** *Let H be pair-maximal, S optimal with $|S| \le |H| + \ell$, and $xy \in U$. Then $x, y \in L_B$ and:*

(i) if xy is deleted, $y \in N_h(x)$, and $|N_h(x)| \le D + \ell$;

(ii) if xy is inserted, x and y have a common neighbour in their cluster, so their distance in $G[L_B]$ is exactly 2;

(iii) in both cases y lies in the ball $\mathrm{Ball}(x, 2)$ *of $G[L_B]$, which has at most Z^2 vertices.*

The same holds with the roles of x and y exchanged.

Proof. $x, y \in L_B$ by Lemma 6.4(a) ($d_h \le D + u \le B$). (i) is the definition together with Lemma 6.4(a). (ii) x, y lie in the exceptional cluster C and are non-adjacent, so Lemma 6.2(b) gives $z \in C$ adjacent to both; $C \subseteq L_B$ by Lemma 6.6(α), so the path $x\,z\,y$ lies in $G[L_B]$ and the distance is 2 (not 1 since $xy \notin E$). (iii) A vertex of $G[L_B]$ has at most B hard neighbours and at most D soft ones, so degree at most $B + D = Z - 1$, and a ball of radius 2 has at most $1 + (Z-1) + (Z-1)^2 \le Z^2$ vertices; deleted pairs have distance 1. □

Lemma 6.2(b) is what makes the radius 2 independent of D and ℓ: the candidates for the other end of a hard edit lie within distance 2 of one end in $G[L_B]$. Corollary 6.7 says that *one* endpoint of a hard edit determines a set of at most Z^2 candidates for the other, for every hard edit, including those of type T2 and deleted edges (Section 8). What it does not bound is the number of candidates for the first endpoint, which is n.

7 An n^ℓ algorithm and the limits of the cluster-identity route

This section uses Corollary 6.7 in two ways: guessing the positions of the hard edits (Theorem 7.2), and following the clusters directly (Theorem 7.3). The first halves the exponent of Theorem 3.5; the second is fixed-parameter tractable for a stronger parameter ω which, by Proposition 7.4, is not bounded by any function of (ℓ, w). **Whether CEaMP is fixed-parameter tractable in $\ell + w$ remains open.**

7.1 The n^ℓ algorithm

Remark 7.1 (Fixed set of hard edits). The proof of Theorem 3.5 shows: for a *fixed* set U' of hard pairs, $m_{U'} = \min \sum_i |x_i|$ over all $x \in \prod \mathcal{D}_i$ with $G \triangle (U' \cup \bigcup x_i)$ a cluster graph is computed in $7^{w+1} n^{O(1)}$ time from a tree decomposition of $I(H)$ of width w, and for *every* U' of hard pairs the value $m_{U'} + |U'|$ is the size of an actual solution.

Theorem 7.2 (n^ℓ algorithm)**.** *Given (G, H, ℓ) for an arbitrary packing H and a tree decomposition of $I(H)$ of width w, CEaMP is solvable in time*

$$(n+1)^\ell \, Z_*^{2\ell} \, 7^{\,w+\ell+1} \, n^{O(1)}, \qquad D_* = D(H) + 2\ell, \quad Z_* = 3D_* + 3\ell + 1 = 3D(H) + 9\ell + 1.$$

Since $D(H) \le 2(w+1)$, $Z_* \le 6w + 9\ell + 7$, so the factor $Z_*^{2\ell}$ is a function of (ℓ, w) only. This improves the exponent 2ℓ of Theorem 3.5 to ℓ; it is still XP in ℓ.

Proof. Compute M and $H' = H \cup M$ by Lemma 2.4; reject if $|M| > \ell$. Then H' is pair-maximal, $\ell' = \ell - |M| \le \ell$, $D(H') \le D(H) + 2|M| \le D_*$, a tree decomposition of $I(H')$ of width $w' \le w + |M| \le w + \ell$ is available, and the instance is equivalent to (G, H', ℓ'). All lemmas of Section 6 apply to H' with $D(H')$ and ℓ'; put B, L_B, Z accordingly. Then $Z = 3D(H') + 3\ell' + 1 \le Z_*$.

Algorithm. Compute L_B and, for $x \in L_B$, the set $P(x)$ of vertices $y \in \mathrm{Ball}_{G[L_B]}(x, 2) \setminus \{x\}$ with xy hard; $|P(x)| \le Z^2$. Enumerate all sequences of at most ℓ' pairs $(x_1, y_1), \ldots, (x_j, y_j)$ with $x_i \in L_B$ and $y_i \in P(x_i)$; there are at most $(1 + nZ^2)^{\ell'} \le (n+1)^{\ell'} Z^{2\ell'}$ of them. For each, let $U' = \{x_i y_i\}$ (a set of at most ℓ' hard pairs), compute $m_{U'}$ by Remark 7.1 for H' in $7^{w'+1} n^{O(1)}$ time, and accept iff $m_{U'} + |U'| \le |H'| + \ell'$.

Soundness. By Remark 7.1, $m_{U'} + |U'|$ is the size of an actual solution of G for every U' of hard pairs of H'; acceptance gives a solution of size $\le |H'| + \ell' = |H| + \ell$.

Completeness. Suppose the instance is a YES instance and let S be an optimal solution, $|S| \le |H'| + \ell'$, with hard edits U, $|U| \le \ell'$ (Lemma 2.2 for H'). For each pair of U take one endpoint x; it lies in L_B and the other endpoint lies in $P(x)$ by Corollary 6.7. So U is among the enumerated sets, and for it $m_U + |U| \le |S| \le |H'| + \ell'$ (the soft part of S is a feasible choice of x for U by Lemma 2.2).

Time. $(n+1)^{\ell'} Z^{2\ell'} 7^{w'+1} n^{O(1)} \le (n+1)^\ell Z_*^{2\ell} 7^{w+\ell+1} n^{O(1)}$. □

The exponent is halved because the candidate list is attached to *one* endpoint (Corollary 6.7); Theorem 3.5 guesses both. Note that ℓ bounds the *number* of hard edits but not the cost of locating them: the n^ℓ comes from the position of the first endpoint of each pair. A fixed-parameter algorithm along this route needs a bound on these positions as well.

7.2 A fixed-parameter algorithm for a stronger parameter

Let H be pair-maximal and $\ell, D, B, Z, L_B, \mathrm{BIG}$ as in Section 6. Let $V' = V \setminus \mathrm{BIG}$ and let Γ' be the graph on V' whose edges are all edges of G and all soft pairs with both ends in V'.

Theorem 7.3 (Fixed-parameter tractability in (ℓ, D, ω))**.** *Let $\omega = \mathrm{tw}(\Gamma')$ and assume a tree decomposition of Γ' of width ω is given. Then one can compute in time $(Z(\omega+1))^{O(\omega)}n + n^{O(1)}$ a number m^* such that*

$$\mathrm{OPT}(G) \le |H| + \ell \iff |E| + \sum_K f(K) + m^* \le |H| + \ell,$$

where K ranges over the hard components of BIG *and f is as in Lemma 6.1. The proof, with the dynamic program, is in Appendix C.*

All quantities in Theorem 7.3 refer to the pair-maximal instance (G, H', ℓ'); they are not functions of the input (G, H, ℓ) (Section 2).

7.3 Why ω is not bounded by (ℓ, w)

Proposition 7.4 (ω is unbounded at $\ell = 0$, $w = 0$)**.** *For every cubic graph F there is an instance $(G_F, H_F, 0)$ with H_F a vertex-disjoint pair-maximal packing ($w = 0$, $c = 1$, $D = 2$), $\mathrm{OPT}(G_F) = |H_F|$ (a YES instance with $\ell = 0$), $\mathrm{BIG} = \emptyset$, every hard component of size at most 6 and every hard degree at most 4, and $\mathrm{tw}(\Gamma') \ge \mathrm{tw}(F)$. By [9] (cited, not re-proved here) there are cubic graphs F (expanders) with $\mathrm{tw}(F) = \Omega(|V(F)|)$. Hence ω is not bounded by any function of (ℓ, w).*

Proof. Orient each edge of F arbitrarily; for each oriented edge $e = (p \to q)$ introduce three vertices a_e, b_e, c_e. Let $\mathcal{K}_v = \{a_e : e \text{ leaves } v\} \cup \{b_e, c_e : e \text{ enters } v\}$ and make each $\mathcal{K}_v$ a clique; add the cross edges $a_e b_e$ and nothing else. Then $|\mathcal{K}_v| = \mathrm{out}(v) + 2\,\mathrm{in}(v) = 3 + \mathrm{in}(v) \le 6$. $H_F = \{P_e = (a_e, b_e, c_e)\}$: $a_e b_e, b_e c_e \in E$ and $a_e c_e \notin E$ since $a_e \in \mathcal{K}_p$, $c_e \in \mathcal{K}_q$, $p \ne q$ and cross edges are of the form $a_{e'} b_{e'}$; so P_e is an induced P_3. The members are vertex-disjoint, so modification-disjoint, $c = 1$, $w = 0$ and $D = 2$.

Pair-maximality. In an induced P_3 $x\,y\,z$ not both edges can lie inside cliques, since then x, z would be in the clique of y (each vertex is in exactly one clique) and adjacent. So one of xy, yz is a cross edge $a_e b_e$, which is soft.

$\mathrm{OPT} = |H_F|$. The partition $\{\mathcal{K}_v\}$ has no inserted pair and deletes exactly the cross edges, $|H_F|$ of them; and $\mathrm{OPT} \ge |H_F|$ by Lemma 2.2.

Hard structure. The soft pairs inside $\mathcal{K}_v$ are $b_e c_e$ (e entering v); all other pairs inside $\mathcal{K}_v$ are hard edges. So each $\mathcal{K}_v$ is one hard component and it is clean. The hard degree of $a_e \in \mathcal{K}_p$ is $|\mathcal{K}_p| - 1 \le 4$ (as a_e is an out-edge, $\mathrm{in}(p) \le 2$, $|\mathcal{K}_p| \le 5$); that of b_e or $c_e \in \mathcal{K}_q$ is $|\mathcal{K}_q| - 2 \le 4$. So $d_h \le 4 = 2D + 3\ell$ at $\ell = 0$, hence no vertex has $d_h > B$ and $\mathrm{BIG} = \emptyset$.

Minor. $V' = V$, each $\mathcal{K}_v$ is connected in Γ' (a clique), and an edge or soft pair of Γ' between $\mathcal{K}_p \ne \mathcal{K}_q$ is one of $a_e b_e$, $a_e c_e$, which exist only for the oriented edge $e = (p \to q)$ of F. Contracting the $\mathcal{K}_v$ gives exactly F, so F is a minor of Γ' and $\mathrm{tw}(\Gamma') \ge \mathrm{tw}(F)$. □

Remark 7.5 (Scope of Proposition 7.4). The proposition refutes only the claim "$\omega \le f(\ell, w)$"; it does *not* refute fixed-parameter tractability. The family has $w = 0$ and $\ell = 0$; it is solved in polynomial time by Theorem 3.5 and has $s = 3$ in Theorem 5.1. The obstruction is to the representation by Γ' (and by any single decomposition used both for the soft pairs and for the clusters): hard edges between far vertices are free (not in the budget), arbitrary in number, and make footprints of clusters disconnected in any decomposition of $I(H)$ or of G_{soft}. At $\ell = 0$ they are harmless because the CSP of Theorem 3.4 needs no identities; for $\ell > 0$ an exceptional cluster can be assembled from vertices that are far apart (a $c = 1$ instance with two vertex-disjoint packed P_3's joined by one hard edge that is deleted). So a dynamic program on one decomposition cannot be used for both purposes without a preprocessing step; a reduction lemma that would supply it is Open Problem 3.

8 Locality of inserted pairs and a non-local obstruction

Section 6 bounds the *number* of hard edits and the *candidates for the second endpoint* of a hard edit, but not the number of candidate *positions* of the first. This section studies the inserted hard pairs of an *optimal* solution. The main tool is the diameter bound of Lemma 6.2, which, combined with pair-maximality, classifies every inserted pair into one of two types (Theorem 8.2). Pairs of the first type are local in a decomposition of $I(H)$ (Corollary 8.3, Proposition 8.5); the second type, which includes the generic merge of two pieces, is *not* local in the same sense (Proposition 8.7). Finally, deleted hard edges can join packed P_3's at arbitrary distance in $I(H)$ (Proposition 8.8). **Nothing in this section gives a fixed-parameter algorithm in $\ell + w$ or a hardness result.**

Setting. H is pair-maximal (for an arbitrary packing read the primed instance), $D(H) = 2c$, S is an *optimal* solution with $|S| \le |H| + \ell$, C is a cluster of $G \triangle S$, and $U = U^- \cup U^+$, $u = |U| \le \ell$ are as in Section 6. An *inserted pair* is a pair of U^+: a hard non-edge of G inside one cluster. A hard edge of G inside a cluster is an edge of $G \triangle S$ and not in S, hence *kept*. The *pieces* of S are the connected components of the graph (V, E_{keep}) where E_{keep} is the set of kept hard edges (isolated vertices are singleton pieces); every piece lies inside one cluster. Pieces depend on S, not only on (G, H).

Lemma 8.1 (Cross pairs)**.** *If $A \neq B$ are pieces contained in a cluster C, then every pair ab ($a \in A, b \in B$) is soft or inserted. This holds for every solution S; neither optimality nor pair-maximality is used.*

Proof. Since $a, b \in C$, ab is an edge of $G \triangle S$, i.e. $ab \in E$ xor $ab \in S$. Suppose ab is hard. If $ab \in E$ then $ab \notin S$, so ab is a kept hard edge and a, b lie in one piece, a contradiction. So $ab \notin E$, hence $ab \in S$, and as ab is hard, $ab \in U^+$. □

(Appendix B bounds the sizes of pieces that share a cluster.)

8.1 The T1/T2 dichotomy

For an inserted pair ab with $a, b \in C$, a *witness* is a vertex $z \in C \setminus \{a, b\}$ with $az, zb \in E$.

Theorem 8.2 (T1/T2)**.** *Let H be pair-maximal, S optimal and ab an inserted pair, $a, b \in C$. Then ab has a witness, and for every witness z at least one of az, zb is soft. Moreover:*

(T1) If az and zb are both soft, they are owned by two distinct *members $P_a \neq P_b$ of H, and $z \in V(P_a) \cap V(P_b)$.*

(T2) If az is hard (the case of zb hard is symmetric), then az is a kept hard edge, so a and z lie in one piece, and zb is soft.

We say that ab is of type T1 *if some witness has both steps soft, and of* type T2 *otherwise; then every witness has exactly one hard step.*

The type is a property of the pair ab (not of a chosen witness): it is T1 iff a witness with two soft steps exists.

Proof. A witness exists by Lemma 6.2(b). If az and zb were both hard for a witness z, then $a\,z\,b$ would be an induced P_3 with all three pairs hard (ab is hard by assumption), contradicting pair-maximality. If both are soft, let P_a, P_b own az, zb. If $P_a = P_b$ then $a, b \in V(P_a)$ and ab is soft, contradicting that ab is hard; so $P_a \neq P_b$, and both contain z. If az is hard, it is a hard edge inside C, hence kept. □

Corollary 8.3 (Locality of soft-free merges)**.** *Let H be pair-maximal, S optimal, and let $A \neq B$ be pieces inside one cluster C such that* no soft pair joins A and B. *Then for every $(a, b) \in A \times B$:*

(i) ab is an inserted pair, of type T1;

(ii) there is $z \in C \setminus (A \cup B)$ with az, zb soft, owned by distinct members $P_a \neq P_b$ with $z \in V(P_a) \cap V(P_b)$;

(iii) given a, the number of possible partners b in any such pair is at most $D(H)(D(H) - 2) = 4c(c - 1)$.

In particular a and b are at distance exactly 2 in the soft graph G_{soft}, and $A \cup B$ lies in the ball of radius 4 of G_{soft} around any of its vertices.

Proof. (i) The cross pair ab is not soft by hypothesis, so it is hard, hence inserted by Lemma 8.1; its type is T1 by (ii). (ii) Apply Theorem 8.2 to ab with a witness z. Suppose az is hard (the other case is symmetric); then z lies in the piece A of a. The pair zb is an edge of G and joins A to B: if it were hard it would be kept and z, b would share a piece, contradicting $A \neq B$; if soft it contradicts the hypothesis. So az and zb are both soft, which is (T1). If $z \in A$, then zb (soft) is a soft pair between A and B; similarly for $z \in B$ and az. So $z \notin A \cup B$. (iii) Fix a. By (ii) $z \in Q(a)$: at most $D(H)$ choices. For such z, let P_a own az. The vertex b satisfies bz soft owned by $P_b \neq P_a$, and $b \notin V(P_a)$ (otherwise $ab \in \text{pairs}(P_a)$ is soft). Now $Q(z) \cap V(P_a) = V(P_a) \setminus \{z\}$ has two elements, so $b \in Q(z) \setminus V(P_a)$, a set of at most $|Q(z)| - 2 \leq D(H) - 2$ vertices. Altogether at most $D(H)(D(H) - 2) = 2c(2c - 2) = 4c(c - 1)$ partners. (For $c = 1$ the count is 0: (T1) needs two members through z.)

The distance statement: ab is not soft and a, z, b is a soft path; so $d_{G_{\text{soft}}}(a, b) = 2$. Every $a' \in A$ has $d_{G_{\text{soft}}}(a', b) = 2$ for a fixed $b \in B$, and every $b' \in B$ has $d_{G_{\text{soft}}}(b', a') = 2$; hence all of $A \cup B$ is within distance 4 of b (and of any other vertex of $A \cup B$, by symmetry). □

Remark 8.4 (Scope). If no soft pair joins A and B then all $|A||B|$ cross pairs are inserted (Lemma 8.1), so $|A||B| \leq u \leq \ell$. The corollary covers *only* merges without a soft pair between the pieces. Merges in which some cross pairs are soft (the generic case when the pieces are neighbours in the soft graph) are not covered, and Proposition 8.7 shows they can genuinely be of type T2.

Proposition 8.5 (Bag locality of type T1 pairs)**.** *Let $(\mathcal{T}, \{\beta_t\})$ be a tree decomposition of $I(H)$ of width w, let $c = c(H)$, and for a set Y of nodes of $I(H)$ let $N[Y]$ be the set of nodes at distance at most 1 from Y (a set of* members of H*).*

(a) If ab is an inserted pair of type T1 with a witness z and members P_a, P_b as in Theorem 8.2, then P_a, P_b are adjacent in $I(H)$, so some bag β_t contains both and $a, b \in \bigcup_{P \in \beta_t} V(P)$, a set of at most $3(w + 1)$ vertices. Moreover the member through z and every member through a or b lies in $N[\beta_t]$.

(b) $(\mathcal{T}, \{N[\beta_t]\})$ is a tree decomposition of the square $I(H)^2$ of width at most $(w+1)(3c-2)-1 \leq (w + 1)(3w + 1) - 1$.

Proof. (a) P_a, P_b share z, so they are adjacent in $I(H)$ and, the edge being covered, lie in a common bag. Every member through a contains $a \in V(P_a)$, hence meets $P_a \in \beta_t$ and lies in $N[\beta_t]$; similarly for b and z.

(b) *Size.* Each member has 3 vertices, each lying in at most c members, so $I(H)$ has maximum degree at most $3(c-1)$, $|N[\beta_t]| \leq (w+1)(1+3(c-1)) = (w+1)(3c-2)$, and $c \leq w+1$. *Connectivity.*

For a node v let $\mathcal{T}_v = \{t : v \in \beta_t\}$ (a subtree). The nodes t with $v \in N[\beta_t]$ form $\bigcup_{y \in N[v]} \mathcal{T}_y$. Each $\mathcal{T}_y$, $y \in N(v)$, meets $\mathcal{T}_v$ because the edge yv lies in some bag; a union of subtrees that all meet the connected set $\mathcal{T}_v$ is connected. *Edges.* Let yv be an edge of $I(H)^2$. If y, v are adjacent in $I(H)$, any $t \in \mathcal{T}_y$ has $y \in \beta_t$ and $v \in N[y]$. If they have a common neighbour m, any $t \in \mathcal{T}_m$ has $y, v \in N[m] \subseteq N[\beta_t]$. □

A type T1 pair is therefore a pair of vertices inside one bag of a decomposition of bounded width, and, given one end, it has at most $4c(c-1)$ candidate partners in the soft-free case. In particular each *cross pair* (a, b) of a soft-free merge lies in a bag; this does not put the pieces A, B themselves in one bag.

Lemma 8.6 (Near-twin agreement)**.** *Let S be any solution (optimal or not), $u = |U|$, and let $z \neq b$ lie in one cluster of $G \triangle S$. Then*

$$A(z, b) = \big\{a \notin \{z, b\} : \ az, ab \text{ both hard}, \ [az \in E] \neq [ab \in E]\big\}$$

has at most u elements. In particular, if zb is soft, the set $\Theta(z, b) = N_h(z) \setminus (N[b] \cup Q(b))$ satisfies $|\Theta(z, b)| \leq u \leq |S| - |H|$ and is computable from G and H alone.

Proof. Let $a \in A(z, b)$. Exactly one of az, ab is a hard edge and the other a hard non-edge. If a lies in the cluster of z, b, the hard non-edge is a hard non-edge inside a cluster, so it is in U. If a lies in another cluster, the hard edge joins two different clusters, so it is deleted and in U. Assign to a this pair. Its ends are a and one of z, b; since $a \notin \{z, b\}$, the map $a \mapsto$ pair is injective. So $|A(z, b)| \leq u$. For the second statement, $a \in \Theta(z, b)$ means $a \neq z$, az a hard edge, $a \notin N[b]$ (so $a \neq b$ and $ab \notin E$) and ab not soft. Thus $a \in A(z, b)$. □

In a type T2 pair with az hard and zb soft, $a \in \Theta(z, b)$: the far end a is one of at most ℓ vertices determined by the soft pair (z, b). The lemma does not say where a is relative to z in $I(H)$: a is a hard neighbour of z.

8.2 Merge pointers need not be bag-local

Proposition 8.7 (Examples of type T2 merges)**.** *Each of the following instances (graph G on vertices $0, \ldots, n-1$ given by its edge list, H given by triples (x, y, z) with centre y) has a pair-maximal modification-disjoint packing H and a unique optimal solution, found by enumerating all set partitions of the vertex set.*

(a) $n = 7$, $E = \{03, 04, 12, 13, 14, 15, 23, 35, 36\}$, $H = \{(4, 1, 5), (0, 3, 2), (1, 3, 6)\}$. $\mathrm{OPT} = 4 = |H| + 1$*; the clusters are* $\{0, 4\}, \{1, 2, 3, 5\}, \{6\}$*. The pieces of* $\{1, 2, 3, 5\}$ *are* $\{1, 2\}$ *and* $\{3, 5\}$ *(kept hard edges* 12 *and* 35*); the cross pairs* $13, 15, 23$ *are soft and* 25 *is inserted.* $Q(2) = \{0, 3\}$ *and* $Q(5) = \{4, 1\}$ *are disjoint, so the pair* 25 *has no witness with two soft steps: it is of type T2.*

(b) $n = 8$, $E = \{02, 06, 14, 16, 17, 23, 25, 37, 45, 46, 47, 57\}$, $H = \{(0, 6, 4), (3, 7, 4), (2, 5, 4), (0, 2, 3), (6, 1, 7)\}$. $\mathrm{OPT} = 6 = |H| + 1$*; the clusters are* $\{0, 6\}, \{1, 4, 5, 7\}, \{2, 3\}$*; the only hard edit of the solution is the inserted pair* 15*; the pieces are* $\{1, 4\}$ *and* $\{5, 7\}$*, the cross pairs* $17, 45, 47$ *are soft,* $Q(1) = \{6, 7\}$*,* $Q(5) = \{2, 4\}$*, so* 15 *has no witness with two soft steps and is of type T2.*

(c) $n = 6$, $E = \{02, 03, 04, 05, 12, 13, 15, 25, 35, 45\}$, $H = \{(3, 5, 4), (0, 2, 1)\}$*. The unique optimum has clusters* $\{0, 1, 2, 3, 5\}, \{4\}$ *and* $\mathrm{OPT} = 4$*; it contains the inserted pair* 23 *(type T2:* 30 *hard and kept,* 02 *soft), but* 2 *and* 3 *lie in the* same *piece (the kept hard edges* $03, 05, 13, 15, 25$ *connect* $\{0, 1, 2, 3, 5\}$*).*

Consequently "every inserted pair with both ends in $V(H)$ has a common soft neighbour" is false, and Corollary 8.3 cannot be extended to merges of two pieces that are joined by a soft pair: (a) and (b) are such merges, with inserted pairs of type T2. Example (c) is T2 but intra-piece.

Proof. Direct enumeration (all 877, 4140 and 203 set partitions of 7, 8 and 6 vertices respectively) of the cost $|\{ab : (ab \in E) \neq (a, b \text{ in one block})\}|$, together with a check that each triple of H is an induced P_3, that the triples have pairwise disjoint pair sets and that no induced P_3 of G has all pairs hard. The structural claims (pieces, soft pairs, Q) are read off from the lists above. In (a) and (b), every witness z of the inserted pair has az or zb hard because $Q(a) \cap Q(b) = \emptyset$, so the pair is not of type T1. □

8.3 An obstruction: deleted hard edges between far members

An optimum may be forced to delete a hard edge whose ends lie in members that are far apart in $I(H)$.

Proposition 8.8 (FAR). *For $m \geq 3$ let G_m be the path $0, 1, \ldots, 2m$ with the extra edge $e^* = \{1, 2m-1\}$ and $H_m = \{P_i = (2i, 2i+1, 2i+2) : 0 \leq i < m\}$ (centres $2i+1$). Then:*

(i) H_m is a pair-maximal modification-disjoint packing, $I(H_m)$ is a path on m nodes ($c = 2$, $w = 1$);

(ii) $\mathrm{OPT}(G_m) = m + 1 = |H_m| + 1$;

(iii) in every *optimum the hard edge e^* is deleted. Its ends lie in P_0 and P_{m-1}, at distance $m-1$ in $I(H_m)$.*

Proof. (i) The chord joins 1 and $2m-1$, which are $2m-2 \geq 4$ apart, so no triple P_i contains both of them and each P_i is an induced P_3; the pair sets are disjoint. Every path edge lies in some P_i and is soft, so e^* is the only hard edge. An induced P_3 has two edges, at most one of which is e^*, so it contains a soft pair: H_m is pair-maximal.

(ii) $|E| = 2m+1$ and G_m is connected with $2m+1$ vertices, hence unicyclic (the unique cycle is on $1, \ldots, 2m-1$, of length ≥ 5). An induced subgraph on s vertices has $e \leq s-1$ edges unless it contains the cycle, in which case $s \geq 5$ and $e \leq s$. With $f(C) = \binom{|C|}{2} - 2e(C)$ this gives $f(C) \geq (s-1)(s-4)/2 \geq -1$ in the first case, with equality iff $s \in \{2, 3\}$ and $G[C]$ is connected, and $f(C) \geq s(s-5)/2 \geq 0$ in the second. Clusters with $f = -1$ have ≥ 2 vertices, so there are at most m of them, and by Lemma 6.1 cost $\geq (2m+1) - m = m+1$. The partition $\{0,1\}, \{2,3\}, \ldots, \{2m-2, 2m-1\}, \{2m\}$ has cost $m+1$.

(iii) In an optimum $\sum f = -m$, so exactly m clusters have $f = -1$ (each connected, of size 2 or 3) and all others have $f = 0$; at most one vertex lies outside the $f = -1$ clusters. A cluster with $f = -1$ that contains the pendant vertex 0 contains its only neighbour 1; likewise for $2m$ and $2m-1$. Suppose $1, 2m-1$ lie in one cluster C^*. If $f(C^*) = -1$ and $|C^*| = 2$, then $C^* = \{1, 2m-1\}$ contains no pendant, and both pendants are outside the $f = -1$ clusters: two vertices outside, a contradiction. If $f(C^*) = -1$ and $|C^*| = 3$, it contains at most one pendant; any pendant not in C^* has its only neighbour in C^* and lies in no other $f = -1$ cluster, so exactly one pendant is outside and the remaining $2m-3$ vertices must form $m-1$ clusters of size ≥ 2, impossible. If $f(C^*) \geq 0$, at most $2m-1$ vertices remain for m clusters of size ≥ 2. So 1 and $2m-1$ are in different clusters, i.e. e^* is deleted. □

So the pointer realised by the deleted edge e^* joins members at distance $m-1$ in $I(H_m)$: its two ends are not adjacent in any $I(H)^r$ with $r < m-1$, and a decomposition of $I(H)^r$ has no

reason to put them in one bag. Corollary 6.7 bounds the candidates for the *other* end of e^* (at most $D+\ell$ hard neighbours), not the position of the first. For $m = 3, 4, 5$ a computer enumeration gives $\mathrm{OPT} = m + 1$ with $7, 9, 11$ optima, each separating the ends of e^*. Appendix D shows that the idea of putting "locally bad" members into every bag does not repair this for distances 2 and 3 (Proposition D.10); for unbounded distance this is open.

9 Open problems and summary of results

Notation as above; $w = \mathrm{tw}(I(H))$.

Open Problem 1 (Main question: (ℓ, w) in general; **OPEN**)**.** Is CEaMP fixed-parameter tractable parameterized by $\ell + w$, i.e. solvable in $f(\ell, w)n^{O(1)}$ time, or is it W[1]-hard in $\ell + w$?

Known: XP in ℓ with single-exponential dependence on w, in time $7^{w+1}n^{2\ell+O(1)}$ and $n^{\ell}f(\ell, w)7^{w+\ell+1}$ (Theorems 3.5 and 7.2); fixed-parameter tractable in ℓ when $w = 0$ [15], and in (ℓ, s) when the components of $I(H)$ span at most s vertices (Theorem 5.1); fixed-parameter tractable in (ℓ, D, ω) (Theorem 7.3), where ω is not bounded by (ℓ, w) (Proposition 7.4); the case $H = \emptyset$ (plain Cluster Editing, $\ell = k$) is fixed-parameter tractable. Adding c to the parameter changes nothing, since $c \leq w + 1$. On our route, a positive answer needs a bound $g(\ell, w)$ on the number of candidate *positions* of the first endpoint of each hard edit: given one endpoint, the other has at most Z^2 candidates (Corollary 6.7), so only the placement of at most ℓ bounded objects is free. Locality is proved for type T1 pairs, in particular soft-free merges (Corollary 8.3), but not for type T2 pairs or deleted edges (Propositions 8.7 and 8.8). The algorithm need not enumerate positions, so this is not a necessary condition.

Open Problem 2 (Pointers of type T2 and deleted hard edges)**.** Can the far ends of inserted pairs of type T2 (a kept hard edge az with a far from z in $I(H)$, and a soft pair zb; by Lemma 8.6 the end a is one of at most ℓ vertices fixed by (z, b)), and of deleted hard edges between far members, be handled in a dynamic program of width bounded by $g(\ell, w)$? Proposition 8.8 shows that an optimum can be forced to delete a hard edge between members at any distance in $I(H)$ (with $w = 1$, $c = 2$, $\ell = 1$); whether a merge pointer between two pieces joined by a soft pair can be forced at unbounded distance is open.

Open Problem 3 (Reduction lemma; open conjecture)**.** Is there a polynomial-time algorithm that, given (G, H, ℓ) with H pair-maximal, either outputs NO correctly or outputs an equivalent instance (G^*, H^*, ℓ^*) with H^* pair-maximal, $\ell^* \leq \ell$, $w(H^*) \leq f_1(\ell, w)$ and $\mathrm{tw}(\Gamma'(G^*, H^*)) \leq f_2(\ell, w)$? A positive answer would give fixed-parameter tractability in $\ell + w$ by Theorem 7.3 (with $D^* \leq 2(w(H^*) + 1)$). Proposition 7.4 shows that some reduction is necessary. For $c = 1$ the rule of [15] is such a reduction; Theorem 5.10 leaves at most 2ℓ components of $I(H)$; Appendix E shows that the same rule applied to pair-closed unions of members removes nothing on a family with $w = 1$, $\ell = 1$ and unbounded $\mathrm{tw}(\Gamma')$.

Open Problem 4 ($w = 1$ with unbounded tree components)**.** If $I(H)$ is a forest ($w \leq 1$, hence $c \leq 2$) with components of unbounded size, is CEaMP fixed-parameter tractable in ℓ? Theorem 5.1 settles bounded components, in particular perfect matchings (Corollary 5.2), but its counting argument is at the level of whole components and gives nothing for a single large tree component.

Open Problem 5 (Packings with $c \leq 22$)**.** Is CEaMP with $c = 2$ fixed-parameter tractable in ℓ? What is the complexity of $\ell = 0$ for $3 \leq c \leq 22$? Both are open in [14], whose authors say a W[1]-hardness result for $c = 2$ "would also not be surprising".

Open Problem 6 (Base of the exponent)**.** For $\ell = 0$, does a $(3-\varepsilon)^w n^{O(1)}$ algorithm exist, or is there a SETH-type lower bound? Theorem 4.1 gives only the exponent.

Open Problem 7 (The chain object)**.** Is there a family of tight instances with c and $\mathrm{tw}(I(H))$ bounded independently of the size and with a core of unbounded size, in the sense of Appendix F? This is the object that a W[1]-hardness proof through hard edges between far vertices (tunnels) would need.

Summary of results and their scope

The following table is the only status summary of the paper. "Input H" means that the statement is in terms of w, ℓ and n of the input packing; "pair-maximal H'" means that it is stated for a pair-maximal packing, and for an arbitrary input applies to a pair-maximal extension H' with the primed parameters of Section 2. "YES" and "optimal" mean that the statement needs a YES instance, respectively an optimal solution.

Result	Applies to	Needs	Status
Thm 3.4, 3.5 (CSP, 3^{w+1}, $7^{w+1}n^{2\ell}$)	input H	–	proved
Thm 4.1, Prop. 4.3 (ETH; w alone)	input H	–	proved from [14]
Thm 5.1 (components, kernel)	input H	–	proved; cites [15] Lemma 6.3, [2], [4]
Thm 5.10 (components of any size)	input H	–	proved; cites [15] Lemma 6.3
Thm 7.2 (n^ℓ algorithm)	input H (via H')	–	proved
Lemmas 6.2–6.6, Cor. 6.7	pair-maximal H'	optimal, YES	proved
Thm 8.2, Cor. 8.3, Prop. 8.5	pair-maximal H'	optimal, YES	proved
Prop. 8.7, 8.8	explicit instances	–	proved (cex: enumeration)
Thm 7.3, Prop. 7.4	pair-maximal H'	YES	proved; expander treewidth cited [9]
Appendices A–D	pair-maximal H' (App. B: any H)	mostly YES	proved
Appendix E, F	explicit families; tunnel model	–	proved
FPT in $\ell + w$ (Open Problem 1)			**open**
Reduction lemma (Open Problem 3)			**open**
Positions of T2 / deleted pointers (Open Problem 2)			**open**

A Twins, vertices outside the packing, and a shrink rule

By Lemma 2.2 the whole difficulty of $\ell > 0$ is the set U of at most ℓ hard edits. This appendix proves what can be said about the vertices outside $V(H)$ for pair-maximal packings H. The results are not used in the main text.

Conventions. Partitions, clusters, costs and optima are as in Section 2; $\Pi(v)$ is the cluster of v. Unless said otherwise, H is pair-maximal and $R = V \setminus V(H)$; every pair with an end in R is hard.

A.1 Twins

Lemma A.1 (Twins, strict form)**.** *Let $u \neq v$ be adjacent with $N[u] = N[v]$. For every partition Π with $\Pi(u) \neq \Pi(v)$, one of the two partitions obtained by moving u into $\Pi(v)$, respectively v into $\Pi(u)$, has cost smaller than Π by at least 1; more precisely the two cost changes sum to exactly -2. Hence u, v lie in one cluster in* every *optimum, with no hypothesis on H.*

Proof. For a vertex y and a set $Y \not\ni y$ let $f(y, Y) = |Y \setminus N(y)| - |Y \cap N(y)|$. Put $C_u = \Pi(u) \setminus \{u\}$ and $C_v = \Pi(v) \setminus \{v\}$ (possibly empty); $C_u \not\ni v$, $C_v \not\ni u$, and neither contains u or v. Moving u out of $\Pi(u)$ changes the status of the pairs uz, $z \in C_u$: before, uz is an edit iff it is a non-edge; after, iff it is an edge; the change is $-f(u, C_u)$. Moving u into $\Pi(v) = C_v \cup \{v\}$ changes, for $z \in C_v \cup \{v\}$, the status from "edit iff edge" to "edit iff non-edge"; the change is $f(u, C_v \cup \{v\}) = f(u, C_v) - 1$ because $uv \in E$. All other pairs keep their status. So

$$\delta_1 = -f(u, C_u) + f(u, C_v) - 1, \qquad \delta_2 = -f(v, C_v) + f(v, C_u) - 1.$$

Since $N(u) \setminus \{v\} = N(v) \setminus \{u\}$ we have $f(u, Y) = f(v, Y)$ for all $Y \subseteq V \setminus \{u, v\}$, hence $\delta_1 + \delta_2 = -2$. Thus $\min(\delta_1, \delta_2) \leq -1$. □

This is the critical-clique property used in kernelizations of Cluster Editing [7, 10] (it is also recalled in [15] when motivating Reduction Rule 6.2); the argument above is self-contained and shows that it is strict.

A.2 Viability of a hard edit

Lemma A.2 (Viability of a hard edit)**.** *Let S be any solution, U its hard part and $uv \in U$. Let $\mathrm{Bd}(uv)$ be the set of $z \notin Q(u) \cup Q(v) \cup \{u, v\}$ such that $\{u, v, z\}$ is an induced P_3 of $G \triangle \{uv\}$. Then for every $z \in \mathrm{Bd}(uv)$ one of uz, vz lies in U, and $|\mathrm{Bd}(uv)| \leq |U| - 1$.*

Proof. All three pairs of $\{u, v, z\}$ are hard ($z \notin Q(u) \cup Q(v)$). If neither uz nor vz is in S, the triple has in $G \triangle S$ the same pattern as in $G \triangle \{uv\}$, an induced P_3. Distinct z give distinct pairs, all different from uv. □

This gives a kernel-style condition (deleting uv needs fewer than ℓ common neighbours outside $Q(u) \cup Q(v)$) but no bound on the number of candidate pairs: all pendant edges are viable.

A.3 The vertices outside $V(H)$

Lemma A.3 (Structure of R)**.** *Let H be pair-maximal. For $r, s \in R$: if $rs \in E$ then $N[r] = N[s]$; if $rs \notin E$ then $N(r) \cap N(s) = \emptyset$. Consequently:*

1. *every component X of $G[R]$ (a* class*) is a clique all of whose vertices have the same closed neighbourhood $N[X]$ (they are true twins);*
2. *$A_X := N(X) \setminus X$ is contained in $V(H)$;*
3. *the sets A_X, over the classes X, are pairwise disjoint.*

Proof. Every pair with an end in R is hard. Let $rs \in E$ and $z \neq r, s$ be adjacent to exactly one of r, s. Then $\{r, s, z\}$ has exactly two edges, i.e. is an induced P_3 with all three pairs hard, contradicting pair-maximality; so $N[r] = N[s]$. If $rs \notin E$ and z is a common neighbour, then $r\,z\,s$ is an induced P_3 whose pairs rz, zs, rs all have an end in R, again a contradiction.

(1) Adjacent vertices of R have equal closed neighbourhoods; along a connected component all closed neighbourhoods coincide, and then any two vertices r, t of the component satisfy $r \in N[r] = N[t]$, so they are adjacent. (2) A neighbour of $x \in X$ lying in R is in the component of x, hence in X. (3) If $a \in A_X \cap A_Y$ for classes $X \neq Y$, then a is a common neighbour of some $x \in X$ and $y \in Y$, which are non-adjacent (different components of $G[R]$); this contradicts the second clause. □

For $x \in X$ and $a \in V(H)$ we thus have $xa \in E$ iff $a \in A_X$; in particular vertices of one class are twins towards $V(H)$ as well. (Lemma A.3 fails for vertex-maximal packings, see Proposition 4.5, whose instance also shows that Theorem A.5 fails without pair-maximality.)

Lemma A.4 (Exchange formula)**.** *Let Π be any partition, X a class with $X \subseteq C$ for a cluster C, $x = |X|$ and $a = |A_X \cap C|$. Let Π' be obtained by moving X out of C into a cluster of its own. Then* $\mathrm{cost}(\Pi') - \mathrm{cost}(\Pi) = x\,(x + 2a - |C|)$.

Proof. Only the pairs between X and $C \setminus X$ change status. By Lemma A.3, a vertex of X has exactly a neighbours in $C \setminus X$, namely $A_X \cap C$. Before the move such a pair is an edit iff it is a non-edge: $x(|C| - x - a)$ edits. After the move it is an edit iff it is an edge: xa edits. The difference is $xa - x(|C| - x - a) = x(x + 2a - |C|)$. □

Theorem A.5 (C1: no edit inside R)**.** *Let H be pair-maximal. In* every *optimum Π, every cluster meets R in the empty set or in exactly one whole class. Hence no edit of Π has both endpoints in R, and for every class X the cluster $\Pi(X)$ equals $X \cup T_X$ with $T_X := \Pi(X) \cap V(H)$.*

Proof. By Lemma A.3(1) and Lemma A.1, every class lies inside one cluster (a class of size 1 trivially). Suppose a cluster C contains two distinct classes X_1, X_2. Put $x_i = |X_i| \geq 1$ and $a_i = |A_{X_i} \cap C|$. The sets $X_1, X_2 \subseteq R$ and $A_{X_1} \cap C, A_{X_2} \cap C \subseteq V(H)$ are pairwise disjoint (Lemma A.3(2),(3)), so $|C| \geq x_1 + x_2 + a_1 + a_2$. Let Δ_i be the cost change of moving X_i out of C. Since Π is an optimum, $\Delta_1 \geq 0$ and $\Delta_2 \geq 0$, i.e. by Lemma A.4 $|C| \leq x_i + 2a_i$ for $i = 1, 2$. With the lower bound on $|C|$: $x_2 + a_2 \leq a_1$ and $x_1 + a_1 \leq a_2$. Adding gives $x_1 + x_2 \leq 0$, absurd. So no cluster contains two classes.

An edit with both ends in R joins two vertices r, s. If they lie in one class, they are adjacent and (by the above) in one cluster, so rs is no edit. If they lie in different classes they are non-adjacent, and rs is an edit iff they share a cluster, which was excluded. □

The strictness of the inequality is what makes the statement hold for every optimum (no ties, and no iteration or termination argument).

A.4 Consequences

Let (G, H, ℓ) be a *YES instance* with H pair-maximal, Π any optimum, S its edit set, so $|S| = \mathrm{OPT} \leq |H| + \ell$. Let U be the hard edits. By Lemma 2.2, $|S| - |H| = \sum_i (|x_i| - 1) + |U|$, so $|U| \leq \ell$. Write $q = \max_u |Q(u)| \leq 2c$.

Corollary A.6 (Normal form)**.** *For every class X let $T_X = \Pi(X) \cap V(H)$.*

(N1) Types of hard edits. *Every hard edit is either an* HH-edit *(both ends in $V(H)$) or an* RH-edit *$\{x, a\}$, $x \in X$ a class, $a \in V(H)$; the latter is a deletion if $a \in A_X \setminus T_X$ and an insertion if $a \in T_X \setminus A_X$, and $\{x, a\}$ is edited iff $\{x', a\}$ is, for all $x, x' \in X$. With h the number of HH-edits, $|U| = h + \sum_X |X| \cdot |A_X \triangle T_X| \leq \ell - \sum_i (|x_i| - 1) \leq \ell$.*

(N2) Frozen classes. *If $|X| > \ell$, then $\Pi(X) = X \cup A_X$.*

(N3) Attachment. *Every $a \in V(H)$ lies in A_X for at most one class X. If $z, z' \in A_X$ are non-adjacent then zz' is soft. If $a \in V(H) \setminus A_X$, $z \in A_X$ and $az \in E$ then az is soft; hence $|N(a) \cap A_X| \leq |Q(a)| \leq q$.*

(N4) Shrink rule. *If $|X| \geq \ell + 2$ and $x \in X$, then (G, H, ℓ) is a YES instance iff $(G - x, H, \ell)$ is. This holds for H pair-maximal without the YES hypothesis. If $A_X = \emptyset$ then X is a component of G and may be deleted without changing* OPT.

(N5) Size. *For a pair-maximal H and after exhausting the two rules in (N4), every class has at most $\ell + 1$ vertices and at most $|V(H)|$ classes have $A_X \neq \emptyset$, so $|R| \leq (\ell + 1)|V(H)|$ and $n \leq (\ell+2)|V(H)| \leq 3(\ell+2)|H|$. ($|V(H)|$ is not bounded in terms of ℓ, so this is not a kernel; see Remark A.8.)*

Proof. (N1) Pairs with an end in R are hard; by Theorem A.5 none has both ends in R. The vertices of X are twins towards $V(H)$ and lie in one cluster, which gives the all-or-none statement; xa is an edit iff exactly one of $a \in T_X$, $a \in A_X$ holds. Counting gives the identity; the bound is the first paragraph of this subsection.

(N2) If $\Pi(X) \neq X \cup A_X$ then, since $\Pi(X) = X \cup T_X$ (Theorem A.5), $A_X \triangle T_X \neq \emptyset$ and (N1) gives $|U| \geq |X| > \ell$, contradicting $|U| \leq \ell$.

(N3) The first claim is Lemma A.3(3). Let $x \in X$. If $z, z' \in A_X$ are non-adjacent, $z\,x\,z'$ is an induced P_3 with pairs zx, xz' hard, so zz' must be soft (pair-maximality). If $a \in V(H) \setminus A_X$ and $z \in A_X$ with $az \in E$, then $xa \notin E$ and $x\,z\,a$ is an induced P_3 with xz, xa hard, so az is soft, i.e. $z \in Q(a)$.

(N4) $x \in R$, so H is still a packing of $G - x$, still pair-maximal (an induced P_3 of $G - x$ is one of G), and $X \setminus \{x\}$ is a class of $G - x$ with the same A_X. ($\Rightarrow$) Restricting a partition of G to $V \setminus x$ does not increase its cost, so $\mathrm{OPT}(G-x) \leq \mathrm{OPT}(G) \leq |H|+\ell$. ($\Leftarrow$) Let Π be an optimum of $G-x$ with cost $\leq |H| + \ell$. Now $|X \setminus x| \geq \ell + 1 > \ell$, so (N2) in $G - x$ gives $\Pi(X \setminus x) = (X \setminus x) \cup A_X$. Add x to this cluster: $N[x] = X \cup A_X$, so every pair at x is satisfied, and the cost does not change. Hence $\mathrm{OPT}(G) \leq |H| + \ell$. If $A_X = \emptyset$, X is a clique component of G, which costs nothing, and H is disjoint from it.

(N5) After the rules each class has at most $\ell + 1$ vertices; the non-empty sets $A_X \subseteq V(H)$ are pairwise disjoint (Lemma A.3), and $|V(H)| \leq 3|H|$. Classes with $A_X = \emptyset$ were deleted. □

Remark A.7 (Tightness of the threshold in (N4)). For $\ell = 0$ the threshold $|X| \geq 2$ cannot be lowered to $|X| \geq 1$: take the 4-cycle of Proposition 4.6, $H = \{(0, 1, 3)\}$, $X = \{2\}$. Then $(G, H, 0)$ is a NO instance, but $G - 2$ is the induced P_3 0 1 3 itself, so $(G - 2, H, 0)$ is a YES instance. For $\ell = 1$ the threshold $|X| \geq \ell + 2 = 3$ cannot be lowered to $|X| \geq \ell + 1 = 2$: replace vertex 2 of the 4-cycle by two adjacent twins x, x', i.e. G has vertices $0, 1, 3, x, x'$ and edges $01, 13, 0x, 0x', 3x, 3x', xx'$, and $H = \{(0, 1, 3)\}$ (pair-maximal, $R = X = \{x, x'\}$). Then $\mathrm{OPT}(G) = 3$, so $(G, H, 1)$ is a NO instance, whereas $\mathrm{OPT}(G - x) = 2 = |H| + 1$, so $(G - x, H, 1)$ is a YES instance (both by enumerating all partitions). For $\ell = 2$ the same family with $|X| = 3$ does not separate ($\mathrm{OPT} = 3$ in both graphs). So the threshold is tight for $\ell \in \{0, 1\}$; tightness for $\ell \geq 2$ is open.

Remark A.8 (The rules and the input packing). The input packing H of CEaMP need not be pair-maximal. Extend it greedily (polynomial time) to a pair-maximal $H' \supseteq H$ with $t = |H'| - |H|$ new members, set $\ell' = \ell - t$ (answer NO if $\ell' < 0$, Lemma 2.4) and apply this section to (G, H', ℓ'). Then (N4), (N5) hold with ℓ' and H'; in particular $n \leq 3(\ell + 2)(|H| + \ell)$ after reduction, since $|H'| \leq |H| + \ell$. But $\mathrm{tw}(I(H'))$ may exceed $\mathrm{tw}(I(H))$, so these rules are a preprocessing for the parameters $(\ell, |H'|)$ or $(\ell, \mathrm{tw}(I(H')))$, not for $(\ell, \mathrm{tw}(I(H)))$ with the given H; and $|V(H')|$ is not bounded by ℓ, so (N5) is not a kernel.

Proposition A.9 (Partners of edits touching R)**.** *In the setting of Corollary A.6, let X be a class, $T = T_X$. Then:*

(P1) Deletion. *If $\{x, a\}$ is an RH-deletion, $a \in A_X$, then X is determined by a (the unique class with $a \in A_X$): it is a unary flag at a of weight $|X|$. Moreover a is then cut off from every $z \in A_X \cap T$ with $z \notin Q(a)$, each such edge az being a hard HH-deletion; so at least $|A_X \cap T| - q$ HH-edits accompany it.*

(P2) No detachment. *If $T \cap A_X = \emptyset$ then $T = \emptyset$.*

(P3) Insertion. *If $a \in T \setminus A_X$ then $T \cap A_X \neq \emptyset$, and for every $z \in T \cap A_X$ either az is soft ($a \in Q(z)$, at most q choices of a per z) or az is a hard non-edge, hence an inserted HH-edit.*

(P4) Wide classes. *If $|A_X| \geq \ell + q$ then $T \subseteq A_X$.*

Consequently, every candidate partner $a \in V(H)$ of a class in an RH-edit is unary (P1), local to some z (P3, soft case) or tied to an HH-edit (P3, second case; P1). This classifies the partners; it does not assert that the HH-edits are independent of the RH-edits.

Proof. (P1) Uniqueness is (N3). Let $a \in A_X \setminus T$ and $z \in A_X \cap T$, $z \notin Q(a)$. If $az \notin E$ then az would be soft by (N3), contradicting $z \notin Q(a)$; so $az \in E$, hard, and $a \notin \Pi(X) \ni z$, so az is deleted. At most q vertices z lie in $Q(a)$.

(P2) Suppose $T \neq \emptyset = T \cap A_X$. The cluster is $C = X \cup T$ (Theorem A.5), $a_X = |A_X \cap C| = 0$, and by Lemma A.4 moving X out changes the cost by $|X|(|X| - |X| - |T|) = -|X||T| < 0$, contradicting optimality.

(P3) $T \neq \emptyset$, so $T \cap A_X \neq \emptyset$ by (P2). Let $z \in T \cap A_X$. If $az \in E$ then az is soft by (N3) ($a \in V(H) \setminus A_X$), i.e. $a \in Q(z)$. Otherwise either az is soft (again $a \in Q(z)$) or it is a hard non-edge inside the cluster $\Pi(X)$, an insertion.

(P4) Suppose $T \not\subseteq A_X$. Put $\alpha = |A_X \cap T|$, $\delta = |A_X \setminus T|$, $\beta = |T \setminus A_X| \geq 1$. By (N1) there are $|X|\beta$ RH-insertions and $|X|\delta$ RH-deletions. For $b \in T \setminus A_X$ and $z \in A_X \cap T$ with $z \notin Q(b)$, the pair bz is hard, lies in a cluster, and is a non-edge by (N3); so it is a hard insertion, an HH-edit. There are at least $\beta \max(0, \alpha - q)$ of them. Hence $\ell \geq |U| \geq \beta(|X| + \max(0, \alpha - q)) + |X|\delta$. If $\alpha \geq q$ this gives $\ell \geq 1 + (\alpha - q) + \delta$, i.e. $|A_X| = \alpha + \delta \leq \ell + q - 1$. If $\alpha < q$ it gives $\delta \leq \ell - 1$ and $|A_X| \leq q - 1 + \ell - 1$. Either way $|A_X| < \ell + q$. □

B Counting lemmas for arbitrary packings

These lemmas hold for every packing H (no maximality unless stated) and describe what a solution of size at most $|H| + \ell$ can look like. All proofs use only pair-disjointness, a budget, and counting. They give neither a fixed-parameter algorithm in $\ell + w$ nor hardness; they are not used in the main text.

B.1 Hubs

In this subsection H is an arbitrary packing, $c = c(H)$, and (G, H, ℓ) is a YES instance where stated.

Lemma B.1 (HUB$^+$)**.** *Let H be any packing, (G, H, ℓ) a YES instance, u a vertex and $N_h(u) = \{x : ux \in E,\ ux \text{ not soft}\}$. Let Λ_u be the graph on $N_h(u)$ whose edges are the pairs $\{x, y\}$ with $xy \notin E$ and xy not soft (hard non-edges). Then:*

(a) $\nu(\Lambda_u) \leq \ell$ (matching number);

(b) Λ_u *has a vertex cover* T *with* $|T| \le 2\ell$*, and every pair of* $N_h(u) \setminus T$ *is an edge of* G *or is soft;*

(c) $\alpha(G[N_h(u)]) \le 2\ell + c + 1$ *where* $c = c(H)$*.*

Proof. (a) Let $x_1y_1, \ldots, x_ty_t$ be a matching of Λ_u and $P_i = (x_i, u, y_i)$. Since $ux_i, uy_i \in E$ and $x_iy_i \notin E$, P_i is an induced P_3. Its three pairs ux_i, uy_i, x_iy_i are hard. The $2t$ vertices x_i, y_i are distinct and different from u, so the pairs ux_i, uy_i ($i \le t$) are pairwise distinct, and the pairs x_iy_i are pairwise distinct and do not contain u. So $\{P_1, \ldots, P_t\}$ is hard over H and $t \le \ell$ by Lemma 2.3.

(b) The vertex set of a maximum matching of Λ_u is a vertex cover (by maximality) of size $2\nu \le 2\ell$. A pair of $N_h(u) \setminus T$ is not an edge of Λ_u, i.e. it is an edge of G or soft.

(c) Let J be independent in $G[N_h(u)]$. By (b), every pair of $J \setminus T$ is a non-edge of G and not an edge of Λ_u, hence soft. Put $J' = J \setminus T$. A soft pair of J' lies in a member P of H and is a non-edge, so it is the pair of the two ends of P (the other two pairs of P are edges); in particular $|V(P) \cap J'| \le 2$. If $J' \ne \emptyset$ fix $x \in J'$; every $y \in J' \setminus \{x\}$ gives a member $P_y \ni x, y$, and $P_y \ne P_{y'}$ for $y \ne y'$ because $|V(P_y) \cap J'| \le 2$. So $|J'| - 1 \le c$. Hence $|J| \le |T| + |J'| \le 2\ell + c + 1$. □

Corollary B.2 (HUB, hard-independent sets)**.** *In the setting of Lemma B.1, if* $Y \subseteq N_h(u)$ *and every pair of* Y *is a hard non-edge, then* $|Y| \le 2\ell + 1$*.*

Proof. Y is a clique of Λ_u, so it contains a matching of size $\lfloor |Y|/2 \rfloor$, and $\lfloor |Y|/2 \rfloor \le \ell$ by Lemma B.1(a). □

Remark B.3 (HUB$^+$ versus an independence-number bound). (1) Bounding $\alpha(G[N_h(u)])$ through the corollary (greedy choice of a subset of an independent set that is also free of soft pairs) gives only $(c+1)(2\ell+1) = (D/2+1)(2\ell+1)$, which is weaker than (c). (2) The matching form (a) is the relevant statement for Vertex Cover on the non-adjacency graph of $N(u)$: let $N_h(u)$ be two disjoint cliques of G of size m with no soft pair between them. Then $\alpha = 2$, but (a_i, u, b_i), a_i, b_i from different cliques, are m pair-disjoint induced P_3's, and $m \le \ell$ in a YES instance. Large independent sets are not the obstruction; large matchings of Λ_u are. (3) For a general H the lemma is applied to H' of Lemma 2.4 with c', ℓ'.

B.2 Pieces and small merges

Fix an arbitrary packing H and a solution S with $|S| \le |H| + \ell$ (neither optimality nor pair-maximality is used). Let $U = U^- \cup U^+$ be its hard part and $u = |U| \le |S| - |H| \le \ell$ (Lemma 2.2). The *pieces of* S are the connected components of the graph of kept hard edges, as in Section 8; every piece lies inside one cluster of $G \triangle S$.

Lemma B.4 (SM: small merge)**.** *Let* S*,* u *and the pieces be as above,* $D \ge \max_v |Q(v)|$ *(e.g.* $D = D(H)$*), and let* C *be a cluster of* $G \triangle S$*.*

(i) If $A \ne B$ *are pieces contained in* C*, then every pair* ab *(*$a \in A, b \in B$*) is soft or inserted.*

(ii) For such A, B *with* $|A| \ge |B|$*:* $|B|\,(|A| - D) \le |U^+ \cap (A \times B)| \le u$*, hence* $|A| \le D + u \le D + \ell$*.*

(iii) If C *is the union of at least two pieces, then* $|C| \le \max\{D + u + 1,\ 2D,\ 2u\}$*.*

Proof. (i) is Lemma 8.1.

(ii) Each vertex of B has at most D soft partners, so at most $D|B|$ pairs of $A \times B$ are soft. By (i) all other pairs of $A \times B$, at least $|A||B| - D|B|$, are inserted, and $|U^+| \le |U| = u$. If $|A| \ge D + u + 1$ then $|B|(|A| - D) \ge u + 1$ because $|B| \ge 1$, a contradiction; so $|A| \le D + u$.

(iii) Let A be a largest piece in C, $a = |A|$ and $r = |C| - a \geq 1$. By (i), every pair $\{x, y\}$ with $x, y \in C$ in different pieces is soft or inserted.

Case $a > D$. Each $x \in C \setminus A$ has at most D soft partners, so at least $a - D \geq 1$ vertices $y \in A$ give an inserted pair xy. These pairs are distinct for distinct (x, y), so $r(a - D) \leq |U^+| \leq u$. Then $r \leq u$ and $a \leq D + u/r$, so $|C| = a + r \leq D + u/r + r$. The function $r \mapsto r + u/r$ is convex on $[1, u]$ and takes the value $u + 1$ at both ends, so $|C| \leq D + u + 1$. (The case does not occur if $u = 0$.)

Case $a \leq D$. All pieces of C have at most D vertices. Each $x \in C$ has at least $|C| - D$ vertices of C outside its piece, at most D of which are soft partners; so x lies in at least $|C| - 2D$ inserted pairs $\{x, y\}$ with $y \in C$. Summing over x counts each pair twice: $2|U^+| \geq |C|(|C| - 2D)$. If $|C| > 2D$, then $|C| - 2D \geq 1$ and $|C| \leq 2|U^+| \leq 2u$. So $|C| \leq \max\{2D, 2u\}$. □

Remark B.5 (Qualifiers and relation to the original bound). (1) The qualifier "pieces sharing a cluster" is essential: a single piece may be an entire cluster of unbounded size, and a statement "pieces have size at most $D+\ell$" is false. (2) The bound (iii) implies the weaker bound $|C| \leq \max\{2D+\ell, 2\ell\}$: $2u \leq 2\ell$ and $2D \leq 2D + \ell$ are clear; $D + u + 1 \leq 2D + \ell$ if $D \geq 1$; if $D = 0$ and $u \geq 1$ then $u + 1 \leq 2u \leq 2\ell$; and if $D = u = 0$ no cluster with two pieces exists (the two cases of the proof are both impossible). (3) The bound (iii) is attained for instance by two singleton pieces and one inserted pair ($D = 0$, $u = 1$, $|C| = 2$); we do not claim it is tight in general. (4) Lemma B.4 says nothing about the *degrees* in a graph of pieces; it bounds the sizes of pieces that share a cluster and the number of cross pairs that are not soft.

B.3 The soft graph has bounded treewidth

Recall the soft graph G_{soft} (Section 2).

Proposition B.6 (Treewidth of the soft graph)**.** $\operatorname{tw}(W) \leq 3(w + 1) - 1 = 3w + 2$ *if $H \neq \emptyset$; G_{soft} is edgeless if $H = \emptyset$. The bound is attained for $w = 0$ (H a single P_3, G_{soft} a triangle).*

Proof. Let $(\mathcal{T}, \{\beta_t\})$ be a tree decomposition of $I(H)$ of width w and $\mathcal{T}_P = \{t : P \in \beta_t\}$, a subtree of $\mathcal{T}$ for each $P \in H$. Put $\beta'_t = \bigcup_{P \in \beta_t} V(P)$, so $|\beta'_t| \leq 3(w + 1)$. For every vertex $x \notin V(H)$ add a new leaf bag $\{x\}$ to an arbitrary node. *Edges:* a soft pair lies in $V(P)$ for some $P \in H$, and $P \in \beta_t$ for some t, so the pair lies in β'_t. *Connectivity:* for $x \in V(H)$ the nodes t with $x \in \beta'_t$ form $\bigcup_{P \ni x} \mathcal{T}_P$. The members through x pairwise share x, so they are pairwise adjacent in $I(H)$ and, the edge being covered by a bag, their subtrees $\mathcal{T}_P$ pairwise intersect. A union of connected subtrees that pairwise intersect is connected (two points of the union lie in subtrees $\mathcal{T}_P, \mathcal{T}_{P'}$ that meet; join them through a common node). The new leaf bags are single bags of size 1. So this is a tree decomposition of G_{soft} of width at most $3(w + 1) - 1$. □

B.4 Completing a fixed soft assignment

Proposition B.7. *Let x assign a subset $x_i \subseteq \operatorname{pairs}(P_i)$ to each P_i and let $b = |H| + \ell - \sum |x_i|$ be the excess budget left. Whether there is a set U of at most b hard pairs with $G \triangle (x \cup U)$ a cluster graph is decidable in $3^b n^{O(1)}$ time.*

Proof. Search tree on $G_x = G \triangle x$: if the current graph has an induced P_3, some pair of it must be flipped; soft pairs are frozen by x, so branch on its at most three hard pairs, reducing b by one (if all its pairs are soft, reject). Depth is at most b; flipping a pair twice is useless, so the search is exhaustive. □

The proposition is a reformulation, not progress towards fixed-parameter tractability in (ℓ, w): finding x is the whole problem, and the property "x is completable" is not local in x.

C The dynamic program for ω and the structure of Γ'

This appendix proves Theorem 7.3, using the notation $f(C) = \binom{|C|}{2} - 2e(C)$ of Lemma 6.1, and records two further statements on the graph Γ'. Throughout, H is pair-maximal.

C.1 Proof of Theorem 7.3

Theorem C.1 (Theorem 7.3 with bounds on m^*)**.** *Let H be pair-maximal and let ℓ, D, B, Z, L_B,* BIG*, V', Γ', ω be as in Section 7, and assume a tree decomposition of Γ' of width ω is given. Then one can compute in time $(Z(\omega+1))^{O(\omega)}n + n^{O(1)}$ a number m^* such that*

$$\mathrm{OPT}(G) \le |H| + \ell \iff |E| + \sum_{K} f(K) + m^* \le |H| + \ell,$$

where K ranges over the hard components of BIG *and f is as in Lemma 6.1. Moreover $m_{\mathrm{all}} \le m^* \le m_{\mathrm{conn}}$, where m_{all} (m_{conn}) is the minimum of $\sum_{C\in\pi'} f(C)$ over all partitions π' of V' with clusters of at most Z vertices (and, for m_{conn}, connected in Γ').*

Proof. *($\Leftarrow$)* Let π' be the partition of V' realising m^* (the dynamic program below produces a real partition) and $\pi = \pi' \cup \{K : K \text{ a component of BIG}\}$. By Lemma 6.1(a), $\mathrm{cost}(\pi) = |E| + \sum_K f(K) + m^* \ge \mathrm{OPT}$, so the right-hand side gives $\mathrm{OPT} \le |H| + \ell$. No structure of G is needed for this direction.

($\Rightarrow$) Let S be optimal with $|S| \le |H| + \ell$. By Lemma 6.6(γ) every component of BIG is a whole cluster; let π' be the restriction of S's partition to V'. Its clusters are Γ'-connected and have at most Z vertices:

- *Exceptional C.* $C \cap \mathrm{BIG} = \emptyset$ by (γ), so $C \subseteq V'$; $G[C]$ is connected by (α) and its edges are edges of Γ'. $|C| \le 2D + \ell + 1 \le Z$.
- *Non-exceptional C not in* BIG. By (β), C is a union of whole hard components, none of which is in BIG (a component of BIG is a cluster by itself), so $C \subseteq V'$. Each component is connected by hard edges, which are edges of Γ'. If there are at least two components, all pairs between them are soft, hence edges of Γ', so C is Γ'-connected, and $|C| \le 2D$. If there is exactly one component $K = C$, it is clean, so every vertex y of K has all other vertices of K as hard neighbours or soft partners, $|K| \le d_h(y) + D + 1 \le B + D + 1 = Z$ because $y \notin \mathrm{BIG}$ has $d_h(y) \le B$.

So π' is feasible for m_{conn} and $\mathrm{cost}(S) = |E| + \sum_K f(K) + \sum_{C\in\pi'} f(C) \ge |E| + \sum_K f(K) + m_{\mathrm{conn}} \ge |E| + \sum_K f(K) + m^*$.

The dynamic program. Take a nice tree decomposition of Γ' of width ω with empty root bag. A state at node t is a partition $\mathcal{P}$ of the bag X_t into blocks together with a size $s(\mathsf{b}) \in \{|\mathsf{b}|, \dots, Z\}$ for each block b (the number of vertices introduced so far, in the subtree of t, of the cluster containing b). Its value is the minimum, over partitions of the introduced vertices into clusters that realise $(\mathcal{P}, s)$ (every cluster either contains a block of $\mathcal{P}$ or has been *closed*, i.e. has no vertex left in the bag), of $\sum_C g(C)$, with $g(C) = \binom{|C|}{2} - 2e(C)$ evaluated on the introduced vertices of C. *Introduce* v into the block b of size s (or into a new block of size 1): the value increases by $s - 2k$, where k is the number of G-neighbours of v in b; here we use that an introduced vertex y that has been forgotten cannot be a G-neighbour of v, because yv would be an edge of Γ' and y, v would share a bag. *Forget* v: if its block has other bag vertices the block shrinks; otherwise the cluster is closed and is never extended. *Join* of two children with the same partition $\mathcal{P}$: for a block with bag part b

and sizes s_1, s_2 in the two children, put $x_1 = s_1 - |\mathsf{b}|$, $x_2 = s_2 - |\mathsf{b}|$; the new size is $s_1 + s_2 - |\mathsf{b}|$ and the value is $g_1 + g_2 - g(\mathsf{b}) + x_1 x_2$ summed over blocks, where $g(\mathsf{b}) = \binom{|\mathsf{b}|}{2} - 2e(\mathsf{b})$ is read off the bag. This is correct because pairs inside b and edges inside b are counted in both children, pairs between the two exclusive parts number $x_1 x_2$, and there is no G-edge between the exclusive parts (such an edge would be an edge of Γ' between vertices forgotten in different subtrees, which share no bag). *The join term $x_1 x_2$ and the subtraction of $g(\mathsf{b})$ are essential*: omitting them gives a different value on some inputs. At the root the value is $\sum_C f(C)$ over a partition of V' with clusters of size $\leq Z$. Call it m^* (the minimum over states; we treat the closed clusters as final).

Every state sequence yields a real partition of V' with cluster sizes $\leq Z$ and correct cost, so $m^* \geq m_{\text{all}}$. A Γ'-connected partition is represented: if the last bag vertex of a cluster C is forgotten at t, the vertex subtrees of C have a connected union (as C is Γ'-connected), which cannot contain the parent of t, so it lies below t and all of C has been introduced; hence no later vertex needs to join a closed cluster. So $m^* \leq m_{\text{conn}}$. There are at most $(\omega+1)^{\omega+1} Z^{\omega+1}$ states per node, a join combines at most $Z^{2(\omega+1)}$ pairs of states with equal partition, and the tree has $O(\omega n)$ nodes; the total time is $(Z(\omega+1))^{O(\omega)} n$. Computing d_h, BIG and Γ' takes $n^{O(1)}$.

Combining: OPT $\leq |H| + \ell$ implies $|E| + \sum_K f(K) + m^* \leq \text{cost}(S) \leq |H| + \ell$ by ($\Rightarrow$), and the converse is ($\Leftarrow$). □

Remark C.2. (1) Cleanliness of the components of BIG need not be tested: it is implied when the instance is a YES instance (Lemma 6.6(γ)), and the inequality itself accounts for the true cost $f(K)$; rejecting an unclean component is a harmless optimisation. (2) The proof uses that an exceptional cluster is disjoint from BIG and hence lies in V'. It is *not* true that $L_B \subseteq V'$: a component of BIG can contain vertices of small hard degree (pendant vertices of a hub component). (3) Here no pointer is guessed: far merges are carried by Γ'-connectivity. As for Theorem 7.2, D, B, Z, ℓ and ω refer to the pair-maximal instance (G, H', ℓ') of Lemma 2.4. Without a tree decomposition of Γ', one is computed in $2^{O(\omega)} n$ time with width $2\omega + 1$ [13].

C.2 Further statements

Lemma C.3 (Cost decomposition along a decomposition of the soft graph)**.** *Let π be a partition of V, let $(\mathcal{T}, \beta)$ be a tree decomposition of G_{soft} (Proposition B.6) and call a hard pair* local *if it lies in a common bag. Then* $\text{cost}(\pi) = \sum_t \text{loc}_t + \text{far}$*, where each edited soft pair and each edited local hard pair is charged to one bag containing it,* loc_t *depends only on $\pi|_{\beta(t)}$, and* far *is the number of edited non-local hard pairs, so* $\text{far} \leq |U|$.

Proof. Bookkeeping: all edited pairs are soft, or hard and local, or hard and non-local; soft pairs always lie in a bag. □

(The $\ell = 0$ CSP of Theorem 3.4 is the local part; all non-locality is in far.)

Proposition C.4 (Structure of Γ')**.** *Let (G, H, ℓ) be a YES instance with H pair-maximal.*

(a) $\Gamma' = G_{\text{soft}}[V'] \cup G_{\text{hard}}[V']$*, where G_{hard} is the graph of hard edges.*

(b) Put $Z^* = \max\{Z, (\ell+1)(2D+\ell+1)\}$. Every hard component K not in BIG *has $|K| \leq Z^*$. So "reject if a non-*BIG *hard component has more than Z^* vertices" is a safe polynomial rule.*

(c) Let $\Gamma'_{/K}$ be Γ' with every hard component contracted. Then $\text{tw}(\Gamma'_{/K}) \leq \text{tw}(\Gamma') \leq Z^*(\text{tw}(\Gamma'_{/K}) + 1) - 1$.

Proof. (a) Every edge of G is hard or soft, and soft pairs are edges of G_{soft}. (b) Take an optimal S with $|S| \le |H| + \ell$ and let U be its edited hard pairs. If no edge of K is deleted, K lies in one cluster C. If C is exceptional, $|C| \le 2D + \ell + 1 \le Z$ (Lemma 6.6(α)). If not, C is a union of whole hard components (Lemma 6.6(β)); with several components $|C| \le 2D$, and a single component $K = C$ is clean, so each $y \in K$ has all other vertices of K as hard neighbours or soft partners and $|K| \le d_h(y) + D + 1 \le B + D + 1 = Z$ ($K \not\subseteq$ BIG). If edges of K are deleted, there are at most $\ell + 1$ pieces of K (components of the kept hard edges); K is connected, so each piece contains an endpoint of a deleted edge and lies in an exceptional cluster of size $\le 2D + \ell + 1$; thus $|K| \le (\ell + 1)(2D + \ell + 1)$. (c) Hard components are connected in Γ' by hard edges, so $\Gamma'_{/K}$ is a minor of Γ'. Conversely replace every node of a tree decomposition of $\Gamma'_{/K}$ by the vertices of its component (of size $\le Z^*$ by (b)). □

By Proposition B.6 the soft graph G_{soft} has treewidth $\le 3w + 2$, so the unbounded treewidth of Γ' in Proposition 7.4 comes from hard edges between far vertices, which are free.

D Conflict weights, pruning and far pointers

This appendix concerns the two kinds of hard edit that the locality results of Section 8 leave open: inserted pairs of type T2 and deleted hard edges. It proves a *conflict lemma* that gives every soft pair a static weight, a defect bound and a polynomial NO-test that follow from it, bounded candidate lists, a pruned version of Theorem 7.3, and the statement that putting "locally bad" members into every bag does not repair the far pointers of Proposition 8.8 for distances 2 and 3. Nothing here proves or refutes fixed-parameter tractability in $\ell + w$.

Setting. H is pair-maximal (otherwise apply Lemma 2.4 and read the primed instance), $Q(v)$ is the set of soft partners, $N_h(v)$ the set of hard neighbours, $D \ge \max_v |Q(v)|$. A solution is identified with a partition π of V; write $x(p,q) = 1$ if p, q lie in one block and 0 otherwise; U is the set of edited hard pairs (so $|U| \le |S| - |H|$ by Lemma 2.2). For a soft pair $\sigma = (z, b)$ put

$$\Delta_{\neq}(z,b) = \{y \notin \{z,b\} : \ zy, by \text{ hard, exactly one an edge of } G\},$$

$$\Delta_{=}(z,b) = N_h(z) \cap N_h(b).$$

Both sets are computed from (G, H) alone. For a solution, the *weight* of the soft pair $\sigma = (z, b)$ is $\mathrm{wt}(\sigma) = |\Delta_{\neq}(z,b)|$ if z, b lie in one cluster (*merged*) and $\mathrm{wt}(\sigma) = |\Delta_{=}(z,b)|$ otherwise. A *conflict* is a pair (σ, y) with y in the relevant set.

D.1 The conflict lemma and its consequences

Lemma D.1 (W1: conflicts)**.** *Let π be* any *solution and $\sigma = (z, b)$ a soft pair.*

(a) If z, b are in one cluster, every $y \in \Delta_{\neq}(z,b)$ has $zy \in U$ or $by \in U$.

(b) If z, b are in different clusters, every $y \in \Delta_{=}(z,b)$ has $zy \in U$ or $by \in U$.

Distinct y give distinct edited pairs, hence $|U| \ge \mathrm{wt}(\sigma)$ for every soft pair σ.

Proof. (a) In a cluster graph, z, b in one cluster implies $x(z,y) = x(b,y)$ for all y. For $y \in \Delta_{\neq}(z,b)$ the G-values of the two hard pairs zy, by differ, so at least one of them is changed. (b) If neither zy nor by is edited, then $x(z,y) = x(b,y) = 1$ (both are edges of G), so z, b, y share a cluster, a contradiction. The edited pair chosen for y contains y and one of z, b, so different y give different pairs. □

Neither maximality nor optimality is used. Lemma 8.6 is the merged case of (a): for z, b in one cluster $\Theta(z,b) \subseteq \Delta_{\neq}(z,b)$, so $|\Theta(z,b)| \le |U|$. (The hypothesis that z, b lie in one cluster is essential: for unmerged z, b the set $\Theta(z,b)$ can be large.)

Lemma D.2 (W2: defect bound)**.** *For every solution,* $\sum_{\sigma\ soft} \mathrm{wt}(\sigma) \le 2D\,|U|$.

Proof. Map each conflict $((z,b),y)$ to an edited pair of $\{zy, by\}$ (Lemma D.1). An edited pair $e = \{p,y\}$ is chosen by conflicts (σ, y) with $\sigma \ni p$, at most $|Q(p)| \le D$ of them, and by conflicts (σ', p) with $\sigma' \ni y$, at most D of them. □

The constant $2D$ is attained for $D = 2$ ($c = 1$) by instances with $n = 6, 7$ found by computer search; for $D = 4$ and $D = 6$ the same search reached only $1/2$ and $1/3$ of the bound, so attainment for $D \ge 4$ is open.

Corollary D.3 (LB: a lower bound from the weights)**.** *Let* $\kappa_i \ge 1$ *be the number of edited pairs among* $\mathrm{pairs}(P_i)$ *and let* $\mathrm{Wt} = \sum_\sigma \mathrm{wt}(\sigma)$ *be the total weight of a solution* π*. Then* $\mathrm{cost}(\pi) \ge \sum_i \kappa_i + \lceil \mathrm{Wt}/2D \rceil$.

Proof. $\mathrm{cost}(\pi) = \sum_i \kappa_i + |U|$ (Lemma 2.2) and $|U| \ge \mathrm{Wt}/2D$ by Lemma D.2. □

The right-hand side is not exact: minimised over assignments of the soft pairs that are transitive on all-soft triples, it can equal $|H|$ although $\mathrm{OPT} > |H|$ (found by computer search on small random instances; the triples with two soft pairs and one hard pair are not checked). We only use it as a lower bound.

Lemma D.4 (Static pruning)**.** *Let* $|S| \le |H| + \ell$*. If* $|\Delta_{\neq}(u,v)| > \ell$*, the soft pair* (u,v) *is split in* S*; if* $|\Delta_{=}(u,v)| > \ell$ *it is merged in* S*.*

Proof. Lemma D.1 and $|U| \le |S| - |H| \le \ell$. □

Lemma D.5 (DEL: rigid hard edges)**.** *If* zy *is a deleted hard edge of a solution, then* $|U| \ge 1 + |N_h(z) \cap N_h(y)|$*. Hence, if* $|S| \le |H| + \ell$*, a hard edge with at least* ℓ *common hard neighbours is never deleted (it is* rigid*), and the hard components of rigid edges lie inside single clusters.*

Proof. z and y are in different clusters. A common hard neighbour w cannot be kept with both, so wz or wy is a deleted hard edge; distinct w give distinct pairs (each contains w), all different from zy. □

A member $P = (x, y, z)$ with centre y has three states of member-cost one: delete xy, delete yz, insert xz. A state fixes whether each of the three pairs of P is merged, hence a weight for each. Call P *locally bad* if in each of its three states some pair of P has positive weight. This is a static property.

Lemma D.6 (BAD: few locally bad members)**.** *For every solution of size* $|H| + \varepsilon$ *the number of locally bad members is at most* $2c\varepsilon$*. Hence more than* $2c\ell$ *locally bad members is a NO-instance of CEaMP with budget* ℓ *(a polynomial test).*

Proof. By Lemma 2.2, $\varepsilon = \sum_i (\kappa_i - 1) + |U|$. A bad member with $\kappa_i \ge 2$ contributes at least 1 to the first sum. A bad member with $\kappa_i = 1$ is in one of its three cost-one states, which has a conflict (σ, y) with σ a pair of P_i; by Lemma D.1 an edit of U is incident to $V(P_i)$. An edit has two ends and each vertex lies in at most c members, so there are at most $2c|U|$ such members. In total at most $\sum_i (\kappa_i - 1) + 2c|U| \le 2c\varepsilon$. □

Proposition D.7 (Candidate lists for (T2) pairs and deleted edges)**.** *Let H be pair-maximal, S an* optimal *solution with $|S| \le |H| + \ell$, and $ab \in U$. Define, from (G, H) alone,*

$$L_1(a) = \begin{cases} N_h(a) \cup \bigcup_{z \in N_h(a)} Q(z) & \text{if } d_h(a) \le D + \ell, \\ \emptyset & \text{otherwise,} \end{cases} \qquad L_2(b) = \bigcup_{z \in Q(b),\ |\Delta_{\neq}(z,b)| \le \ell} \Delta_{\neq}(z, b).$$

Then $|L_1(a)| \le (D + \ell)(D + 1)$ and $|L_2(b)| \le D\ell$, and:

(i) if ab is a deleted hard edge, then $b \in L_1(a)$ and $a \in L_1(b)$;

(ii) if ab is an inserted pair of type (T2) with middle vertex z, az hard (kept) and zb soft (Theorem 8.2), then $b \in L_1(a)$ and $a \in L_2(b)$.

Pairs of type (T1) are not covered; they are bag-local by Proposition 8.5.

Proof. Every endpoint x of a pair of U has $d_h(x) \le D + |U| \le D + \ell$ (Lemma 6.4(a)), so the guard in L_1 holds for the ends. (i) The deleted pair is a hard edge, so $b \in N_h(a)$, and symmetrically. (ii) $z \in N_h(a)$ and $b \in Q(z)$, so $b \in L_1(a)$, and $|L_1(a)| \le (D + \ell) + (D + \ell)D$. For the soft end: $z \in Q(b)$ ($\le D$ choices); since az is kept and ab is inserted, a, z, b lie in one cluster, so zb is merged, and az is a hard edge while ab is a hard non-edge, so $a \in \Delta_{\neq}(z, b)$, and $|\Delta_{\neq}(z, b)| \le |U| \le \ell$ by Lemma D.1(a). Hence $a \in L_2(b)$ and $|L_2(b)| \le D\ell$. □

Optimality is used only through Theorem 8.2. The lists say where the partner of a given end can be; they do not say where the first end is (Corollary 6.7).

D.2 A pruned form of Theorem C.1

Let BIG, V', Z be as in Theorem C.1 and let Γ^ℓ be the graph on V' whose edges are all hard edges inside V' and all soft pairs (u, v) inside V' with $|\Delta_{\neq}(u, v)| \le \ell$. It is a spanning subgraph of Γ', so $\mathrm{tw}(\Gamma^\ell) \le \mathrm{tw}(\Gamma')$, and it is computable from (G, H) alone.

Theorem D.8 (P: pruned ω-theorem)**.** *Let H be pair-maximal and let $\omega_\ell = \mathrm{tw}(\Gamma^\ell)$, with a tree decomposition of that width given. Then one can compute in time $(Z(\omega_\ell+1))^{O(\omega_\ell)}n+n^{O(1)}$ a number m^* with*

$$\mathrm{OPT}(G) \le |H| + \ell \iff |E| + \sum_K f(K) + m^* \le |H| + \ell,$$

K ranging over the hard components of BIG*. So CEaMP is FPT in $(\ell, D, \mathrm{tw}(\Gamma^\ell))$.*

Proof. Let $e^+(C)$ be the number of edges of G inside C that are edges of Γ^ℓ, and $\tilde{f}(C) = \binom{|C|}{2} - 2e^+(C)$. Run the dynamic program of the proof of Theorem C.1 on Γ^ℓ, with e^+ in place of e everywhere (in the introduce step k counts the Γ^ℓ-neighbours of v in the block, and in the join step there is no Γ^ℓ-edge between the exclusive parts). Exactly as there, it computes $m^* =$ the minimum of $\sum_C \tilde{f}(C)$ over Γ^ℓ-connected partitions of V' with clusters of at most Z vertices, in the stated time.

($\Leftarrow$) $e^+ \le e$, so $f(C) \le \tilde{f}(C)$ and the partition realising m^*, completed by the components of BIG, has true cost at most $|E| + \sum_K f(K) + m^*$ (Lemma 6.1(a)); no structure is needed.

($\Rightarrow$) Let S be optimal, $|S| \le |H| + \ell$. By the proof of Theorem C.1, the clusters of S inside V' have at most Z vertices and are connected through edges of G inside the cluster and (for a cluster with several hard components) soft pairs inside the cluster. A soft pair inside a cluster is merged, so $|\Delta_{\neq}| \le \ell$ by Lemma D.4 and it is an edge of Γ^ℓ; every hard edge is an edge of Γ^ℓ. Hence these clusters are Γ^ℓ-connected and $e^+(C) = e(C)$, $\tilde{f}(C) = f(C)$. So $\mathrm{cost}(S) = |E| + \sum_K f(K) + \sum_C f(C) \ge |E| + \sum_K f(K) + m^*$. □

Theorem D.8 does not use optimality of the merges, only their cost, and it therefore cannot decide the open question.

Proposition D.9 (Theorem D.8 is strictly stronger than Theorem 7.3)**.** *Let F be a cubic graph with an orientation in which every in-degree is* 1 *or* 2 *(so every vertex has in-degree ≥ 1; for a* 2*-edge-connected F such an orientation exists by Robbins' theorem), and let (G_F, H_F) be the instance of Proposition 7.4. For every budget $\ell \leq 4$, every component of Γ^ℓ has at most* 5 *vertices (so $\mathrm{tw}(\Gamma^\ell) \leq 4$), whereas $\mathrm{tw}(\Gamma') \geq \mathrm{tw}(F)$.*

Proof. For the oriented edge $e = (p \to q)$ the soft pairs $a_e b_e$ and $a_e c_e$ join $\mathcal{K}_p$ and $\mathcal{K}_q$. The soft partners of a_e are b_e, c_e and those of b_e are a_e, c_e (the three pairs of P_e). Hence $\Delta_{\neq}(a_e, b_e)$ consists of the vertices $y \in \mathcal{K}_p \setminus \{a_e\}$ (the pair $a_e y$ is a hard edge and $b_e y$ a hard non-edge), of which there are $2 + \mathrm{in}(p)$, and of the vertices $y \in \mathcal{K}_q \setminus \{b_e, c_e\}$ (here $b_e y$ is a hard edge and $a_e y$ a hard non-edge), of which there are $1 + \mathrm{in}(q)$. So $|\Delta_{\neq}(a_e, b_e)| = 3 + \mathrm{in}(p) + \mathrm{in}(q) \geq 5$, and the same count holds for $a_e c_e$. For $\ell \leq 4$ both cross pairs are pruned, the only edges of Γ^ℓ are those inside the cliques $\mathcal{K}_v$, and $|\mathcal{K}_v| = 3 + \mathrm{in}(v) \leq 5$. The bound on $\mathrm{tw}(\Gamma')$ is Proposition 7.4. □

The in-degree condition matters: with a source p (in-degree 0) the weight can be 4.

D.3 Far pointers and locally bad members

An edited pair $e = (a, b)$ of a solution S is *two-sided* if there are vertices t, t' with (t, a) soft, (t', b) soft, and $\{a, b, t\}$, $\{a, b, t'\}$ induced P_3's of $G \triangle (S \setminus \{e\})$. It is *far* (at distance d) if the members containing a and b are at distance $\geq d$ in $I(H)$.

Proposition D.10 (NF-bad fails for distance 2 and 3)**.** *The following statement is false for "far" meaning distance ≥ 2 and for "far" meaning distance ≥ 3 in $I(H)$:* there is an optimum in which every far two-sided edit touches a locally bad member.

Witnesses. Both instances have $n = 10$, a pair-maximal modification-disjoint packing, a unique optimum, $\mathrm{OPT} = |H| + 1 = 6$, no locally bad member, and exactly one edit, a hard deleted edge, which is two-sided (found by enumerating all set partitions). *Distance* 2*:* $E = \{01, 06, 12, 23, 29, 34, 45, 56, 67, 78, 79\}$, $H = \{(0,1,2), (2,3,4), (4,5,6), (6,7,8), (2,9,7)\}$; the optimum $\{0,1\}, \{2,9\}, \{3,4\}, \{5,6\}, \{7,8\}$ deletes $\{0,6\}$ (witnesses 1 and 5). *Distance* 3*:* $E = \{01, 08, 12, 14, 23, 34, 45, 49, 56, 67, 78\}$, $H = \{(0,1,2), (2,3,4), (4,5,6), (6,7,8), (1,4,9)\}$; the optimum $\{0,1\}, \{2,3\}, \{4,9\}, \{5,6\}, \{7,8\}$ deletes $\{0,8\}$ (witnesses 1 and 7). In both, $I(H)$ has $c = 3$. □

Remark D.11 (Scope). Pointers at distance $\leq K$ are bag-local in a decomposition of $I(H)^K$, whose width is a function of (w, c, K), so the version of the statement that matters is the one with "far" larger than any function of (ℓ, w). Proposition D.10 does not refute it, and Proposition 8.8 does not either: the far ends of its edit are exactly the two locally bad members (for $m = 3, 4, 5$ these are the only locally bad members, by enumeration). So the idea "put the locally bad members into every bag" is excluded for distances 2 and 3 only; for unbounded distance it is open.

E Reduction on unions of members fails: the spine family

Theorem 5.10 applies Rule 6.2 of [15] to components of $I(H)$. This appendix shows that the natural generalisation to pair-closed unions of *members* (Section 5) cannot prove the reduction lemma (Open Problem 3), already for $w = 1$ and $\ell = 1$. Soundness of the rule is cited from [15]; the proposition itself is proved here, and does not use the rule. Throughout, Γ', BIG, B, Z are as in Section 7; a set $\mathcal{F}$ of members is pair-closed, reducible and irreducible as in Section 5.

Proposition E.1 (The spine family $\mathrm{Sp}(F)$)**.** *Let F be a cubic graph with an orientation of in-degree at most 2 at every vertex. Let F' be F plus a new vertex $*$ and an edge $e_0 = (t \to *)$, $t \in V(F)$. Order the edges of F' as $e_0, e_1, \ldots, e_N$ so that e_0 comes first,* consecutive edges have different heads and $\mathrm{head}(e_1) \neq \mathrm{tail}(e_0) = t$ *(possible, since every head occurs at most twice and t has in-degree at most 2). Write $h(i)$ and $\mathrm{t}(i)$ for the head and tail of e_i. The instance $\mathrm{Sp}(F)$ has vertices a_i, b_i, c_i, u_i $(0 \le i \le N)$, z and y; for $v \in V(F')$ the sets $\mathcal{K}_v^+ = \{a_i : \mathrm{t}(i) = v\} \cup \{b_i, c_i, u_i : h(i) = v\}$ (so $\mathcal{K}_*^+ = \{b_0, c_0, u_0\}$). Edges: all pairs inside each $\mathcal{K}_v^+$; the cross edges $a_i b_i$; the spine edges $b_i u_{i+1}$ $(i < N)$ and $b_N z$; and $y u_0$, $y c_0$ (and no other edge at y). Packing: $P_i = (a_i, b_i, c_i)$ and $X_i = (u_i, b_i, u_{i+1})$, both with centre b_i ($u_{N+1} := z$); $\ell = 1$. Then:*

(S1) the members are induced P_3's with pairwise disjoint pair sets, H is pair-maximal, $I(H)$ is a caterpillar (spine $X_0, \ldots, X_N$, leaf P_i at X_i), so $w = 1$, $c = 2$, $D = 4$;

(S2) $(G, H, 1)$ is a YES instance and $\mathrm{OPT} = |H| + 1$;

(S3) all hard degrees are at most $7 \le B = 11$, so $\mathrm{BIG} = \emptyset$;

(S4) F is a minor of Γ', so $\mathrm{tw}(\Gamma') \ge \mathrm{tw}(F)$, which is $\Omega(|V(F)|)$ for cubic expanders [9] (cited);

(S5) $I(H)$ is connected and tight: $\tau = |H|$;

(S6) every nonempty pair-closed set $\mathcal{F}$ of members (connected or not) is irreducible.

Proof. (S1) Every induced P_3 of G uses a cross edge or a spine edge or a y-edge, because two edges inside cliques at a common vertex lie in one clique. P_i is induced: $a_i b_i$ is a cross edge, $b_i c_i$ lies in $\mathcal{K}_{h(i)}^+$, $a_i c_i \notin E$ because $\mathrm{t}(i) \neq h(i)$. X_i is induced: $u_i b_i$ lies in $\mathcal{K}_{h(i)}^+$, $b_i u_{i+1}$ is a spine edge, and $u_i u_{i+1} \notin E$ since consecutive heads differ (z is adjacent to b_N only). The pair sets $\{a_i b_i, b_i c_i, a_i c_i\}$, $\{u_i b_i, b_i u_{i+1}, u_i u_{i+1}\}$ are pairwise disjoint. Pair-maximality: no two hard edges form an induced P_3. The soft edges are $a_i b_i$, $b_i c_i$, $u_i b_i$, $b_i u_{i+1}$, so the hard edges are the other clique edges and $y u_0$, $y c_0$. Two hard clique edges at a vertex lie in one clique, so their other ends are adjacent; the hard edges at y go to u_0, c_0, which are adjacent; and a hard edge at u_0 or c_0 inside $\mathcal{K}_*^+ = \{b_0, c_0, u_0\}$ can only be $u_0 c_0$, whose ends are both adjacent to y.

(S2) The partition $\{\mathcal{K}_v^+\}$ together with $\{z\}$, with y put into $\mathcal{K}_*^+$, deletes exactly the $2(N + 1) = |H|$ cross and spine edges (all soft) and inserts the single pair $y b_0$; its cost is $|H| + 1$. For $\mathrm{OPT} \ge |H| + 1$ see (S6) below.

(S3) A vertex of $\mathcal{K}_v^+$ has at most $|\mathcal{K}_v^+| - 1 = 2 + 2\,\mathrm{in}(v) \le 6$ hard neighbours inside its clique, and u_0, c_0 have one more hard neighbour, y (which has hard degree 2).

(S4) Contract every $\mathcal{K}_v^+$ (a clique). The cross edges $a_i b_i$ join $\mathcal{K}_{\mathrm{t}(i)}^+$ and $\mathcal{K}_{h(i)}^+$, so the quotient contains F.

(S5) In the partition of (S2), the pair $a_i b_i$ is deleted, $b_i c_i$ kept, $a_i c_i$ a non-edge in different clusters; $u_i b_i$ kept, $b_i u_{i+1}$ deleted, $u_i u_{i+1}$ a non-edge in different clusters. So its restriction to $V^- = V \setminus \{y\}$ edits exactly one pair per member: $\tau \le |H|$, hence $\tau = |H|$.

(S6) For pair-closed $\mathcal{F}$ the restriction of the solution of (S2) to $W_{\mathcal{F}}$ has at most $|\mathcal{F}|$ edits (no member outside $\mathcal{F}$ has a pair inside $W_{\mathcal{F}}$, and $y b_0$ leaves $W_{\mathcal{F}}$), so $\tau(G[W_{\mathcal{F}}]) = |\mathcal{F}|$ and the optimal $S_{\mathcal{F}}$ are the *tight* ones: exactly one edited pair per member of $\mathcal{F}$ and no hard pair is edited. Suppose such an $S_{\mathcal{F}}$ satisfied the boundary condition. We use repeatedly: two vertices $p, q \in W_{\mathcal{F}}$ joined by a hard non-edge are non-adjacent in $G[W_{\mathcal{F}}] \triangle S_{\mathcal{F}}$, and a hard edge is kept.

Case 1: $\mathcal{F}$ contains a spine member; let X_i be the one of smallest index.

$i \ge 1$. Then $X_{i-1} \notin \mathcal{F}$, and $P_{i-1} \notin \mathcal{F}$: otherwise $b_{i-1}, u_i \in W_{\mathcal{F}}$ and the pair $b_{i-1} u_i$ of the outside member X_{i-1} would lie inside $W_{\mathcal{F}}$. So $a_{i-1}, b_{i-1}, u_{i-1} \notin W_{\mathcal{F}}$ and $N(b_{i-1}) \cap W_{\mathcal{F}} = \{u_i\} \cup Y$

with $Y = \mathcal{K}^+_{h(i-1)} \cap W_{\mathcal{F}}$. For $m \in Y$ the pair $u_i m$ is a non-edge ($h(i) \neq h(i-1)$) and is soft only if $m = u_{i+1}$. If Y contains some $m \neq u_{i+1}$, then u_i, m are two non-adjacent vertices of $N(b_{i-1}) \cap W_{\mathcal{F}}$: the boundary condition fails. If $Y = \{u_{i+1}\}$, the set $\{u_i, u_{i+1}\}$ must be a cluster, so X_i inserts $u_i u_{i+1}$ and keeps $u_i b_i$, $b_i u_{i+1}$; then b_i lies in this cluster, but $b_i \notin N(b_{i-1})$: contradiction. So $Y = \emptyset$ and $\{u_i\}$ is a cluster, i.e. X_i deletes $u_i b_i$. If $P_i \notin \mathcal{F}$, the vertex $c_i \notin W_{\mathcal{F}}$ is adjacent to u_i and b_i, which lie in different clusters: the boundary condition fails at c_i. If $P_i \in \mathcal{F}$, the hard edge $u_i c_i$ is kept, contradicting that u_i is a singleton.

$i = 0$. The vertex $y \notin W_{\mathcal{F}}$ has $N(y) \cap W_{\mathcal{F}} = \{u_0\}$ if $P_0 \notin \mathcal{F}$ and $\{u_0, c_0\}$ if $P_0 \in \mathcal{F}$. First case: $\{u_0\}$ is a cluster, so X_0 deletes $u_0 b_0$, and $c_0 \notin W_{\mathcal{F}}$, adjacent to $u_0, b_0 \in W_{\mathcal{F}}$ in different clusters, violates the condition. Second case: $\{u_0, c_0\}$ must be a cluster, so b_0 is separated from both: X_0 deletes $u_0 b_0$ and P_0 deletes $b_0 c_0$. Then $a_0 b_0$ and $b_0 u_1$ are kept, so a_0, u_1 are adjacent to b_0. Now $a_0 \in \mathcal{K}^+_t$ and $u_1 \in \mathcal{K}^+_{h(1)}$ with $h(1) \neq t$ *by the ordering condition*, so $a_0 u_1$ is a hard non-edge (the soft partners of a_0 are b_0, c_0), and $a_0 b_0 u_1$ is an induced P_3 of $G[W_{\mathcal{F}}] \triangle S_{\mathcal{F}}$: contradiction.

Case 2: $\mathcal{F}$ contains only leaves. Take $P_i \in \mathcal{F}$; $u_i, u_{i+1} \notin W_{\mathcal{F}}$ (for $i = N$ read z for u_{i+1}). If $i = N$ then $N(z) \cap W_{\mathcal{F}} = \{b_N\}$ must be a cluster, i.e. b_N is a singleton; but each of the three single edits of P_N (delete $a_N b_N$, delete $b_N c_N$, insert $a_N c_N$) keeps b_N with a_N or c_N. So let $i < N$. Then $N(u_{i+1}) \cap W_{\mathcal{F}} = \{b_i\} \cup Y$ with $Y = \mathcal{K}^+_{h(i+1)} \cap W_{\mathcal{F}}$. For $m \in Y$ the pair $b_i m$ is a non-edge ($h(i+1) \neq h(i)$) unless $m = a_i$ (possible when $\mathrm{t}(i) = h(i+1)$), and it is soft only if $m = a_i$ (the soft partners of b_i are a_i, c_i, u_i, u_{i+1}). *(2a)* If Y contains some $m \neq a_i$, then b_i, m are non-adjacent in $N(u_{i+1}) \cap W_{\mathcal{F}}$: failure. *(2b)* If $Y = \emptyset$, then $\{b_i\}$ is a cluster, impossible as above. *(2c)* If $Y = \{a_i\}$, then $P_{i+1} \notin \mathcal{F}$ and $b_{i+1} \in \mathcal{K}^+_{h(i+1)}$ is outside $W_{\mathcal{F}}$ with $N(b_{i+1}) \cap W_{\mathcal{F}} = \{a_i\}$; this must be a cluster, so a_i is a singleton and $a_i b_i$ is deleted. But then $\{a_i, b_i\} = N(u_{i+1}) \cap W_{\mathcal{F}}$ is not a cluster ($a_i b_i$ is not an edge of $G \triangle S_{\mathcal{F}}$): contradiction.

So $\mathcal{F}$ is irreducible. For the lower bound of (S2): if $\mathrm{OPT} = |H|$, an optimum S has $|S| = \tau$ for the whole component $\mathcal{C} = I(H)$, so $S = S_{\mathcal{C}}$ is tight and optimal on $V^- = V \setminus \{y\}$; by (S6) it violates the boundary condition at y, and Lemma 5.5 gives an edited pair $\{y, u\}$, which contradicts $S = S_{\mathcal{C}}$. So $\mathrm{OPT} \geq |H| + 1$. □

Remark E.2 (Finite checks). Claims (S1)–(S5) and $\mathrm{OPT} = |H| + 1$ were verified by computer for F the triangle ($n = 18$, OPT by an exact subset dynamic program) and for K_4 ($n = 30$). Claim (S6) was checked, in agreement with the proof, for the triangle on all 90 pair-closed subfamilies with $|\mathcal{F}| \leq 6$ and on the whole component, and for K_4 on all 632 pair-closed subfamilies with $|\mathcal{F}| \leq 4$ for 8 random admissible orderings. The proof above is the justification.

Consequence: an obstruction, not a refutation of the reduction lemma. No reduction obtained by iterating Rule 6.2 on pair-closed unions of members (the form used for components in Section 5) can prove Open Problem 3 by itself: on $\mathrm{Sp}(F)$ it removes nothing, although $\ell = 1$, $w = 1$, $\mathrm{BIG} = \emptyset$ and $\mathrm{tw}(\Gamma') = \Omega(|V(F)|)$. In particular the counting argument of Lemma 5.6 does not extend from components to members: all $2(N+1)$ members are irreducible although $\ell = 1$. The mechanism is Lemma 5.5: along the spine each member is made irreducible by an edit at a shared vertex of its neighbour, down to the trigger y. Other generalisations of the rule (to vertex sets that are not unions of members) are not excluded.

F Tunnels: locks, memory and a restricted model

The most natural attempt at W[1]-hardness in $\ell + w$ is a reduction from Grid Tiling in which hard edges between far-apart vertices (*tunnels*) carry equalities between distant parts of $I(H)$ of

bounded width. This appendix proves what a tunnel can and cannot do and isolates one missing gadget (Open Problem 7). Neither W[1]-hardness nor fixed-parameter tractability of CEaMP in $\ell + w$ follows from anything below.

F.1 Zero-excess solutions and tunnels

Soft and hard pairs and $Q(v)$ are as in Section 2. A solution S is *zero-excess* (ZE) if $|S| = |H|$. A member (a, b, c) (centre b) is in state 1, 2 or 3 if its edited pair is ab, bc or ac. By Lemma 2.2 a ZE solution has exactly one edit in each member and no other edit, so it is determined by its vector of states; let $\Sigma = \Sigma(G, H) \subseteq \{1, 2, 3\}^H$ be the set of these vectors. None of the lemmas of this section except Lemma F.7 needs pair-maximality.

Lemma F.1 (Zero-excess structure)**.** *Let S be a ZE solution. (a) Every hard edge lies inside a cluster and every hard non-edge joins two clusters. (b) Every member is in one of the states $1, 2, 3$. (c) Every hard component lies in one cluster, and two vertices of one cluster in different hard components form a soft pair.*

Proof. (a),(b) are Lemma 2.2 with $|S| = |H|$: no hard pair is edited and each member has exactly one edit. A member (a, b, c) with togetherness pattern (t_{ab}, t_{bc}, t_{ac}) has the cluster patterns $000, 100, 010, 111$ of costs $2, 1, 1, 1$, so the one-edit patterns are the states $2, 1, 3$. (c) A hard edge is unedited, so its ends are together; two vertices of one cluster in different hard components are not joined by a hard edge, and a hard non-edge inside a cluster would be edited, so the pair is soft. □

A *tunnel* is a hard edge xx' with $N_h(x) = \{x'\}$, $N_h(x') = \{x\}$ and $Q(x) \cap Q(x') = \emptyset$; tunnels are always taken pairwise vertex-disjoint. A *stub* at x is a new vertex x' adjacent to x only (then $Q(x') = \emptyset$). A vertex is an *anchor* if it lies in a member and has no hard edge. *Cutting* a tunnel means editing xx' (one hard edit).

Lemma F.2 (A tunnel is a lock)**.** *Let xx' be a tunnel and S a solution in which all hard pairs at x and at x' are unedited. Then the cluster of x is $\{x, x'\}$. If S is ZE, then no member has x as centre, every member through x has x as an end with the pair at x that is an edge deleted, and the other two vertices of that member are together.*

Proof. Let $t \neq x, x'$ be in the cluster of x. If $t \notin Q(x)$ then xt is hard, hence a non-edge as $N_h(x) = \{x'\}$, hence edited, a contradiction; so $t \in Q(x)$, and likewise $t \in Q(x')$, contradicting $Q(x) \cap Q(x') = \emptyset$. In a ZE solution a member (y, x, z) with centre x would have both pairs at x apart, hence two edits. If x is an end, the pair x–centre is apart (an edit), the non-edge at x is apart (no edit), so the remaining pair is together. □

Lemma F.3 (Anchors: singleton = forced states; inactive anchors cost nothing)**.** *Let x be an anchor and σ a ZE solution of (G, H). Then x is a singleton of σ iff every member through x has x as an end and is in the state that deletes the edge at x. Moreover: (a) if G' is G with a stub at x, then G' has a ZE solution iff G has a ZE solution in which x is a singleton; (b) if x_1, x_2 are anchors that lie in no common member, are non-adjacent and have $Q(x_1) \cap Q(x_2) = \emptyset$, and G' is G plus the edge x_1x_2, then G' has a ZE solution iff G has a ZE solution in which x_1 and x_2 are both singletons.*

Proof. The first statement: ($\Rightarrow$) as in Lemma F.2; ($\Leftarrow$) the soft partners of x are its mates in members, and these are apart from x; x has no hard edge, and its hard non-edges are unedited. (a),(b) ($\Rightarrow$): by Lemma F.2 the cluster is $\{x, x'\}$ (resp. $\{x_1, x_2\}$); split it into singletons. No other pair changes status or togetherness. ($\Leftarrow$) join the singleton to its stub (resp. join the two singletons). □

Thus an anchor costs nothing when it is a singleton in some ZE solution, and the point of a tunnel is precisely that this singleton requirement is a constraint on the member states through x: a *lock*. (A lock fixes the states of the members through x to unit values; it never forces more.)

Lemma F.4 (Rigidity at a lock)**.** *Let S be ZE. (a) If y, z are together then for every $q \notin \{y, z\}$, q is together with y iff with z. Hence if qz is hard and qy soft, the soft pair qy is forced to be together iff $qz \in E$; if both are soft, the owners of yz, qy, qz form a clique of $I(H)$ (a triangle if distinct). (b) If u, w and w, v are together, then uv is an edge or a soft pair.*

Proof. Transitivity of "together"; for (b) a hard pair inside a cluster is an unedited edge (Lemma F.1(a)). □

Consequently the two vertices that a lock bridges (the mates y, z of a locked member) are *occupied*: $y \sim z$ is a cluster that cannot also be bridged to a third vertex unless the pair is an edge or soft.

F.2 Memory across a tree decomposition

Let $(\mathcal{T}, \beta)$ be a tree decomposition of $I(H)$, $e = tt'$ an edge of $\mathcal{T}$, $\mathcal{B}_e$ the members in both bags of e, and $\mathcal{M}_1$, $\mathcal{M}_2$ the members that occur only on the side of t, resp. t'. A set S of anchors is a *face* if some ZE solution of (G, H) makes every $x \in S$ a singleton; equivalently (Lemma F.3) the partial vector of forced states is the restriction of an element of Σ. A *minimal non-face* (a *core*) is a non-face all of whose proper subsets are faces.

Lemma F.5 (Splice)**.** *(a) If $\sigma, \sigma' \in \Sigma$ agree on $\mathcal{B}_e$, the vector equal to σ on $\mathcal{M}_1 \cup \mathcal{B}_e$ and to σ' on $\mathcal{M}_2$ is in Σ. (b) Let $S = S_1 \cup S_2$ be a core with all members through anchors of S_i in $\mathcal{M}_i$ ($i = 1, 2$, both parts non-empty). For $b \in S$ let $\sigma^b \in \Sigma$ make all anchors of $S \setminus \{b\}$ singletons. Then $\{\sigma^b|_{\mathcal{B}_e} : b \in S_1\}$ and $\{\sigma^a|_{\mathcal{B}_e} : a \in S_2\}$ are disjoint.*

Proof. (a) $\sigma \in \Sigma$ iff no vertex triple is an induced P_3 after editing; the status of a triple's pairs depends on the owners of its soft pairs, a clique of $I(H)$ (Observation 3.2), so contained in one bag, hence in $\mathcal{M}_1 \cup \mathcal{B}_e$ or in $\mathcal{M}_2 \cup \mathcal{B}_e$; all constraints are satisfied by the combined vector. (b) If $\sigma^b|_{\mathcal{B}_e} = \sigma^a|_{\mathcal{B}_e}$ with $b \in S_1$, $a \in S_2$, splice σ^a on $\mathcal{M}_1 \cup \mathcal{B}_e$ with σ^b on $\mathcal{M}_2$: the anchors of S_1 are singletons (they are in $S \setminus \{a\}$, and their members are in M_1), those of S_2 likewise, so S is a face, a contradiction. □

Lemma F.6 (Majority closure bounds cores)**.** *Suppose Σ is closed under* some *majority operation $m : \{1, 2, 3\}^3 \to \{1, 2, 3\}$ applied coordinatewise ($m(a, a, b) = m(a, b, a) = m(b, a, a) = a$). Then every core has at most two anchors.*

Proof. Let S be a core with $|S| \geq 3$; we derive a contradiction.

(1) Each member contains at most one anchor of S, and as an end. If an anchor $x \in S$ is the centre of a member, x is never a singleton of a ZE solution (Lemma F.2), so $\{x\}$ is a non-face, contradicting minimality of S. If both ends a, c of a member (a, b, c) are anchors of S, then making a a singleton forces state 1 and making c a singleton forces state 2 (Lemma F.3), so $\{a, c\}$ is a non-face, again contradicting minimality, since $|S| \geq 3$. (A member can carry two end-anchors, so a pair of members can carry up to four anchors; this case analysis is what reduces the count to two.)

(2) The forced vector. Let T be the set of members through anchors of S. By Lemma F.3 and (1), the requirement that every anchor of S be a singleton fixes the state of each member of T to a single value, giving a vector $t \in \{1, 2, 3\}^T$ without conflicts, and S is a face iff $t \in \Sigma_T := \mathrm{proj}_T(\Sigma)$,

which is closed under m. The same holds for every $S' \subseteq S$ with its forced vector t' on its members T'.

(3) Binary projections determine t. Claim: if for every two members $P, P' \in T$ the pair $(t_P, t_{P'})$ lies in $\mathrm{proj}_{P,P'}(\Sigma)$, then $t \in \Sigma_T$. (This is the Baker–Pixley argument [1].) Induction on $|T|$; for $|T| \leq 2$ it is trivial. For $|T| \geq 3$ choose three members i_1, i_2, i_3 and, by induction applied to $T \setminus \{i_j\}$ and the m-closed relation $\mathrm{proj}_{T\setminus\{i_j\}}(\Sigma)$, choose $u_j \in \Sigma_T$ that agrees with t on $T \setminus \{i_j\}$. Then $m(u_1, u_2, u_3) = t$: at i_1 the vectors u_2, u_3 agree with t, similarly at i_2, i_3, and elsewhere all three agree.

(4) Since S is a non-face, $t \notin \Sigma_T$, so by (3) some pair of members $P, P' \in T$ has $(t_P, t_{P'}) \notin \mathrm{proj}_{P,P'}(\Sigma)$. Let S' be the set of anchors of S lying in P or P'; by (1), $|S'| \leq 2 < |S|$. The vector forced by S' agrees with t on P and P'. If S' were a face, some $\sigma \in \Sigma$ would extend it and $(\sigma_P, \sigma_{P'}) = (t_P, t_{P'})$, a contradiction. So S' is a non-face and a proper subset of S, contradicting that S is a core. □

F.3 Fan-out of released tunnels, and the tunnel-release model

Lemma F.7 (Released ends have small hard degree)**.** *Let H be pair-maximal, $D \geq \max_v |Q(v)|$, and S an optimal solution with $|S| \leq |H| + \ell$. (a) Every endpoint of an edited hard pair (in particular of a cut hard edge) has hard degree at most $D + \ell$. (b) Every hard component containing a vertex of hard degree larger than $2D + 3\ell$ is a whole cluster of S in which no* hard *pair is edited.*

Proof. (a) Lemma 6.4(a) gives $d_h(x) \leq D + u$ and $u \leq |S| - |H| \leq \ell$. (b) Lemma 6.6($\gamma$) with $B = 2D + 3\ell$; hard pairs inside the cluster are unedited by Lemma 6.6(β). Soft pairs inside the component may be edited (take K_9 minus one edge ac with $H = \{(a, b, c)\}$). □

Proposition F.8 (Bounded-degree Grid Tiling is easy)**.** *Grid Tiling [6] (find $a_1, \ldots, a_k, b_1, \ldots, b_k \in [n]$ with $(a_i, b_j) \in S_{ij}$ for all i, j) is solvable in time $n\, d^{2k}\,$*poly *on instances where every value has at most d partners in every S_{ij}.*

Proof. Guess a_1 (n ways); each b_j has at most d candidates, being a partner of a_1 in S_{1j}; then each a_i has at most d candidates, being a partner of b_1 in S_{i1}. □

F.4 The tunnel-release model

Fix a packing H and a set $\mathcal{J}$ of tunnels (stubs allowed). A *TR-solution* is a solution with exactly one edit in every member whose hard edits are cuts of tunnels; its excess is the number of cuts. For $Y \subseteq \mathcal{J}$ let $f(Y) = 1$ iff G minus the tunnel edges of Y has a ZE solution; call $S \subseteq \mathcal{J}$ *lockable* if $f(\mathcal{J} \setminus S) = 1$, and a minimal non-lockable set a *core*.

Proposition F.9 (Bounded cores give FPT in the model)**.** *(i) f is monotone. (ii) $f(Y) = 1$ iff Y meets every core. (iii) The minimum number of cuts of a TR-solution is the minimum size of a hitting set of the cores. (iv) If every core has at most d tunnels, this minimum is computable in time $d^\ell 3^{w+1} n^{O(1)}$ (or decides that it exceeds ℓ), given a tree decomposition of $I(H)$ of width w.*

Proof. (i) Given a ZE solution of G minus Y' and $Y' \subseteq Y$, each tunnel of $Y \setminus Y'$ is a cluster $\{x, x'\}$ (Lemma F.2); splitting it into singletons changes no edit. (ii),(iii): $f(Y) = 1$ iff $\mathcal{J} \setminus Y$ is lockable iff it contains no core, as non-lockability is upward closed; a TR-solution with cut set Y exists iff $f(Y) = 1$. (iv) Test f with the $\ell = 0$ dynamic program of Theorem 3.4; find a core inside $\mathcal{J} \setminus Y$ by deleting tunnels one at a time while the rest stays non-lockable; branch on which of its at most d tunnels is cut, to depth ℓ. □

This is a statement about a restricted model, not about CEaMP: solutions in which a member receives two edits, or a hard pair other than a tunnel edge is edited, are excluded, and they can be strictly cheaper. Inside the model, a W[1]-hardness proof therefore needs cores of unbounded size (Open Problem 7).

F.5 Examples

Example F.10 (A local flip gadget)**.** Let G^* have vertices $0, 1, 3, 4, 5, 6, 7$ and edges $01, 04, 05, 14, 15, 45, 03, 13, 36, 37, 06, 56$, with members $(6, 3, 7), (4, 1, 3), (5, 0, 3), (4, 5, 6), (6, 0, 1)$ and anchor $x = 7$ ($c = 3$; the only hard edges are 04 and 15, and 7 has none). It is valid, pair-maximal and tight (OPT $= |H| = 5$), and has exactly five ZE solutions (found by enumerating all partitions): with x a singleton, $\{0, 1, 3, 4, 5, 6\}$ and $\{0, 1, 4, 5\}, \{3, 6\}$; with x not a singleton, $\{0, 1, 4, 5, 6\}, \{3, 7\}$, $\{0, 1, 4, 5\}, \{3, 6, 7\}$ and $\{0, 1, 4, 5\}, \{3, 7\}, \{6\}$. On the pair of members $m_3 = (4, 5, 6)$, $m_1 = (4, 1, 3)$ the locked relation (anchor a singleton) is the equality on $\{2, 3\}$, while releasing the anchor admits $(3, 2)$: the stored value flips only when the tunnel at x is cut. The relation of G^* is not determined by its binary projections (22233 is pairwise consistent but not a solution).

Example F.11 (The uniform majority operation need not close Σ)**.** Let $n = 6$, $E = \{03, 05, 12, 14, 23, 24, 25, 34, 35, 45\}$ and $H = \{(0, 5, 4), (1, 2, 5), (0, 3, 2), (1, 4, 3)\}$ ($c = 2$, tight). Then $\Sigma = \{1111, 1313, 3131, 2222\}$ (by enumeration of all partitions). The uniform majority m_3 ($m_3(a, b, c) = 3$ for pairwise distinct a, b, c) maps $(1313, 3131, 2222)$ to $3333 \notin \Sigma$, so Σ is not closed under m_3. It is, however, closed under other majority operations, for instance the one with value 1 on all pairwise distinct triples, so Lemma F.6 still applies to this instance. In particular the example refutes closure under the specific operation m_3 for $c \leq 2$, not closure under every majority operation.

The object that a hardness proof through tunnels would need is a family of tight instances with bounded c and $\mathrm{tw}(I(H))$ and a core of unbounded size (Open Problem 7). Lemma F.4(b) explains why the obvious compositions of copies of G^* fail: the two vertices bridged by a lock are occupied and cannot be shared with the next gadget. By Lemma F.6, such a family needs a set Σ that is closed under no majority operation.

G Computational checks

The proofs in this paper do not depend on computation, except that the statements about explicit small instances are verified by exhaustive enumeration, as indicated where they occur. This appendix lists what was enumerated and what was tested.

Exhaustive enumeration of explicit instances. For each of the following instances all set partitions of the vertex set were enumerated and the cost of each partition computed: the instances of Remark 3.6 (domain size), Propositions 4.5 and 4.6, the shrink-rule example of Appendix A (for twin classes of size $1, \ldots, 4$), the three instances of Proposition 8.7, the family of Proposition 8.8 for $m = 3, 4, 5$, the two instances of Proposition D.10, the gadget G^* and the majority example of Appendix F. In each case the packing was checked to consist of induced P_3's with pairwise disjoint pair sets and to be pair-maximal, and the optima and the structural claims were read off from the enumeration.

Randomised tests of lemmas. The lemmas of Sections 6 and 8 and of Appendices A–D were tested on random small graphs ($n \leq 10$; random, clustered, hub and sparse graphs; random and

maximal packings) against all partitions of cost at most $|H| + 3$, with code written independently of the proofs. No violation was found: the decomposition identity, Lemma 6.2, Theorem 8.2, Corollary 8.3, Lemma 8.6, Lemmas 6.4 and 6.6, Corollary 6.7 (inserted pairs had distance at most 2 in every test), Theorem A.5, Corollary A.6, Lemmas B.1, B.4, D.1 and D.2, and Lemma 5.5. Control experiments in which a constant is weakened by one (for example $\alpha \le 2\ell + c$ in Lemma B.1, or $|C| \le \max\{D + u, 2D, 2u\}$ in Lemma B.4), or in which partitions of cost $\mathrm{OPT} + 1$ or $\mathrm{OPT} + 2$ are tested, produce violations, so the tests are not vacuous. The dynamic program of Theorem 7.3 was implemented and agreed with brute force on 2,778 pairs of an instance ($n \le 10$) and a budget ℓ. These tests are evidence for the statements; the proofs are the justification. The scripts are available from the author.

References

[1] Kirby A. Baker and Alden F. Pixley. Polynomial interpolation and the Chinese Remainder Theorem for algebraic systems. *Mathematische Zeitschrift*, 143:165–174, 1975.

[2] Sebastian Böcker. A golden ratio parameterized algorithm for cluster editing. *Journal of Discrete Algorithms*, 16:79–89, 2012.

[3] Hans L. Bodlaender. A linear-time algorithm for finding tree-decompositions of small treewidth. *SIAM Journal on Computing*, 25(6):1305–1317, 1996.

[4] Jianer Chen and Jie Meng. A $2k$ kernel for the cluster editing problem. *Journal of Computer and System Sciences*, 78(1):211–220, 2012.

[5] Christophe Crespelle, Pål Grønås Drange, Fedor V. Fomin, and Petr A. Golovach. A survey of parameterized algorithms and the complexity of edge modification. *Computer Science Review*, 48:100556, 2023.

[6] Marek Cygan, Fedor V. Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.

[7] Michael R. Fellows, Michael A. Langston, Frances A. Rosamond, and Peter Shaw. Efficient parameterized preprocessing for cluster editing. In *Fundamentals of Computation Theory (FCT 2007)*, LNCS, pages 312–321, 2007.

[8] Alexander Firbas, Alexander Dobler, Fabian Holzer, Jakob Schafellner, Manuel Sorge, Anaïs Villedieu, and Monika Wißmann. The complexity of cluster vertex splitting and company. In *SOFSEM 2024: Theory and Practice of Computer Science*, LNCS, pages 226–239, 2024.

[9] Martin Grohe and Dániel Marx. On tree width, bramble size, and expansion. *Journal of Combinatorial Theory, Series B*, 99(1):218–228, 2009.

[10] Jiong Guo. A more effective linear kernelization for cluster editing. *Theoretical Computer Science*, 410(8–10):718–726, 2009.

[11] Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *Journal of Computer and System Sciences*, 63(4):512–530, 2001.

[12] Christian Komusiewicz and Johannes Uhlmann. Cluster editing with locally bounded modifications. *Discrete Applied Mathematics*, 160(15):2259–2270, 2012.

[13] Tuukka Korhonen. A single-exponential time 2-approximation algorithm for treewidth. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 184–192, 2021.

[14] Shaohua Li, Marcin Pilipczuk, and Manuel Sorge. Cluster editing parameterized above modification-disjoint P_3-packings. *ACM Transactions on Algorithms*, 20(1):1–43, 2023. Conference version: STACS 2021, LIPIcs 187, Art. 49.

[15] René van Bevern, Vincent Froese, and Christian Komusiewicz. Parameterizing edge modification problems above lower bounds. *Theory of Computing Systems*, 62(3):739–770, 2018. Conference version: CSR 2016.