SORTIE: Fair Vehicle Routing When Route Durations Are Random.

Ex-post ordered fairness, rank prices and risk-consistent tours

Quang-Vinh Dang
British University Vietnam, Hung Yen, Vietnam
vinh.dq4@buv.edu.vn ORCID 0000-0002-3877-8024

Working paper, November 2026

Abstract

Fair vehicle routing balances drivers' workloads with ordered measures of route lengths, such as the range or the Gini mean difference. The recent exact approaches, which also require every route to be a shortest tour of its customers, assume deterministic travel times. Drivers, however, experience realised durations. We study fair routing when arc travel times are random, with a fleet-wide day factor and arc-level noise. We distinguish ex-ante fairness (of expected workloads) from ex-post fairness (the expected unfairness of the realised day, sorted day by day) and show that every deterministic fair plan overstates ex-post fairness, with an explicit bound in terms of the routes' quadratic lengths. We prove that multiplicative common shocks leave expected ordered fairness unchanged. We then give an exact, sampling-free formula for the expected value of any ordered measure of independent route durations, a Poisson–binomial order-statistic integral that also handles the negative weights of dispersion measures, and derive from it *rank prices*: the derivative of expected unfairness with respect to a shift of each route's duration. Negative rank prices identify exactly the routes a fairness objective would like to lengthen. Under uncertainty even the min-max measure can prefer a tour that is not the shortest, so route consistency must be imposed explicitly; we define it through each driver's own mean–standard deviation and prove that such tours are supported Pareto tours, computable by parametric Held–Karp. The resulting bilevel heuristic, SORTIE, screens all relocate and swap moves with the rank prices. On instances derived from the CVRPLIB benchmark used by the deterministic literature, [RESULTS SUMMARY].

Keywords: vehicle routing; fairness; stochastic travel times; ordered weighted averaging; order statistics; bilevel optimisation.

1 Introduction

Routing plans that minimise total distance often give some drivers much longer days than others. Unequal workloads hurt driver satisfaction and retention and can be seen as unfair [Matl et al., 2018]. The fairness literature in routing therefore optimises a *fairness measure* of the vector of route workloads: the longest route, the range, the Gini mean difference (GMD), the variance or a mean absolute deviation [Matl et al., 2018, Karsu and Morton, 2015, Bektaş and Letchford, 2020]. Ordered measures, which weight the sorted workloads [Yager, 1988, Ogryczak and Śliwiński, 2003, Nickel and Puerto, 2005], cover all of these in one framework [Blanco et al., 2026].

Non-monotone measures such as the range or the GMD create a well-known pathology: an optimiser can lower the measure by making a light route *longer*, giving a driver a needless detour

[Matl et al., 2018]. van Rossum et al. [2026a,b] and Ljubić et al. [2026] close this loophole exactly, by requiring every route to be a shortest (TSP-optimal) tour of its customers. Ljubić et al. [2026] write the problem as a bilevel programme in which a planner assigns customers and each driver solves a TSP, and solve instances with up to 35 customers to optimality for seven measures.

All of these works assume deterministic travel times. A driver's day, however, is a realised duration: congestion, weather, parking and service-time variability differ between routes and between days. Fairness is then a statement about random variables, and three questions arise that the deterministic theory does not ask.

1. *What should be balanced?* The expected workloads (ex-ante fairness), or the workloads drivers actually experience, sorted day by day (ex-post fairness)? The identity of the worst-off driver changes from day to day, and the two notions can disagree.

2. *How can ex-post fairness be optimised?* Sample-average approximation (SAA) is the default, but sorting per scenario makes the objective noisy and its gradient uninformative.

3. *What is a consistent route?* Under random durations "the shortest tour" is not obviously what a driver would choose, and we show that monotone measures no longer protect drivers from detours.

Contributions.

- **Model.** We formulate fair routing with random travel times (a fleet-wide day factor times arc-level noise) as a bilevel problem with an ex-post ordered fairness objective, an expected-cost budget and per-driver risk-consistent tours. With no noise it reduces to the X-CVRP-TSP of Ljubić et al. [2026].

- **Fairness illusion (Proposition 5).** For every ordered measure with nondecreasing weights, ex-ante fairness underestimates ex-post unfairness, by at most $\|\lambda\|_1 v(\sum_k W_k)^{1/2}$, where W_k is the route's quadratic length. Common multiplicative shocks leave the expected value of every ordered measure unchanged (Proposition 2); they do inflate the variance measure and the tail of unfairness.

- **PBOI (Theorem 3).** An exact, sampling-free formula for the expected ordered measure of independent route durations with given marginals (we use moment-matched gamma marginals), as an integral of a Poisson–binomial distribution against tail sums of the weights. It handles negative weights and costs $O(Gm^2)$.

- **Rank prices (Proposition 4).** The derivative of expected unfairness with respect to a location shift of a route's duration equals the expected weight of its rank. Prices of dispersion measures sum to zero, and a negative price is precisely the stochastic incentive to lengthen a route.

- **Risk-consistent tours (Proposition 6, Theorem 8).** Even $\mathbb{E}\max$ can prefer a non-shortest tour. We define consistency through the driver's own mean–standard deviation and prove that the consistent tour is a supported Pareto tour of (length, quadratic length), found exactly by dichotomic parametric Held–Karp.

- **Algorithm and evidence.** SORTIE, a bilevel local search whose moves are screened by rank prices, is compared with the deterministic fair plan, SAA and ablations on the instances used by the deterministic literature, with exact enumeration on small instances.

2 Literature review

Workload equity in deterministic routing. Matl et al. [2018] survey equity measures for routing workloads and show that their monotonicity governs whether optimal plans can contain needlessly inefficient routes; Matl et al. [2019a] study how the choice of workload resource changes balanced plans, and Matl et al. [2019b] approximate cost–balance Pareto fronts. Min-max routing has a long exact tradition [Applegate et al., 2002, Bertazzi et al., 2015], and bi-objective route balancing is solved exactly by Halvorsen-Weare and Savelsbergh [2016]; ℓ_p norms as fairness measures appear in Bektaş and Letchford [2020]. Recent work on non-monotone and ordered objectives includes branching rules for range and order-based objectives [van Rossum et al., 2024, 2026a], cutting planes that enforce TSP-optimality for the range [van Rossum et al., 2026b], route-cost equity constraints [Agius et al., 2026], Gini-bounded mTSP variants [Bhadoriya et al., 2026], fair cooperative fleets [López Sánchez et al., 2024], driver satisfaction [Bruinink et al., 2025], multi-period equity [Nekooghadirli et al., 2026] and workload/pay design for heterogeneous or crowdsourced drivers [Mahmoodian et al., 2025, Sobhanan et al., 2026]. The ordered machinery comes from location theory [Nickel and Puerto, 2005, Ogryczak, 2000, Blanco et al., 2024, Ljubić et al., 2024, Blanco et al., 2026], the ordered median TSP [Ljubić et al., 2026], OWA [Yager, 1988, Ogryczak and Śliwiński, 2003] and top-k sums [Ogryczak and Tamir, 2003]. Closest to us, Ljubić et al. [2026] propose a unified ordered framework with per-vehicle TSP-optimality through bilevel optimisation. All of these works are deterministic.

Fairness under uncertainty. In routing, Kinable et al. [2026] study a stochastic VRP with partial deliveries where fairness is among *customers* (expected fill rates) under uncertain demand, and Gao et al. [2026] combine sequential fair allocation and routing with ex-post max-min objectives across locations. Consistency in the sense of the consistent VRP (same driver, similar arrival times) is treated under stochastic customers and demands by Alvarez et al. [2024] and with robust demand by Wu et al. [2025], following Groër et al. [2009], Kovacs et al. [2015], Yu et al. [2024]; this notion differs from the TSP-optimality consistency studied here. For drivers, Anh [2022] bound workload differences in task assignment with a distributionally robust chance constraint and Lin et al. [2022] add robust workload fairness to two-echelon meal-delivery routing; gig and rideshare work considers fairness of assignment probabilities and matching under stochastic arrivals [Singh et al., 2023, Basik et al., 2021, Nanda et al., 2020, Hosseini and Schierreich, 2025]. The ex-ante/ex-post distinction is established in welfare economics [Fleurbaey and Peragine, 2012], in inequity-averse stochastic optimisation [Gutjahr, 2021], in Gini-based stochastic shelter location and humanitarian logistics [Mostajabdaveh et al., 2019, Alem et al., 2022, Mostajabdaveh et al., 2025] and in sequential allocation [Sinclair et al., 2023]; see also Shehadeh [2023], Tsang and Shehadeh [2026], Ye et al. [2024], Barak and Talgam-Cohen [2026]. To our knowledge no paper optimises an ordered measure of *random route durations*, separates its ex-ante and ex-post versions, or defines route consistency for random travel times.

Stochastic travel times. A-priori routing with random travel times goes back to Laporte et al. [1992] and Kenyon and Morton [2003], the latter with makespan-type objectives; see Gómez et al. [2016], Adulyasak and Jaillet [2016], Errico et al. [2016, 2018], Zhang et al. [2013], Ehmke et al. [2015], Osorio-Mora et al. [2025] and the surveys Oyola et al. [2018], Gendreau et al. [2016]. Correlated travel times are handled by Rostami et al. [2021], Bakach et al. [2021], Rajabi-Bahaabadi et al. [2019], Bomboi et al. [2021], robust and distributionally robust variants by Han et al. [2014], Jaillet et al. [2016], Zhang et al. [2021], Ghosal and Wiesemann [2020], and SAA by Verweij et al. [2003]. Mean–risk path choice [Nikolova and Stier-Moses, 2014, Sen et al., 2001] motivates our

driver-level consistency criterion. CVaR [Rockafellar and Uryasev, 2000, 2002] and the price of fairness [Bertsimas et al., 2011, Mo and Walrand, 2000] complete the background.

Order statistics and parametric optimisation. Order statistics of independent, non-identically distributed variables have permanent representations [Vaughan and Venables, 1972, Bapat and Beg, 1989, Balakrishnan, 2007, Glueck et al., 2008] and are related to the Poisson–binomial count of successes in independent trials [Hoeffding, 1956, Fernandez and Williams, 2010, Hong, 2013, Biscarri et al., 2018]; see David and Nagaraja [2003], Sen [1970]. Supported efficient solutions of bi-criteria problems are found by dichotomic search [Aneja and Nair, 1979, Ehrgott, 2005, Ehrgott and Gandibleux, 2000, Ralphs et al., 2006], and the TSP on few nodes by the Held–Karp recursion [Held and Karp, 1962, Bellman, 1962].

3 Model

Let $G = (V, A)$ be a complete graph with depot 0, customers $V' = \{1, \dots, n\}$, symmetric costs c_{ij} satisfying the triangle inequality, demands q_i and m identical vehicles of capacity Q. A plan consists of a partition $S_1, \dots, S_m$ of V' into nonempty capacity-feasible sets and a tour r_k of $\{0\} \cup S_k$ for each vehicle.

Random workloads. The travel time of arc (i, j) is $\tau_{ij} = Y c_{ij} \varepsilon_{ij}$, where $Y = \exp(s_Y Z - s_Y^2/2)$, $Z \sim N(0, 1)$, is a fleet-wide day factor with $\mathbb{E}Y = 1$, and the arc disturbances ε_{ij} are independent of each other and of Y with mean 1 and variance v^2 (gamma in our experiments). The workload of driver k is $\Phi_k = Y X_k$ with

$$X_k = \sum_{(i,j)\in r_k} c_{ij}\varepsilon_{ij}, \qquad \mathbb{E}X_k = L_k = \sum_{(i,j)\in r_k} c_{ij}, \qquad \operatorname{Var} X_k = \sigma_k^2 = v^2 W_k, \quad W_k = \sum_{(i,j)\in r_k} c_{ij}^2.$$

L_k is the tour length and W_k its *quadratic length*. Given Y the workloads are independent; unconditionally they are positively correlated.

Ordered fairness. For $a \in \mathbb{R}^m$ with order statistics $a_{(1)} \le \cdots \le a_{(m)}$ and weights $\lambda \in \mathbb{R}^m$, $\ell_\lambda(a) = \sum_j \lambda_j a_{(j)}$. We use the maximum ($\lambda = e_m$), the range ($e_m - e_1$), the GMD ($\lambda_j = 2(2j - m - 1)/m^2$) and the mean absolute deviation from the median (MAD; $\lambda = (-1, \dots, -1, 0, 1, \dots, 1)/m$ for odd m), together with the empirical variance $\operatorname{Var}_{\text{emp}}(a) = m^{-1} \sum_k (a_k - \bar{a})^2$. All four λ are nondecreasing.

Definition 1 (Three notions of fairness)**.** For a plan with random workloads Φ: *ex-ante* unfairness $\ell_\lambda(\mathbb{E}\Phi)$; *ex-post* unfairness $\mathbb{E}\ell_\lambda(\Phi)$; *risk of unfairness* $\operatorname{CVaR}_\beta \ell_\lambda(\Phi)$.

A deterministic fair model optimises the ex-ante notion. Drivers experience the ex-post one. SORTIE optimises ex-post unfairness and reports the CVaR out of sample.

4 Theory

Proposition 2 (Common shocks)**.** *For every* λ, $\mathbb{E}\ell_\lambda(\Phi) = \mathbb{E}\ell_\lambda(X)$, *and* $\mathbb{E}\operatorname{Var}_{\text{emp}}(\Phi) = e^{s_Y^2}\,\mathbb{E}\operatorname{Var}_{\text{emp}}(X)$.

Proof. Sorting commutes with multiplication by $y > 0$, so $\ell_\lambda(yX) = y\ell_\lambda(X)$; take expectations using independence of Y and X and $\mathbb{E}Y = 1$. The empirical variance is homogeneous of degree two and $\mathbb{E}Y^2 = e^{s_Y^2}$. □

Fleet-wide shocks therefore drop out of the planning objective for ordered measures, which reduces evaluation to the conditionally independent vector X.

Theorem 3 (Poisson–binomial order-statistic integral)**.** *Let $X_1, \dots, X_m \geq 0$ be independent with CDFs F_k. For $t \geq 0$ let $N(t) = \#\{k : X_k \leq t\}$, a Poisson–binomial count with success probabilities $F_k(t)$, and $\pi_r(t) = \mathbb{P}(N(t) = r)$. With tail sums $\Lambda_r = \sum_{j=r+1}^{m} \lambda_j$,*

$$\mathbb{E}\ell_\lambda(X) = \int_0^\infty \sum_{r=0}^{m-1} \Lambda_r \, \pi_r(t) \, dt. \tag{1}$$

Proof. $X_{(j)} > t$ if and only if fewer than j of the X_k are at most t, so $\mathbb{E}X_{(j)} = \int_0^\infty \mathbb{P}(N(t) \leq j-1)\, dt$. Hence $\mathbb{E}\ell_\lambda(X) = \int_0^\infty \sum_j \lambda_j \sum_{r \leq j-1} \pi_r(t)\, dt = \int_0^\infty \sum_r \pi_r(t) \sum_{j \geq r+1} \lambda_j \, dt$, where Tonelli's theorem is applied to the positive and negative parts of λ separately. □

The vector $\pi(t)$ is the coefficient vector of $\prod_k (1 - F_k(t) + F_k(t)z)$, obtained by m convolution steps. Below the smallest lower quantile the integrand equals Λ_0, above the largest upper quantile it vanishes ($\Lambda_m := 0$), and in between we use Gauss–Legendre panels whose breakpoints are quantiles of *every* marginal, so that a short, low-variance route is resolved even when the other routes are long. We use moment-matched gamma marginals $X_k \sim \Gamma(L_k^2/\sigma_k^2, \sigma_k^2/L_k)$. The identity behind (1) is classical for order statistics of non-identical variables [Bapat and Beg, 1989, Balakrishnan, 2007]; our contribution is to make it an exact planning objective for ordered fairness, where the tail sums Λ_r absorb the negative weights of dispersion measures, and to differentiate it.

Proposition 4 (Rank prices)**.** *Let R_k be the rank of X_k and define $\varpi_k = \sum_j \lambda_j \mathbb{P}(R_k = j)$, where $\mathbb{P}(R_k = j) = \int f_k(t)\, \mathbb{P}(N_{-k}(t) = j-1)\, dt$ and N_{-k} counts the other drivers. Then $\frac{d}{d\delta}\mathbb{E}\ell_\lambda(X + \delta e_k)\big|_{\delta=0} = \varpi_k$ and $\sum_k \varpi_k = \sum_j \lambda_j$.*

Proof. ℓ_λ is piecewise linear with $\partial \ell_\lambda / \partial a_k = \lambda_{R_k(a)}$ off the null set of ties and is $\|\lambda\|_1$-Lipschitz, so differentiation and expectation commute by dominated convergence. The ranks form a permutation, so $\sum_k \mathbb{P}(R_k = j) = 1$. □

For the range, GMD and MAD the prices sum to zero: burdened drivers carry positive prices and light drivers negative ones. A negative price is exactly the incentive to lengthen a route. Deterministically, the price of the lightest route under the range is -1, and a detour of length δ reduces the range by δ; stochastically, the incentive is spread over all drivers that might be the lightest on a given day. SORTIE uses the prices, together with $\psi_k = \partial \mathbb{E}\ell_\lambda / \partial \sigma_k$, to predict the effect of a move.

Proposition 5 (Fairness illusion)**.** *If λ is nondecreasing then ℓ_λ is convex and*

$$0 \leq \mathbb{E}\ell_\lambda(X) - \ell_\lambda(L) \leq \|\lambda\|_1 \, \mathbb{E}\|X - L\|_\infty \leq \|\lambda\|_1 \, v\Big(\sum_k W_k\Big)^{1/2}.$$

Moreover $\mathbb{E}\,\mathrm{Var}_{\mathrm{emp}}(X) = \mathrm{Var}_{\mathrm{emp}}(L) + \frac{m-1}{m^2}\sum_k \sigma_k^2$, and for Gaussian marginals $\mathbb{E}\,\mathrm{GMD}(X) = m^{-2}\sum_{i \neq j}\left[s_{ij}\sqrt{2/\pi}\, e^{-d_{ij}^2/(2s_{ij}^2)} + d_{ij}(1 - 2\Phi(-d_{ij}/s_{ij}))\right]$ with $d_{ij} = L_i - L_j$ and $s_{ij}^2 = \sigma_i^2 + \sigma_j^2$.

Proof. For nondecreasing λ the rearrangement inequality gives $\ell_\lambda(a) = \max_\sigma \sum_j \lambda_j a_{\sigma(j)}$, a maximum of linear functions; Jensen's inequality gives the lower bound. Since $|a_{(j)} - b_{(j)}| \leq \|a - b\|_\infty$ for every j, ℓ_λ is $\|\lambda\|_1$-Lipschitz in the sup norm, and $\mathbb{E}\max_k |Z_k| \leq (\mathbb{E}\sum_k Z_k^2)^{1/2}$. The variance identity follows from $\mathbb{E}X_k^2 = L_k^2 + \sigma_k^2$ and $\mathbb{E}\bar{X}^2 = \bar{L}^2 + m^{-2}\sum_k \sigma_k^2$; the GMD formula is the mean of a folded normal. □

A deterministic fair plan therefore *overstates* the fairness drivers will experience. The overstatement grows with the arc noise and with the quadratic lengths: long arcs, not long routes, are what make a balanced plan unbalanced in practice.

Proposition 6 (Monotonicity no longer protects drivers)**.** *With deterministic workloads, every monotone measure ($\lambda \geq 0$) has an optimal plan made of TSP-optimal routes [Matl et al., 2018]. With random workloads this fails even for* $\mathbb{E}\max$.

Example 7. Two drivers; $X_1 \equiv 100$. Driver 2 can drive tour A ($L = 100$, $\sigma = 10$) or tour B ($L = 101$, $\sigma \approx 0$). With Gaussian X_2, $\mathbb{E}\max = 100 + 10\varphi(0) \approx 103.99$ under A and ≈ 101 under B, so the optimiser picks the longer tour B.

Route consistency must therefore be imposed for every measure, and it needs a risk attitude. We let each driver rank the tours of their customer set by

$$\rho_\kappa(r) = \frac{\mathbb{E}[\Phi(r) \mid Y] + \kappa\,\mathrm{SD}[\Phi(r) \mid Y]}{Y} = L(r) + \kappa v\sqrt{W(r)},$$

which ranks tours identically on every day. $\kappa = 0$ is TSP-optimality (ties broken by smaller W).

Theorem 8 (Risk-consistent tours are supported)**.** *For $\kappa \geq 0$, some minimiser of ρ_κ over the tours of a fixed customer set is a supported extreme tour: it minimises $L(r) + \theta W(r)$ for some $\theta \in [0, \infty)$, or minimises W lexicographically after L ($\theta = \infty$). All supported extreme tours are enumerated by dichotomic search, each step being one Held–Karp run with arc weights $c_{ij} + \theta c_{ij}^2$.*

Proof. $f(L, W) = L + \kappa v\sqrt{W}$ is concave and nondecreasing on $\mathbb{R}^2_+$. A concave function bounded below on a pointed polyhedron attains its minimum at an extreme point; since f is nondecreasing, the minimum of f over the finite set $P = \{(L(r), W(r))\}$ equals its minimum over $\mathrm{conv}(P) + \mathbb{R}^2_+$ and is attained at an extreme point of that set. These extreme points are exactly the unique minimisers of $w_1 L + w_2 W$ with $(w_1, w_2) \geq 0$, $(w_1, w_2) \neq 0$ (for $w_1 = 0$ with ties broken by smaller L), and dichotomic search enumerates all of them [Aneja and Nair, 1979]. Tours in the relative interior of a hull edge are dominated by the edge's end points and need not be enumerated. With heterogeneous noise (ε_{ij} with variance $v^2 H_{ij}^2$) the same holds with $W = \sum (H_{ij} c_{ij})^2$ and arc weights $c_{ij} + \theta H_{ij}^2 c_{ij}^2$. □

The SORTIE problem.

$$\begin{aligned} \min\ & \mathbb{E}\,\ell_\lambda\big(X(r_1, \ldots, r_m)\big) \\ \text{s.t.}\ & (S_1, \ldots, S_m) \text{ capacity-feasible partition},\ S_k \neq \emptyset, \qquad \textstyle\sum_k L(r_k) \leq B \cdot C^*, \\ & r_k \in \arg\min\{\rho_\kappa(r) : r \text{ a tour of } \{0\} \cup S_k\} \quad \forall k, \end{aligned}$$

where C^* is the minimum expected total length and $B \geq 1$ the budget factor. With $v = 0$ and $\kappa = 0$ this is the X-CVRP-TSP of Ljubić et al. [2026].

5 The SORTIE algorithm

The lower level is exact for routes of up to 15 customers (numba Held–Karp, memoised by customer set, about 4 ms at 13 customers) and uses 2-opt/Or-opt from several starts beyond that. The evaluator costs about 0.5 ms per plan for $m = 5$. A full screen of all relocate and swap moves costs a few milliseconds at $n = 25$ and under 1 ms per route-pair precomputation at $n = 100$, so the exact evaluations, not the screen, dominate.

Algorithm 1 SORTIE

$P \leftarrow$ cost-efficient plan; $P^* \leftarrow P$
while time remains **do**
    compute rank prices ϖ_k and dispersion prices ψ_k of P (Prop. 4)
    score every relocate and swap move by $\sum_{k\in\{a,b\}} \varpi_k \Delta L_k + \psi_k \Delta\sigma_k$ (insertion deltas)
    evaluate the K best-scored budget- and capacity-feasible moves exactly: lower-level tours (Thm. 8) and PBOI (Thm. 3)
    if one improves P **then** apply the best
    else ruin (price-weighted or related removal) and recreate (price-scored regret-2); repair cost if over budget; accept by simulated annealing
    end if
    update P^*
end while
return P^*

6 Computational study

6.1 Set-up

Instances. We follow the slicing rule of the deterministic fair routing literature [Ljubić et al., 2026]. Block b of size n consists of the depot and the n consecutive customers $bn+1, \dots, (b+1)n$ of CVRPLIB X-n641-k35 [Uchoa et al., 2017], with ceiled Euclidean distances, $m = 5$ vehicles and capacity $Q = \lfloor \sum_i q_i/4 \rfloor$, so one vehicle's worth of slack is available. The main grid uses $n \in \{15, 20, 25, 30, 35\}$ and blocks $b = 0, \dots, 4$ (25 instances); the scale test uses $n \in \{50, 75, 100\}$ and blocks 0–2; the exactness test uses ten instances with $n = 10$, $m = 3$ and $Q = \lfloor \sum_i q_i/2.4 \rfloor$.

Uncertainty. Unless stated otherwise $v = 0.3$ (gamma arc noise) and $s_Y = 0.15$. In the congestion experiment (E8) the arc coefficient of variation is vH_{ij}, where $H_{ij} \in [1/3, 2]$ is largest for arcs near a randomly drawn congestion centre, so noise is no longer proportional to length.

Budget. C^* is the cost of the best of three restarts of the cost-minimising search; the total length of every fair plan must satisfy $\sum_k L_k \le B\, C^*$ with $B \in \{1.05, 1.10\}$.

Methods. All methods share the search of Algorithm 1, the start plan, the time limit (15 s for $n \le 25$, 25 s for $n \ge 30$, 60 s for $n \ge 50$) and the random seed. They differ only in the objective and in the lower level. *Det* minimises the deterministic ordered measure $\ell_\lambda(L)$ of the tour lengths, i.e. the model of Ljubić et al. [2026] solved heuristically. *SAA* minimises the sample average of $\ell_\lambda(X)$ over 200 fixed arc scenarios. *SORTIE-0* minimises the exact ex-post value by PBOI with TSP-optimal tours ($\kappa = 0$); *SORTIE-1* uses the risk-consistent lower level with $\kappa = 1$. In Det and SAA the moves are screened by the deterministic and the sample objective respectively.

Evaluation. Every plan is evaluated out of sample by arc-level Monte Carlo with 20 000 scenarios (10 000 for $n \ge 50$), including the day factor, with common random numbers across methods. The reported *ex-post value* is the Monte Carlo estimate of $\mathbb{E}\ell_\lambda(\Phi)$. Paired comparisons use the Wilcoxon signed-rank test. All tables are generated from the raw results by the released scripts.

Table 1: E1: PBOI against arc-level Monte Carlo (50 000 samples) on 168 plans; z: difference over its Monte Carlo standard error.

Measure	Plans	Mean |err| (%)	Max |err| (%)	Share $\|z\| < 3$
Max	168	0.041	0.137	0.994
Range	168	0.063	0.249	1.000
GMD	168	0.058	0.241	1.000
MAD	168	0.065	0.272	0.994
Var	168	0.109	0.440	1.000

Table 2: E3: what each method believes its unfairness is, relative to the true ex-post value (Monte Carlo without the day factor, which is what each model represents), mean signed error over 50 runs per cell. Last column: largest absolute error of the PBOI prediction for any method's plan.

	Believed vs true (mean signed %)				PBOI
Measure	Det	SAA	SORTIE-0	SORTIE-1	max |err| %
Max	-8.00	-0.69	-0.01	-0.01	0.20
Range	-16.97	-0.83	+0.02	+0.02	0.60
GMD	-14.63	-0.83	+0.04	+0.05	0.61
MAD	-16.25	-0.99	+0.05	+0.07	0.67
Var	-26.52	-1.57	+0.00	+0.04	0.98

6.2 Accuracy and cost of the evaluator (E1, E2)

Table 1 compares PBOI with 50 000-sample arc-level Monte Carlo on 168 plans: the cost-efficient plans and random feasible plans whose routes are sequenced by the lower level, for $n \in \{15, 25, 35, 50\}$. The mean absolute relative error is below 0.11% for all five measures and below 0.44% in the worst case, and 99–100% of the differences lie within three Monte Carlo standard errors. The mean z-scores are not exactly zero (up to 0.34 for the MAD): PBOI carries a systematic bias of a few hundredths of a percent, which we attribute to the moment-matched gamma marginal of each route (a sum of arc gammas with different scales is not exactly gamma). PBOI takes 0.46 ms per plan for $m = 5$, against 0.28 s for 20 000-scenario Monte Carlo. A 200-scenario SAA is about seven times faster than PBOI but is a noisy estimate. The advantage of PBOI is therefore that it is exact and noise-free at a cost compatible with thousands of evaluations per second, not raw speed.

On the ten $n = 10$ instances we enumerated all capacity- and budget-feasible partitions (4 to 145 per instance) with exact Held–Karp tours. The cost search found the optimal C^* on all ten. For both range and GMD, SORTIE-0 found the plan with the smallest exact ex-post value on nine of ten instances (mean gap 0.39%, maximum 3.9%). Det found the deterministic optimum on nine of ten; on the tenth instance, which has only 20 budget-feasible partitions, it stopped at a plan with twice the optimal deterministic range.

6.3 The fairness illusion and the value of ex-post optimisation (E3)

Table 2 quantifies Proposition 5. A deterministic fair plan believes its unfairness is the ex-ante value; the truth is 8% larger for the maximum, 15–17% larger for the range, GMD and MAD, and 27% larger for the variance. The in-sample SAA value is optimistic by about 1%, which is the optimiser's curse on 200 scenarios. The PBOI value believed by SORTIE has a mean signed error below 0.1%, a mean absolute error of 0.07–0.23% and a worst case of 1%.

Table 3 reports the out-of-sample ex-post value relative to Det. SORTIE-0 and SORTIE-1 each

Table 3: E3: improvement of the out-of-sample ex-post value over Det (mean %, positive is better; in brackets the number of instances on which the method is strictly better). *: Wilcoxon signed-rank $p < 0.05$ in favour of the method; the test is on ranks and can disagree in sign with the mean when one instance dominates (Var, $B = 1.05$, SORTIE-1).

Measure	B	SAA	SORTIE-0	SORTIE-1
GMD	1.05	+0.66 (8/25)	+0.26* (7/25)	-0.43 (8/25)
GMD	1.10	+0.20 (9/25)	+0.74* (12/25)	+0.62* (16/25)
MAD	1.05	+0.98* (10/25)	+1.30* (13/25)	+1.48* (13/25)
MAD	1.10	+0.77 (12/25)	+1.21* (13/25)	+1.42* (20/25)
Max	1.05	+0.77* (15/25)	+0.63* (15/25)	+0.82* (15/25)
Max	1.10	+1.71* (23/25)	+2.11* (24/25)	+1.96* (24/25)
Range	1.05	+0.55 (12/25)	-0.15 (13/25)	+0.74 (13/25)
Range	1.10	+0.96* (14/25)	+1.34* (16/25)	+1.31* (16/25)
Var	1.05	-0.45 (7/25)	-0.36 (8/25)	-0.01* (12/25)
Var	1.10	-0.17 (6/25)	+0.68* (12/25)	+1.31* (18/25)




Figure 1: E3, $B = 1.10$: per-instance improvement of the ex-post value over Det.

have a positive mean improvement over Det in 8 of the 10 cells, and the signed-rank test favours them in 8 cells each. Pooled over all 250 runs, SORTIE-0 improves on Det by 0.77% on average (better in 133 runs, tied in 102, $p < 10^{-16}$) and SORTIE-1 by 0.92% (better in 155, tied in 69). SAA improves by 0.60% and is the least consistent: its median improvement is zero, it is favoured significantly in only 4 cells, and it is worse than Det for the variance. Head to head, SORTIE-0 beats SAA in 138 and SORTIE-1 in 150 of 250 runs ($p < 10^{-14}$). The largest gains are for the maximum at $B = 1.10$ (2.1%, 24 of 25 instances), where the extra budget lets the planner trade a slightly longer but less variable longest route.

The improvements are small, about one percent on average, and this is itself a finding. With homogeneous noise ($\sigma_k^2 = v^2 W_k$) the standard deviation of a route grows with its length, so the plans that balance lengths also nearly balance risks, and a deterministic fair plan is close to ex-post optimal even though its *assessment* of its own fairness is badly wrong. For context, the cost-optimal plan has an ex-post range of 2075 against about 1340 for the fair plans, at a cost 6.6% higher: fairness itself is worth optimising; the stochastic refinement adds about one percent on top. For range-optimised plans the $\mathrm{CVaR}_{0.9}$ of the realised range, which none of the methods optimises, is 2054 for Det, 2046 for SAA, 2047 for SORTIE-0 and 2034 for SORTIE-1; only SORTIE-1 improves it at both budgets (SORTIE-0 is slightly worse than Det at $B = 1.05$).

Table 4: E4: free tours against lower-level tours, $B = 1.10$, 25 instances per row. Excess length: total length above the TSP tours of the same customer sets.

Measure	Method	Runs w/ non-TSP	Non-TSP routes	Excess length	Belief err (%)	Ex-post value
Max	det-lower	0/25	0.00	0.0	-8.92	2220.8
Max	det-free	0/25	0.00	0.0	-8.96	2225.7
Max	pboi-lower	0/25	0.00	0.0	+0.00	2175.3
Max	pboi-free	9/25	0.48	3.7	+0.00	2179.5
Range	det-lower	0/25	0.00	0.0	-22.16	1107.4
Range	det-free	23/25	1.00	91.9	-23.55	1084.8
Range	pboi-lower	0/25	0.00	0.0	+0.05	1092.8
Range	pboi-free	25/25	1.68	114.7	+0.05	1069.4
GMD	det-lower	0/25	0.00	0.0	-18.82	435.7
GMD	det-free	21/25	1.08	87.4	-19.01	428.0
GMD	pboi-lower	0/25	0.00	0.0	+0.08	433.2
GMD	pboi-free	23/25	1.36	78.9	+0.07	438.3

6.4 Route consistency (E4)

E4 removes the lower level: the planner may sequence each route freely (*free*) instead of using the TSP-optimal tour (*lower*). For the maximum, free sequencing changes almost nothing, as monotonicity predicts in the deterministic case. For the range and the GMD the free planner lengthens the short routes: 23 of 25 Det plans and all 25 SORTIE plans for the range contain a route that is not a shortest tour of its customers, on average by 92–115 distance units. This lowers the ex-post value by about 2% for both methods on the range and for Det on the GMD (Table 4); for SORTIE on the GMD it is 1.2% worse, because the larger space slows the search. Wherever it helps, the reduction comes from making some drivers drive detours. This is exactly the behaviour signalled by negative rank prices and excluded by the bilevel constraint. The deterministic belief error persists with free tours (-24% for the range); the PBOI belief remains exact.

6.5 Rank-price screening (E5)

Replacing the price screen by $K = 8$ random feasible moves, with everything else unchanged, gives a similar number of iterations but worse plans: price screening lowers the ex-post value by 6.6% for the range and 7.2% for the GMD on average (medians 1.7% and 2.5%; better on 14 and 15 of 25 instances, worse on 1 and 0, ties otherwise; $p < 10^{-3}$). The mean is driven by a few instances on which random screening stalls far from a good plan. The rank prices are thus an effective first-order guide for the search, and not only an interpretive device.

6.6 Noise level (E6)

Table 5 varies v for $n = 25$. The illusion grows monotonically with v, as the bound of Proposition 5 suggests: at $v = 0.5$ a deterministic plan understates its range by 43%. The gain of SORTIE over Det also grows with v, from 0.25% at $v = 0.1$ to 1.75% at $v = 0.5$ for the range (better on all five instances).

6.7 Larger instances (E7)

Table 6 reports $n \in \{50, 75, 100\}$ with a 60 s limit. Here the routes have 10 to 20 customers on average, so most lower-level calls fall back to local search, and the number of iterations drops from

Table 5: E6: arc noise v, $n = 25$, $B = 1.10$, five instances per row.

v	Measure	Det belief err (%)	SORTIE-0 vs Det
0.1	GMD	-6.4	+0.48 (1/5)
0.1	Range	-8.7	+0.25 (2/5)
0.3	GMD	-22.0	+0.64 (1/5)
0.3	Range	-28.7	+0.96 (3/5)
0.5	GMD	-35.8	+1.50 (4/5)
0.5	Range	-42.5	+1.75 (5/5)




Figure 2: Left: range believed by Det (ex-ante) and by SORTIE (PBOI) against the true expected range, E3. Right: share of the true ex-post unfairness that a deterministic plan does not see, against v (E6).

about 1500–2000 at $n = 50$ to 100–400 at $n \geq 75$. The deterministic belief error stays at about -22% for all three methods, and PBOI remains accurate (mean absolute error 0.2%, at most 1%). The comparison of the methods, however, is dominated by search noise: individual differences from Det range from -65% to $+40\%$. SORTIE-0 is better than Det on 5 of 6 runs at $n = 50$ and at $n = 100$ but on only 1 of 6 at $n = 75$, where on one instance it ends at an ex-post range 65% worse than Det; SAA is just as erratic. We read E7 as showing that the evaluator scales but the bilevel search does not yet: the bottleneck at $n \geq 75$ is the lower level, not the fairness evaluation, and a faster lower level (or longer runs) is needed before conclusions can be drawn about the objective at this size.

7 Conclusions

Fair routing under random travel times has to say whose fairness it measures. Drivers experience realised durations, so the relevant quantity is the expected unfairness of the realised day. A deterministic fair plan cannot see this quantity, and in our experiments it understated it by 8% (maximum) to 27% (variance) at $v = 0.3$, and by up to 43% at $v = 0.5$. A multiplicative common shock, by contrast, leaves the expected value of every ordered measure unchanged (Proposition 2), although it does inflate the variance measure and the tail of unfairness.

Table 6: E7: improvement of the ex-post value over Det, $B = 1.10$, 60 s, three instances per row (format as Table 3).

n	Measure	SAA	SORTIE-0
50	GMD	+7.31 (1/3)	+7.75 (3/3)
50	Range	+7.73 (2/3)	+2.50 (2/3)
75	GMD	-2.25 (1/3)	+0.54 (1/3)
75	Range	-9.06 (0/3)	-22.45 (0/3)
100	GMD	+14.19 (3/3)	+7.84 (2/3)
100	Range	-3.91 (2/3)	+9.22 (3/3)

The Poisson–binomial order-statistic integral evaluates any ordered measure of independent route durations exactly, in under a millisecond for five vehicles, and its derivatives, the rank prices, are both interpretable and useful: they identify the routes a fairness objective would like to lengthen, and screening moves with them gave plans 7% better on average (2% at the median) than random screening. Under uncertainty, route consistency no longer follows from monotonicity, so it has to be part of the model; the mean–standard-deviation lower level makes it computable by parametric Held–Karp. Without a lower level, both the deterministic and the stochastic planners produced detours in almost every range-optimised plan.

Optimising ex-post fairness improved the realised fairness by about one percent over a deterministic fair plan under homogeneous noise (up to 2% for the maximum), consistently but modestly, because balancing route lengths already roughly balances route risks when variance grows with length.

Limitations. The lower level is exact only for routes of up to 15 customers; beyond that it relies on local search, and the bilevel constraint is then satisfied only approximately. PBOI assumes conditionally independent route durations; dependence beyond a common multiplicative factor (for example shared congested arcs) would need the copula or simulation layer we did not develop. The time limits are per run and are slightly exceeded by the final iteration. All instances are slices of one benchmark file with five vehicles.

Future work. Exact methods for the stochastic bilevel problem (branch-and-cut with PBOI-based bounds), correlated route durations, multi-day fairness, in which unfairness can be compensated across days, and time-dependent travel times.

References

Yossiri Adulyasak and Patrick Jaillet. Models and Algorithms for Stochastic and Robust Vehicle Routing with Deadlines. *Transportation Science*, 50(2):608–626, 2016. doi: 10.1287/trsc.2014.0581.

Maxime Agius, Nabil Absi, Dominique Feillet, and Thierry Garaix. Solution algorithms for a vehicle routing problem with route-cost equity constraints. *Annals of Operations Research*, 359 (3):2309–2349, 2026. doi: 10.1007/s10479-026-07145-6.

Douglas Alem, Aakil M. Caunhye, and Alfredo Moreno. Revisiting Gini for equitable humanitarian logistics. *Socio-Economic Planning Sciences*, 82:101312, 2022. doi: 10.1016/j.seps.2022.101312.

Aldair Alvarez, Jean-François Cordeau, and Raf Jans. The consistent vehicle routing problem with stochastic customers and demands. *Transportation Research Part B: Methodological*, 186:102968, 2024. doi: 10.1016/j.trb.2024.102968.

Yash P. Aneja and Kunhiraman P. K. Nair. Bicriteria Transportation Problem. *Management Science*, 25(1):73–78, 1979. doi: 10.1287/mnsc.25.1.73.

Nguyen Duy Anh. Workload Balancing Among Heathcare Workers Under Uncertain Service Time Using Distributionally Robust Optimization. *Asia-Pacific Journal of Operational Research*, 39 (05):2150045, 2022. doi: 10.1142/s0217595921500457.

David Applegate, William Cook, Sanjeeb Dash, and André Rohe. Solution of a Min-Max Vehicle Routing Problem. *INFORMS Journal on Computing*, 14(2):132–143, 2002. doi: 10.1287/ijoc.14.2.132.118.

Iurii Bakach, Ann Melissa Campbell, Jan Fabian Ehmke, and Timothy L. Urban. Solving vehicle routing problems with stochastic and correlated travel times and makespan objectives. *EURO Journal on Transportation and Logistics*, 10:100029, 2021. doi: 10.1016/j.ejtl.2021.100029.

N. Balakrishnan. Permanents, Order Statistics, Outliers, and Robustness. *Revista Matemática Complutense*, 20(1):7–107, 2007. doi: 10.5209/rev_rema.2007.v20.n1.16528.

R. B. Bapat and M. I. Beg. Order statistics for nonidentically distributed variables and permanents. *Sankhyā: The Indian Journal of Statistics, Series A*, 51(1):79–93, 1989. No DOI; JSTOR 25050725 (id as listed in search results; indexed in IAS repository).

Zohar Barak and Inbal Talgam-Cohen. Ex-ante versus Ex-post: Egalitarian Facility Location Mechanism Design. arXiv preprint, 2026.

Fuat Basik, Bugra Gedik, Hakan Ferhatosmanoglu, and Kun-Lung Wu. Fair Task Allocation in Crowdsourced Delivery. *IEEE Transactions on Services Computing*, 14(4):1040–1053, 2021. doi: 10.1109/tsc.2018.2854866.

Tolga Bektaş and Adam N. Letchford. Using ℓ_p-norms for fairness in combinatorial optimisation. *Computers & Operations Research*, 120:104975, 2020. doi: 10.1016/j.cor.2020.104975.

Richard Bellman. Dynamic Programming Treatment of the Travelling Salesman Problem. *Journal of the ACM*, 9(1):61–63, 1962. doi: 10.1145/321105.321111.

Luca Bertazzi, Bruce Golden, and Xingyin Wang. Min–Max vs. Min–Sum Vehicle Routing: A worst-case analysis. *European Journal of Operational Research*, 240(2):372–381, 2015. doi: 10.1016/j.ejor.2014.07.025.

Dimitris Bertsimas, Vivek F. Farias, and Nikolaos Trichakis. The Price of Fairness. *Operations Research*, 59(1):17–31, 2011. doi: 10.1287/opre.1100.0865.

Abhay Singh Bhadoriya, Deepjyoti Deka, and Kaarthik Sundar. Equitable routing–rethinking the Multiple Traveling Salesman Problem. *Journal of the Operational Research Society*, pages 1–20, 2026. doi: 10.1080/01605682.2026.2708027. Published online 2026; first posted on arXiv April 2024.

William Biscarri, Sihai Dave Zhao, and Robert J. Brunner. A simple and fast method for computing the Poisson binomial distribution function. *Computational Statistics & Data Analysis*, 122:92–100, 2018. doi: 10.1016/j.csda.2018.01.007.

Víctor Blanco, Alfredo Marín, and Justo Puerto. Intra-facility equity in discrete and continuous p-facility location problems. *Computers & Operations Research*, 162:106487, 2024. doi: 10.1016/j.cor.2023.106487.

Víctor Blanco, Miguel A. Pozo, Justo Puerto, and Alberto Torrejon. Optimization-based computation and integration of ordered measures. *Annals of Operations Research*, 364(3):2799–2834, 2026. doi: 10.1007/s10479-026-07314-7.

Federica Bomboi, Christoph Buchheim, and Jonas Pruente. On the stochastic vehicle routing problem with time windows, correlated travel times, and time dependency. *4OR*, 20(2):217–239, 2021. doi: 10.1007/s10288-021-00476-z.

Tom Bruinink, Lotte Berghman, and Twan Dollevoet. Beyond efficiency: Exploring the cost effects of prioritizing driver satisfaction in vehicle routing. *Operations Research Letters*, 63:107344, 2025. doi: 10.1016/j.orl.2025.107344.

H. A. David and H. N. Nagaraja. *Order Statistics*. Wiley Series in Probability and Statistics. Wiley, 3 edition, 2003. doi: 10.1002/0471722162.

Jan Fabian Ehmke, Ann Melissa Campbell, and Timothy L. Urban. Ensuring service levels in routing problems with time windows and stochastic travel times. *European Journal of Operational Research*, 240(2):539–550, 2015. doi: 10.1016/j.ejor.2014.06.045.

Matthias Ehrgott. *Multicriteria Optimization*. Springer, Berlin, Heidelberg, 2 edition, 2005. doi: 10.1007/3-540-27659-9. Author field absent in DOI metadata; ISBN 3540213988.

Matthias Ehrgott and Xavier Gandibleux. A survey and annotated bibliography of multiobjective combinatorial optimization. *OR Spectrum*, 22(4):425–460, 2000. doi: 10.1007/s002910000046.

F. Errico, G. Desaulniers, M. Gendreau, W. Rei, and L.-M. Rousseau. A priori optimization with recourse for the vehicle routing problem with hard time windows and stochastic service times. *European Journal of Operational Research*, 249(1):55–66, 2016. doi: 10.1016/j.ejor.2015.07.027.

F. Errico, G. Desaulniers, M. Gendreau, W. Rei, and L.-M. Rousseau. The vehicle routing problem with hard time windows and stochastic service times. *EURO Journal on Transportation and Logistics*, 7(3):223–251, 2018. doi: 10.1007/s13676-016-0101-4.

Manuel Fernandez and Stuart Williams. Closed-Form Expression for the Poisson-Binomial Probability Density Function. *IEEE Transactions on Aerospace and Electronic Systems*, 46(2):803–817, 2010. doi: 10.1109/taes.2010.5461658.

Marc Fleurbaey and Vito Peragine. Ex Ante Versus Ex Post Equality of Opportunity. *Economica*, 80(317):118–130, 2012. doi: 10.1111/j.1468-0335.2012.00941.x.

Haiqing Gao, Seyed M. R. Iravani, and Sean R. Sinclair. Sequential Fair Allocation and Routing in Nonprofit Operations. arXiv preprint, 2026.

Michel Gendreau, Ola Jabali, and Walter Rei. 50th Anniversary Invited Article—Future Research Directions in Stochastic Vehicle Routing. *Transportation Science*, 50(4):1163–1173, 2016. doi: 10.1287/trsc.2016.0709.

Shubhechyya Ghosal and Wolfram Wiesemann. The Distributionally Robust Chance-Constrained Vehicle Routing Problem. *Operations Research*, 68(3):716–732, 2020. doi: 10.1287/opre.2019.1924.

D. H. Glueck, A. Karimpour-Fard, J. Mandel, L. Hunter, and K. E. Muller. Fast Computation by Block Permanents of Cumulative Distribution Functions of Order Statistics from Several Populations. *Communications in Statistics - Theory and Methods*, 37(18):2815–2824, 2008. doi: 10.1080/03610920802001896.

Chris Groër, Bruce Golden, and Edward Wasil. The Consistent Vehicle Routing Problem. *Manufacturing & Service Operations Management*, 11(4):630–643, 2009. doi: 10.1287/msom.1080.0243.

Walter J. Gutjahr. Inequity-averse stochastic decision processes. *European Journal of Operational Research*, 288(1):258–270, 2021. doi: 10.1016/j.ejor.2020.05.045.

Andrés Gómez, Ricardo Mariño, Raha Akhavan-Tabatabaei, Andrés L. Medaglia, and Jorge E. Mendoza. On Modeling Stochastic Travel and Service Times in Vehicle Routing. *Transportation Science*, 50(2):627–641, 2016. doi: 10.1287/trsc.2015.0601.

Elin E. Halvorsen-Weare and Martin W.P. Savelsbergh. The bi-objective mixed capacitated general routing problem with different route balance criteria. *European Journal of Operational Research*, 251(2):451–465, 2016. doi: 10.1016/j.ejor.2015.11.024.

Jinil Han, Chungmok Lee, and Sungsoo Park. A Robust Scenario Approach for the Vehicle Routing Problem with Uncertain Travel Times. *Transportation Science*, 48(3):373–390, 2014. doi: 10.1287/trsc.2013.0476.

Michael Held and Richard M. Karp. A Dynamic Programming Approach to Sequencing Problems. *Journal of the Society for Industrial and Applied Mathematics*, 10(1):196–210, 1962. doi: 10.1137/0110015.

Wassily Hoeffding. On the Distribution of the Number of Successes in Independent Trials. *The Annals of Mathematical Statistics*, 27(3):713–721, 1956. doi: 10.1214/aoms/1177728178.

Yili Hong. On computing the distribution function for the Poisson binomial distribution. *Computational Statistics & Data Analysis*, 59:41–51, 2013. doi: 10.1016/j.csda.2012.10.006.

Hadi Hosseini and Šimon Schierreich. The Algorithmic Landscape of Fair and Efficient Distribution of Delivery Orders in the Gig Economy. arXiv preprint, 2025.

Patrick Jaillet, Jin Qi, and Melvyn Sim. Routing Optimization Under Uncertainty. *Operations Research*, 64(1):186–200, 2016. doi: 10.1287/opre.2015.1462.

Özlem Karsu and Alec Morton. Inequity averse optimization in operational research. *European Journal of Operational Research*, 245(2):343–359, 2015. doi: 10.1016/j.ejor.2015.02.035.

Astrid S. Kenyon and David P. Morton. Stochastic Vehicle Routing with Random Travel Times. *Transportation Science*, 37(1):69–82, 2003. doi: 10.1287/trsc.37.1.69.12820.

Joris Kinable, Natasja Sluijk, Michel Gendreau, Walter Rei, and Tom Van Woensel. Fair Stochastic Vehicle Routing with Partial Deliveries. *Transportation Science*, 60(2):264–283, 2026. doi: 10.1287/trsc.2024.0556.

Attila A. Kovacs, Bruce L. Golden, Richard F. Hartl, and Sophie N. Parragh. The Generalized Consistent Vehicle Routing Problem. *Transportation Science*, 49(4):796–816, 2015. doi: 10.1287/trsc.2014.0529.

Gilbert Laporte, François Louveaux, and Hélène Mercure. The Vehicle Routing Problem with Stochastic Travel Times. *Transportation Science*, 26(3):161–170, 1992. doi: 10.1287/trsc.26.3.161.

Shaochong Lin, Monique Bakker, and Eman Leung. Robust meal delivery service for the elderly: a case study in Hong Kong. *Scientific Reports*, 12(1):21546, 2022. doi: 10.1038/s41598-022-25924-6.

Ivana Ljubić, Justo Puerto, and Alberto Torrejón. Fair Vehicle Routing via Bilevel Optimization. Optimization Online preprint, submitted 5 March 2026, 2026. URL https://optimization-online.org/2026/03/fair-vehicle-routing-via-bilevel-optimization/.

Ivana Ljubić, Miguel A. Pozo, Justo Puerto, and Alberto Torrejón. Benders decomposition for the discrete ordered median problem. *European Journal of Operational Research*, 317(3):858–874, 2024. doi: 10.1016/j.ejor.2024.04.030.

Ivana Ljubić, Alfredo Marín, Justo Puerto, and Francisco Temprano. Ordered Median Traveling Salesman Problem. *Networks*, 88(1):59–84, 2026. doi: 10.1002/net.70036.

Aitor López Sánchez, Marin Lujak, Frederic Semet, and Holger Billhardt. How to achieve fair and efficient cooperative vehicle routing? *AI Communications*, 37(2):223–245, 2024. doi: 10.3233/aic-220315.

Vahid Mahmoodian, Hadi Charkhgard, and Iman Dayarian. Equitable Workload Allocation in Vehicle Routing Problem With Heterogeneous Drivers. *Production and Operations Management*, 34(7):1875–1900, 2025. doi: 10.1177/10591478241305873.

P. Matl, R. F. Hartl, and T. Vidal. Workload Equity in Vehicle Routing Problems: A Survey and Analysis. *Transportation Science*, 52(2):239–260, 2018. doi: 10.1287/trsc.2017.0744.

P. Matl, R.F. Hartl, and T. Vidal. Workload equity in vehicle routing: The impact of alternative workload resources. *Computers & Operations Research*, 110:116–129, 2019a. doi: 10.1016/j.cor.2019.05.016.

Piotr Matl, Richard F. Hartl, and Thibaut Vidal. Leveraging single-objective heuristics to solve bi-objective problems: Heuristic box splitting and its application to vehicle routing. *Networks*, 73(4):382–400, 2019b. doi: 10.1002/net.21876.

J. Mo and J. Walrand. Fair end-to-end window-based congestion control. *IEEE/ACM Transactions on Networking*, 8(5):556–567, 2000. doi: 10.1109/90.879343.

Mahdi Mostajabdaveh, Walter J. Gutjahr, and F. Sibel Salman. Inequity-averse shelter location for disaster preparedness. *IISE Transactions*, 51(8):809–829, 2019. doi: 10.1080/24725854.2018.1496372.

Mahdi Mostajabdaveh, F. Sibel Salman, and Walter J. Gutjahr. A branch-and-price algorithm for fast and equitable last-mile relief aid distribution. *European Journal of Operational Research*, 324(2):522–537, 2025. doi: 10.1016/j.ejor.2025.01.032.

Vedant Nanda, Pan Xu, Karthik Abhinav Sankararaman, John Dickerson, and Aravind Srinivasan. Balancing the Tradeoff between Profit and Fairness in Rideshare Platforms during High-Demand Hours. In *Proceedings of the AAAI Conference on Artificial Intelligence*, pages 2210–2217. Association for the Advancement of Artificial Intelligence (AAAI), 2020. doi: 10.1609/aaai.v34i02.5597.

Najmeh Nekooghadirli, Michel Gendreau, Jean-Yves Potvin, and Thibaut Vidal. Workload equity in multiperiod vehicle routing problems. *International Transactions in Operational Research*, 33 (1):245–267, 2026. doi: 10.1111/itor.70012. Published online 2025; volume 33 (2026).

Stefan Nickel and Justo Puerto. *Location Theory: A Unified Approach*. Springer, Berlin, Heidelberg, 2005. doi: 10.1007/3-540-27640-8. Hardcover 2005; eBook 2006 (cited as 2006 in Ljubić et al.).

E. Nikolova and N. E. Stier-Moses. A Mean-Risk Model for the Traffic Assignment Problem with Stochastic Travel Times. *Operations Research*, 62(2):366–382, 2014. doi: 10.1287/opre.2013.1246.

Wlodzimierz Ogryczak and Arie Tamir. Minimizing the sum of the k largest functions in linear time. *Information Processing Letters*, 85(3):117–122, 2003. doi: 10.1016/s0020-0190(02)00370-8.

Włodzimierz Ogryczak. Inequality measures and equitable approaches to location problems. *European Journal of Operational Research*, 122(2):374–391, 2000. doi: 10.1016/s0377-2217(99)00240-4.

Włodzimierz Ogryczak and Tomasz Śliwiński. On solving linear programs with the ordered weighted averaging objective. *European Journal of Operational Research*, 148(1):80–91, 2003. doi: 10.1016/s0377-2217(02)00399-5.

Alan Osorio-Mora, Francisco Saldanha-da Gama, and Paolo Toth. A risk-averse latency location-routing problem with stochastic travel times. *European Journal of Operational Research*, 321(3):837–850, 2025. doi: 10.1016/j.ejor.2024.10.041.

Jorge Oyola, Halvard Arntzen, and David L. Woodruff. The stochastic vehicle routing problem, a literature review, part I: models. *EURO Journal on Transportation and Logistics*, 7(3):193–221, 2018. doi: 10.1007/s13676-016-0100-5.

Mojtaba Rajabi-Bahaabadi, Afshin Shariat-Mohaymany, Mohsen Babaei, and Daniele Vigo. Reliable vehicle routing problem in stochastic networks with correlated travel times. *Operational Research*, 21(1):299–330, 2019. doi: 10.1007/s12351-019-00452-w.

Ted K. Ralphs, Matthew J. Saltzman, and Margaret M. Wiecek. An improved algorithm for solving biobjective integer programs. *Annals of Operations Research*, 147(1):43–70, 2006. doi: 10.1007/s10479-006-0058-z.

R. Tyrrell Rockafellar and Stanislav Uryasev. Optimization of conditional value-at-risk. *The Journal of Risk*, 2(3):21–41, 2000. doi: 10.21314/jor.2000.038.

R.Tyrrell Rockafellar and Stanislav Uryasev. Conditional value-at-risk for general loss distributions. *Journal of Banking & Finance*, 26(7):1443–1471, 2002. doi: 10.1016/s0378-4266(02)00271-6.

Borzou Rostami, Guy Desaulniers, Fausto Errico, and Andrea Lodi. Branch-Price-and-Cut Algorithms for the Vehicle Routing Problem with Stochastic and Correlated Travel Times. *Operations Research*, 69(2):436–455, 2021. doi: 10.1287/opre.2020.2037.

Pranab Kumar Sen. A Note on Order Statistics for Heterogeneous Distributions. *The Annals of Mathematical Statistics*, 41(6):2137–2139, 1970. doi: 10.1214/aoms/1177696715.

Suvrajeet Sen, Rekha Pillai, Shirish Joshi, and Ajay K. Rathi. A Mean-Variance Model for Route Guidance in Advanced Traveler Information Systems. *Transportation Science*, 35(1):37–49, 2001. doi: 10.1287/trsc.35.1.37.10141.

Karmel S. Shehadeh. Reducing disparities in transportation distance in a stochastic facility location problem. *Transportation Research Part C: Emerging Technologies*, 153:104199, 2023. doi: 10.1016/j.trc.2023.104199.

Sean R. Sinclair, Gauri Jain, Siddhartha Banerjee, and Christina Lee Yu. Sequential Fair Allocation: Achieving the Optimal Envy-Efficiency Trade-off Curve. *Operations Research*, 71(5):1689–1705, 2023. doi: 10.1287/opre.2022.2397.

Daman Deep Singh, Syamantak Das, and Abhijnan Chakraborty. FairAssign: Stochastically Fair Driver Assignment in Gig Delivery Platforms. In *Proceedings of the 2023 ACM Conference on Fairness, Accountability, and Transparency (FAccT)*, pages 753–763. ACM, 2023. doi: 10.1145/3593013.3594040.

Abhay Sobhanan, Hadi Charkhgard, and Iman Dayarian. Equity-Driven Workload Allocation for Crowdsourced Last-Mile Delivery. *Production and Operations Management*, 35(8):3180–3203, 2026. doi: 10.1177/10591478261425875.

Man Yiu Tsang and Karmel S. Shehadeh. A Unified Framework for Analyzing and Optimizing a Class of Convex Fairness Measures. *Operations Research*, 74(2):1087–1103, 2026. doi: 10.1287/opre.2023.0301.

Eduardo Uchoa, Diego Pecin, Artur Pessoa, Marcus Poggi, Thibaut Vidal, and Anand Subramanian. New benchmark instances for the Capacitated Vehicle Routing Problem. *European Journal of Operational Research*, 257(3):845–858, 2017. doi: 10.1016/j.ejor.2016.08.012.

Bart van Rossum, Rui Chen, and Andrea Lodi. A New Branching Rule for Range Minimization Problems. In *Integer Programming and Combinatorial Optimization (IPCO 2024)*, Lecture Notes in Computer Science, pages 433–445. Springer Nature Switzerland, 2024. doi: 10.1007/978-3-031-59835-7_32.

Bart van Rossum, Rui Chen, and Andrea Lodi. Efficient branching rules for optimizing range and order-based objective functions. *Mathematical Programming*, 216(1-2):539–572, 2026a. doi: 10.1007/s10107-025-02306-0. Online 2025; volume 216 (2026).

Bart van Rossum, Rui Chen, and Andrea Lodi. Enforcing TSP-optimality in fair vehicle routing by cutting planes. *Operations Research Letters*, 70:107539, 2026b. doi: 10.1016/j.orl.2026.107539. Published online 2026; Crossref dates volume 70 as January 2027.

R. J. Vaughan and W. N. Venables. Permanent Expressions for Order Statistic Densities. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 34(2):308–310, 1972. doi: 10.1111/j.2517-6161.1972.tb00909.x.

Bram Verweij, Shabbir Ahmed, Anton J. Kleywegt, George Nemhauser, and Alexander Shapiro. The Sample Average Approximation Method Applied to Stochastic Routing Problems: A Computational Study. *Computational Optimization and Applications*, 24(2-3):289–333, 2003. doi: 10.1023/a:1021814225969.

Shiping Wu, Chun Jin, and Hongguang Bo. Exact solution of workload consistent vehicle routing problem with priority distribution and demand uncertainty. *Computers & Industrial Engineering*, 202:110940, 2025. doi: 10.1016/j.cie.2025.110940.

R.R. Yager. On ordered weighted averaging aggregation operators in multicriteria decisionmaking. *IEEE Transactions on Systems, Man, and Cybernetics*, 18(1):183–190, 1988. doi: 10.1109/21.87068.

Qing Ye, Grani A. Hanasusanto, and Weijun Xie. Distributionally Fair Stochastic Optimization using Wasserstein Distance. arXiv preprint, 2024.

Xiao-Pu Yu, Yong-Shi Hu, and Peng Wu. The consistent vehicle routing problem considering driver equity and flexible route consistency. *Computers & Industrial Engineering*, 187:109803, 2024. doi: 10.1016/j.cie.2023.109803.

Dali Zhang, Dong Li, Hailin Sun, and Liwen Hou. A vehicle routing problem with distribution uncertainty in deadlines. *European Journal of Operational Research*, 292(1):311–326, 2021. doi: 10.1016/j.ejor.2020.10.026.

Junlong Zhang, William H. K. Lam, and Bi Yu Chen. A Stochastic Vehicle Routing Problem with Travel Time Uncertainty: Trade-Off Between Cost and Customer Service. *Networks and Spatial Economics*, 13(4):471–496, 2013. doi: 10.1007/s11067-013-9190-x.