SORTIE: Fair Vehicle Routing When Route Durations Are Random.

Ex-post ordered fairness, rank prices and risk-consistent tours

Quang-Vinh Dang
British University Vietnam, Hung Yen, Vietnam
vinh.dq4@buv.edu.vn ORCID 0000-0002-3877-8024

Working paper, November 2026

Abstract

Fair vehicle routing balances drivers' workloads with ordered measures of route lengths, such as the range or the Gini mean difference. The recent exact approaches, which also require every route to be a shortest tour of its customers, assume deterministic travel times. Drivers, however, experience realised durations. We study fair routing when arc travel times are random, with a fleet-wide day factor and arc-level noise. We distinguish ex-ante fairness (of expected workloads) from ex-post fairness (the expected unfairness of the realised day, sorted day by day) and show that every deterministic fair plan overstates ex-post fairness, with an explicit bound in terms of the routes' quadratic lengths. We prove that multiplicative common shocks leave expected ordered fairness unchanged. We then give an exact, sampling-free formula for the expected value of any ordered measure of independent route durations, a Poisson–binomial order-statistic integral that also handles the negative weights of dispersion measures, and derive from it *rank prices*: the derivative of expected unfairness with respect to a shift of each route's duration. Negative rank prices identify exactly the routes a fairness objective would like to lengthen. Under uncertainty even the min-max measure can prefer a tour that is not the shortest, so route consistency must be imposed explicitly; we define it through each driver's own mean–standard deviation and prove that such tours are supported Pareto tours, computable by parametric Held–Karp. The resulting bilevel heuristic, SORTIE, screens all relocate and swap moves with the rank prices. On instances derived from the CVRPLIB benchmark used by the deterministic literature, deterministic fair plans understate their realised unfairness by 8–27%, while the integral agrees with Monte Carlo to about 0.1% at under a millisecond per plan. Optimising ex-post fairness improves realised fairness by about 1% when noise is proportional to length and by about 1.5% under congestion-driven heterogeneous noise; screening with rank prices gives plans 6–9% better than random screening; and without the lower level, nearly all range-optimised plans buy fairness with detours.

Keywords: vehicle routing; fairness; stochastic travel times; ordered weighted averaging; order statistics; bilevel optimisation.

1 Introduction

Routing plans that minimise total distance often give some drivers much longer days than others. Unequal workloads hurt driver satisfaction and retention and can be seen as unfair [Matl et al., 2018]. The fairness literature in routing therefore optimises a *fairness measure* of the vector of route workloads: the longest route, the range, the Gini mean difference (GMD), the variance or a

mean absolute deviation [Matl et al., 2018, Karsu and Morton, 2015, Bektaş and Letchford, 2020]. Ordered measures, which weight the sorted workloads [Yager, 1988, Ogryczak and Śliwiński, 2003, Nickel and Puerto, 2005], cover all of these in one framework [Blanco et al., 2026].

Non-monotone measures such as the range or the GMD create a well-known pathology: an optimiser can lower the measure by making a light route *longer*, giving a driver a needless detour [Matl et al., 2018]. van Rossum et al. [2026a,b] and Ljubić et al. [2026] close this loophole exactly, by requiring every route to be a shortest (TSP-optimal) tour of its customers. Ljubić et al. [2026] write the problem as a bilevel programme in which a planner assigns customers and each driver solves a TSP, and solve instances with up to 35 customers to optimality for seven measures.

All of these works assume deterministic travel times. A driver's day, however, is a realised duration: congestion, weather, parking and service-time variability differ between routes and between days. Fairness is then a statement about random variables, and three questions arise that the deterministic theory does not ask.

1. *What should be balanced?* The expected workloads (ex-ante fairness), or the workloads drivers actually experience, sorted day by day (ex-post fairness)? The identity of the worst-off driver changes from day to day, and the two notions can disagree.

2. *How can ex-post fairness be optimised?* Sample-average approximation (SAA) is the default, but sorting per scenario makes the objective noisy and its gradient uninformative.

3. *What is a consistent route?* Under random durations "the shortest tour" is not obviously what a driver would choose, and we show that monotone measures no longer protect drivers from detours.

Contributions.

- **Model.** We formulate fair routing with random travel times (a fleet-wide day factor times arc-level noise) as a bilevel problem with an ex-post ordered fairness objective, an expected-cost budget and per-driver risk-consistent tours. With no noise it reduces to the X-CVRP-TSP of Ljubić et al. [2026].

- **Fairness illusion (Proposition 5).** For every ordered measure with nondecreasing weights, ex-ante fairness underestimates ex-post unfairness, by at most $\|\lambda\|_1 v(\sum_k W_k)^{1/2}$, where W_k is the route's quadratic length. Common multiplicative shocks leave the expected value of every ordered measure unchanged (Proposition 2); they do inflate the variance measure and the tail of unfairness.

- **PBOI (Theorem 3).** An exact, sampling-free formula for the expected ordered measure of independent route durations with given marginals (we use moment-matched gamma marginals), as an integral of a Poisson–binomial distribution against tail sums of the weights. It handles negative weights and costs $O(Gm^2)$.

- **Rank prices (Proposition 4).** The derivative of expected unfairness with respect to a location shift of a route's duration equals the expected weight of its rank. Prices of dispersion measures sum to zero, and a negative price is precisely the stochastic incentive to lengthen a route.

- **Risk-consistent tours (Proposition 6, Theorem 8).** Even $\mathbb{E}\max$ can prefer a non-shortest tour. We define consistency through the driver's own mean–standard deviation and

prove that the consistent tour is a supported Pareto tour of (length, quadratic length), found exactly by dichotomic parametric Held–Karp.

- **Algorithm and evidence.** SORTIE, a bilevel local search whose moves are screened by rank prices, is compared with the deterministic fair plan, SAA and ablations on the instances used by the deterministic literature, with exact enumeration on small instances.

2 Literature review

Workload equity in deterministic routing. Matl et al. [2018] survey equity measures for routing workloads and show that their monotonicity governs whether optimal plans can contain needlessly inefficient routes; Matl et al. [2019a] study how the choice of workload resource changes balanced plans, and Matl et al. [2019b] approximate cost–balance Pareto fronts. Min-max routing has a long exact tradition [Applegate et al., 2002, Bertazzi et al., 2015], and bi-objective route balancing is solved exactly by Halvorsen-Weare and Savelsbergh [2016]; ℓ_p norms as fairness measures appear in Bektaş and Letchford [2020]. Recent work on non-monotone and ordered objectives includes branching rules for range and order-based objectives [van Rossum et al., 2024, 2026a], cutting planes that enforce TSP-optimality for the range [van Rossum et al., 2026b], route-cost equity constraints [Agius et al., 2026], Gini-bounded mTSP variants [Bhadoriya et al., 2026], fair cooperative fleets [López Sánchez et al., 2024], driver satisfaction [Bruinink et al., 2025], multi-period equity [Nekooghadirli et al., 2026] and workload/pay design for heterogeneous or crowdsourced drivers [Mahmoodian et al., 2025, Sobhanan et al., 2026]. The ordered machinery comes from location theory [Nickel and Puerto, 2005, Ogryczak, 2000, Blanco et al., 2024, Ljubić et al., 2024, Blanco et al., 2026], the ordered median TSP [Ljubić et al., 2026], OWA [Yager, 1988, Ogryczak and Śliwiński, 2003] and top-k sums [Ogryczak and Tamir, 2003]. Closest to us, Ljubić et al. [2026] propose a unified ordered framework with per-vehicle TSP-optimality through bilevel optimisation. All of these works are deterministic.

Fairness under uncertainty. In routing, Kinable et al. [2026] study a stochastic VRP with partial deliveries where fairness is among *customers* (expected fill rates) under uncertain demand, and Gao et al. [2026] combine sequential fair allocation and routing with ex-post max-min objectives across locations. Consistency in the sense of the consistent VRP (same driver, similar arrival times) is treated under stochastic customers and demands by Alvarez et al. [2024] and with robust demand by Wu et al. [2025], following Groër et al. [2009], Kovacs et al. [2015], Yu et al. [2024]; this notion differs from the TSP-optimality consistency studied here. For drivers, Anh [2022] bound workload differences in task assignment with a distributionally robust chance constraint and Lin et al. [2022] add robust workload fairness to two-echelon meal-delivery routing; gig and rideshare work considers fairness of assignment probabilities and matching under stochastic arrivals [Singh et al., 2023, Basik et al., 2021, Nanda et al., 2020, Hosseini and Schierreich, 2025]. The ex-ante/ex-post distinction is established in welfare economics [Fleurbaey and Peragine, 2012], in inequity-averse stochastic optimisation [Gutjahr, 2021], in Gini-based stochastic shelter location and humanitarian logistics [Mostajabdaveh et al., 2019, Alem et al., 2022, Mostajabdaveh et al., 2025] and in sequential allocation [Sinclair et al., 2023]; see also Shehadeh [2023], Tsang and Shehadeh [2026], Ye et al. [2024], Barak and Talgam-Cohen [2026]. To our knowledge no paper optimises an ordered measure of *random route durations*, separates its ex-ante and ex-post versions, or defines route consistency for random travel times.

Stochastic travel times. A-priori routing with random travel times goes back to Laporte et al. [1992] and Kenyon and Morton [2003], the latter with makespan-type objectives; see Gómez et al. [2016], Adulyasak and Jaillet [2016], Errico et al. [2016, 2018], Zhang et al. [2013], Ehmke et al. [2015], Osorio-Mora et al. [2025] and the surveys Oyola et al. [2018], Gendreau et al. [2016]. Correlated travel times are handled by Rostami et al. [2021], Bakach et al. [2021], Rajabi-Bahaabadi et al. [2019], Bomboi et al. [2021], robust and distributionally robust variants by Han et al. [2014], Jaillet et al. [2016], Zhang et al. [2021], Ghosal and Wiesemann [2020], and SAA by Verweij et al. [2003]. Mean–risk path choice [Nikolova and Stier-Moses, 2014, Sen et al., 2001] motivates our driver-level consistency criterion. CVaR [Rockafellar and Uryasev, 2000, 2002] and the price of fairness [Bertsimas et al., 2011, Mo and Walrand, 2000] complete the background.

Order statistics and parametric optimisation. Order statistics of independent, non-identically distributed variables have permanent representations [Vaughan and Venables, 1972, Bapat and Beg, 1989, Balakrishnan, 2007, Glueck et al., 2008] and are related to the Poisson–binomial count of successes in independent trials [Hoeffding, 1956, Fernandez and Williams, 2010, Hong, 2013, Biscarri et al., 2018]; see David and Nagaraja [2003], Sen [1970]. Supported efficient solutions of bi-criteria problems are found by dichotomic search [Aneja and Nair, 1979, Ehrgott, 2005, Ehrgott and Gandibleux, 2000, Ralphs et al., 2006], and the TSP on few nodes by the Held–Karp recursion [Held and Karp, 1962, Bellman, 1962].

3 Model

Let $G = (V, A)$ be a complete graph with depot 0, customers $V' = \{1, \dots, n\}$, symmetric costs c_{ij} satisfying the triangle inequality, demands q_i and m identical vehicles of capacity Q. A plan consists of a partition $S_1, \dots, S_m$ of V' into nonempty capacity-feasible sets and a tour r_k of $\{0\} \cup S_k$ for each vehicle.

Random workloads. The travel time of arc (i, j) is $\tau_{ij} = Y c_{ij} \varepsilon_{ij}$, where $Y = \exp(s_Y Z - s_Y^2/2)$, $Z \sim N(0, 1)$, is a fleet-wide day factor with $\mathbb{E}Y = 1$, and the arc disturbances ε_{ij} are independent of each other and of Y with mean 1 and variance v^2 (gamma in our experiments). The workload of driver k is $\Phi_k = Y X_k$ with

$$X_k = \sum_{(i,j)\in r_k} c_{ij}\varepsilon_{ij}, \qquad \mathbb{E}X_k = L_k = \sum_{(i,j)\in r_k} c_{ij}, \qquad \operatorname{Var} X_k = \sigma_k^2 = v^2 W_k, \quad W_k = \sum_{(i,j)\in r_k} c_{ij}^2.$$

L_k is the tour length and W_k its *quadratic length*. Given Y the workloads are independent; unconditionally they are positively correlated.

Ordered fairness. For $a \in \mathbb{R}^m$ with order statistics $a_{(1)} \le \cdots \le a_{(m)}$ and weights $\lambda \in \mathbb{R}^m$, $\ell_\lambda(a) = \sum_j \lambda_j a_{(j)}$. We use the maximum ($\lambda = e_m$), the range ($e_m - e_1$), the GMD ($\lambda_j = 2(2j - m - 1)/m^2$) and the mean absolute deviation from the median (MAD; $\lambda = (-1, \dots, -1, 0, 1, \dots, 1)/m$ for odd m), together with the empirical variance $\operatorname{Var}_{\text{emp}}(a) = m^{-1} \sum_k (a_k - \bar{a})^2$. All four λ are nondecreasing.

Definition 1 (Three notions of fairness)**.** For a plan with random workloads Φ: *ex-ante* unfairness $\ell_\lambda(\mathbb{E}\Phi)$; *ex-post* unfairness $\mathbb{E}\ell_\lambda(\Phi)$; *risk of unfairness* $\operatorname{CVaR}_\beta \ell_\lambda(\Phi)$.

A deterministic fair model optimises the ex-ante notion. Drivers experience the ex-post one. SORTIE optimises ex-post unfairness and reports the CVaR out of sample.

4 Theory

Proposition 2 (Common shocks)**.** *For every λ, $\mathbb{E}\ell_\lambda(\Phi) = \mathbb{E}\ell_\lambda(X)$, and $\mathbb{E}\,\mathrm{Var}_{\mathrm{emp}}(\Phi) = e^{s_Y^2}\,\mathbb{E}\,\mathrm{Var}_{\mathrm{emp}}(X)$.*

Proof. Sorting commutes with multiplication by $y > 0$, so $\ell_\lambda(yX) = y\ell_\lambda(X)$; take expectations using independence of Y and X and $\mathbb{E}Y = 1$. The empirical variance is homogeneous of degree two and $\mathbb{E}Y^2 = e^{s_Y^2}$. □

Fleet-wide shocks therefore drop out of the planning objective for ordered measures, which reduces evaluation to the conditionally independent vector X.

Theorem 3 (Poisson–binomial order-statistic integral)**.** *Let $X_1, \dots, X_m \geq 0$ be independent with CDFs F_k. For $t \geq 0$ let $N(t) = \#\{k : X_k \leq t\}$, a Poisson–binomial count with success probabilities $F_k(t)$, and $\pi_r(t) = \mathbb{P}(N(t) = r)$. With tail sums $\Lambda_r = \sum_{j=r+1}^{m} \lambda_j$,*

$$\mathbb{E}\ell_\lambda(X) = \int_0^\infty \sum_{r=0}^{m-1} \Lambda_r\, \pi_r(t)\, dt. \tag{1}$$

Proof. $X_{(j)} > t$ if and only if fewer than j of the X_k are at most t, so $\mathbb{E}X_{(j)} = \int_0^\infty \mathbb{P}(N(t) \leq j-1)\, dt$. Hence $\mathbb{E}\ell_\lambda(X) = \int_0^\infty \sum_j \lambda_j \sum_{r \leq j-1} \pi_r(t)\, dt = \int_0^\infty \sum_r \pi_r(t) \sum_{j \geq r+1} \lambda_j\, dt$, where Tonelli's theorem is applied to the positive and negative parts of λ separately. □

The vector $\pi(t)$ is the coefficient vector of $\prod_k (1 - F_k(t) + F_k(t)z)$, obtained by m convolution steps. Below the smallest lower quantile the integrand equals Λ_0, above the largest upper quantile it vanishes ($\Lambda_m := 0$), and in between we use Gauss–Legendre panels whose breakpoints are quantiles of *every* marginal, so that a short, low-variance route is resolved even when the other routes are long. We use moment-matched gamma marginals $X_k \sim \Gamma(L_k^2/\sigma_k^2, \sigma_k^2/L_k)$. The identity behind (1) is classical for order statistics of non-identical variables [Bapat and Beg, 1989, Balakrishnan, 2007]; our contribution is to make it an exact planning objective for ordered fairness, where the tail sums Λ_r absorb the negative weights of dispersion measures, and to differentiate it.

Proposition 4 (Rank prices)**.** *Let R_k be the rank of X_k and define $\varpi_k = \sum_j \lambda_j \mathbb{P}(R_k = j)$, where $\mathbb{P}(R_k = j) = \int f_k(t)\, \mathbb{P}(N_{-k}(t) = j-1)\, dt$ and N_{-k} counts the other drivers. Then $\frac{d}{d\delta}\mathbb{E}\ell_\lambda(X + \delta e_k)\big|_{\delta=0} = \varpi_k$ and $\sum_k \varpi_k = \sum_j \lambda_j$.*

Proof. ℓ_λ is piecewise linear with $\partial\ell_\lambda/\partial a_k = \lambda_{R_k(a)}$ off the null set of ties and is $\|\lambda\|_1$-Lipschitz, so differentiation and expectation commute by dominated convergence. The ranks form a permutation, so $\sum_k \mathbb{P}(R_k = j) = 1$. □

For the range, GMD and MAD the prices sum to zero: burdened drivers carry positive prices and light drivers negative ones. A negative price is exactly the incentive to lengthen a route. Deterministically, the price of the lightest route under the range is -1, and a detour of length δ reduces the range by δ; stochastically, the incentive is spread over all drivers that might be the lightest on a given day. SORTIE uses the prices, together with $\psi_k = \partial\mathbb{E}\ell_\lambda/\partial\sigma_k$, to predict the effect of a move.

Proposition 5 (Fairness illusion)**.** *If λ is nondecreasing then ℓ_λ is convex and*

$$0 \leq \mathbb{E}\ell_\lambda(X) - \ell_\lambda(L) \leq \|\lambda\|_1\, \mathbb{E}\|X - L\|_\infty \leq \|\lambda\|_1\, v\Big(\sum_k W_k\Big)^{1/2}.$$

Moreover $\mathbb{E}\,\mathrm{Var}_{\mathrm{emp}}(X) = \mathrm{Var}_{\mathrm{emp}}(L) + \frac{m-1}{m^2}\sum_k \sigma_k^2$, and for Gaussian marginals $\mathbb{E}\,\mathrm{GMD}(X) = m^{-2}\sum_{i \neq j}\left[s_{ij}\sqrt{2/\pi}\, e^{-d_{ij}^2/(2s_{ij}^2)} + d_{ij}(1 - 2\Phi(-d_{ij}/s_{ij}))\right]$ with $d_{ij} = L_i - L_j$ and $s_{ij}^2 = \sigma_i^2 + \sigma_j^2$.

Proof. For nondecreasing λ the rearrangement inequality gives $\ell_\lambda(a) = \max_\sigma \sum_j \lambda_j a_{\sigma(j)}$, a maximum of linear functions; Jensen's inequality gives the lower bound. Since $|a_{(j)} - b_{(j)}| \le \|a - b\|_\infty$ for every j, ℓ_λ is $\|\lambda\|_1$-Lipschitz in the sup norm, and $\mathbb{E}\max_k |Z_k| \le (\mathbb{E}\sum_k Z_k^2)^{1/2}$. The variance identity follows from $\mathbb{E}X_k^2 = L_k^2 + \sigma_k^2$ and $\mathbb{E}\bar{X}^2 = \bar{L}^2 + m^{-2}\sum_k \sigma_k^2$; the GMD formula is the mean of a folded normal. □

A deterministic fair plan therefore *overstates* the fairness drivers will experience. The overstatement grows with the arc noise and with the quadratic lengths: long arcs, not long routes, are what make a balanced plan unbalanced in practice.

Proposition 6 (Monotonicity no longer protects drivers)**.** *With deterministic workloads, every monotone measure ($\lambda \ge 0$) has an optimal plan made of TSP-optimal routes [Matl et al., 2018]. With random workloads this fails even for* $\mathbb{E}\max$.

Example 7. Two drivers; $X_1 \equiv 100$. Driver 2 can drive tour A ($L = 100$, $\sigma = 10$) or tour B ($L = 101$, $\sigma \approx 0$). With Gaussian X_2, $\mathbb{E}\max = 100 + 10\varphi(0) \approx 103.99$ under A and ≈ 101 under B, so the optimiser picks the longer tour B.

Route consistency must therefore be imposed for every measure, and it needs a risk attitude. We let each driver rank the tours of their customer set by

$$\rho_\kappa(r) = \frac{\mathbb{E}[\Phi(r) \mid Y] + \kappa\,\mathrm{SD}[\Phi(r) \mid Y]}{Y} = L(r) + \kappa v\sqrt{W(r)},$$

which ranks tours identically on every day. $\kappa = 0$ is TSP-optimality (ties broken by smaller W).

Theorem 8 (Risk-consistent tours are supported)**.** *For $\kappa \ge 0$, some minimiser of ρ_κ over the tours of a fixed customer set is a supported extreme tour: it minimises $L(r) + \theta W(r)$ for some $\theta \in [0, \infty)$, or minimises W lexicographically after L ($\theta = \infty$). All supported extreme tours are enumerated by dichotomic search, each step being one Held–Karp run with arc weights $c_{ij} + \theta c_{ij}^2$.*

Proof. $f(L, W) = L + \kappa v\sqrt{W}$ is concave and nondecreasing on $\mathbb{R}^2_+$. A concave function bounded below on a pointed polyhedron attains its minimum at an extreme point; since f is nondecreasing, the minimum of f over the finite set $P = \{(L(r), W(r))\}$ equals its minimum over $\mathrm{conv}(P) + \mathbb{R}^2_+$ and is attained at an extreme point of that set. These extreme points are exactly the unique minimisers of $w_1 L + w_2 W$ with $(w_1, w_2) \ge 0$, $(w_1, w_2) \ne 0$ (for $w_1 = 0$ with ties broken by smaller L), and dichotomic search enumerates all of them [Aneja and Nair, 1979]. Tours in the relative interior of a hull edge are dominated by the edge's end points and need not be enumerated. With heterogeneous noise (ε_{ij} with variance $v^2 H_{ij}^2$) the same holds with $W = \sum (H_{ij} c_{ij})^2$ and arc weights $c_{ij} + \theta H_{ij}^2 c_{ij}^2$. □

The SORTIE problem.

$$\begin{aligned} \min\ & \mathbb{E}\,\ell_\lambda\big(X(r_1, \dots, r_m)\big) \\ \text{s.t.}\ & (S_1, \dots, S_m) \text{ capacity-feasible partition},\ S_k \ne \emptyset, \qquad \textstyle\sum_k L(r_k) \le B \cdot C^*, \\ & r_k \in \arg\min\{\rho_\kappa(r) : r \text{ a tour of } \{0\} \cup S_k\} \quad \forall k, \end{aligned}$$

where C^* is the minimum expected total length and $B \ge 1$ the budget factor. With $v = 0$ and $\kappa = 0$ this is the X-CVRP-TSP of Ljubić et al. [2026].

Algorithm 1 SORTIE

$P \leftarrow$ cost-efficient plan; $P^* \leftarrow P$
while time remains **do**
    compute rank prices ϖ_k and dispersion prices ψ_k of P (Prop. 4)
    score every relocate and swap move by $\sum_{k\in\{a,b\}} \varpi_k \Delta L_k + \psi_k \Delta\sigma_k$ (insertion deltas)
    evaluate the K best-scored budget- and capacity-feasible moves with screening tours and PBOI (Thm. 3)
    confirm the improving candidates in order with the exact lower level (Thm. 8)
    if a confirmed move improves P **then** apply it
    else ruin (price-weighted or related removal, at most 12 customers) and recreate (price-scored regret-2); repair cost if over budget; confirm every changed route; accept by simulated annealing
    end if
    update P^*
end while
return P^*

5 The SORTIE algorithm

The lower level is exact for routes of up to 15 customers (numba Held–Karp, memoised by customer set, about 5 ms at 15 customers) and uses 2-opt/Or-opt from several starts beyond that, including a warm start built from the route's previous tour by cheapest insertion. Solving it for every candidate move would dominate the run time, so candidates are evaluated with *screening tours*: a single warm-started 2-opt/Or-opt descent (for $\kappa > 0$, inside the same dichotomic search). A move is applied only after both changed routes have been re-solved by the exact lower level and the move still improves the objective and respects the budget; perturbed plans are confirmed route by route in the same way. Every plan the search accepts therefore satisfies the bilevel constraint; the screening tours only decide which moves are examined. Because a screening tour is never shorter than the exact one, a move that improves only with exact tours can be missed; this costs search quality, not feasibility. The evaluator costs about 0.4 ms per plan for $m = 5$. Together with a cap of 12 customers on the ruin size, this raises the iteration rate of SORTIE-0 at $n = 100$ from about 250 to about 1700 per minute, and at $n = 75$ from about 170 to about 1800.

6 Computational study

6.1 Set-up

Instances. We follow the slicing rule of the deterministic fair routing literature [Ljubić et al., 2026]. Block b of size n consists of the depot and the n consecutive customers $bn+1, \ldots, (b+1)n$ of CVRPLIB X-n641-k35 [Uchoa et al., 2017], with ceiled Euclidean distances, $m = 5$ vehicles and capacity $Q = \lfloor \sum_i q_i/4 \rfloor$, so one vehicle's worth of slack is available. The main grid uses $n \in \{15, 20, 25, 30, 35\}$ and blocks $b = 0, \ldots, 4$ (25 instances); the scale test uses $n \in \{50, 75, 100\}$ and blocks 0–5; the exactness test uses ten instances with $n = 10$, $m = 3$ and $Q = \lfloor \sum_i q_i/2.4 \rfloor$.

Uncertainty. Unless stated otherwise $v = 0.3$ (gamma arc noise) and $s_Y = 0.15$. In the congestion experiment (E8) the arc coefficient of variation is vH_{ij}, where $H_{ij} \in [1/3, 2]$ is largest for arcs near a randomly drawn congestion centre, so noise is no longer proportional to length.

Table 1: E1: PBOI against arc-level Monte Carlo (50 000 samples) on 168 plans; z: difference over its Monte Carlo standard error.

Measure	Plans	Mean |err| (%)	Max |err| (%)	Share $\|z\| < 3$
Max	168	0.041	0.137	0.994
Range	168	0.062	0.249	1.000
GMD	168	0.057	0.241	1.000
MAD	168	0.064	0.272	0.988
Var	168	0.108	0.440	0.994

Budget. C^* is the cost of the best of three restarts of the cost-minimising search; the total length of every fair plan must satisfy $\sum_k L_k \le B\, C^*$ with $B \in \{1.05, 1.10\}$.

Methods. All methods share the search of Algorithm 1, the start plan, the time limit (15 s for $n \le 25$, 25 s for $n \ge 30$, 120 s for $n \ge 50$) and the random seed. They differ only in the objective and in the lower level. *Det* minimises the deterministic ordered measure $\ell_\lambda(L)$ of the tour lengths, i.e. the model of Ljubić et al. [2026] solved heuristically. *SAA* minimises the sample average of $\ell_\lambda(X)$ over 200 fixed arc scenarios. *SORTIE-0* minimises the exact ex-post value by PBOI with TSP-optimal tours ($\kappa = 0$); *SORTIE-1* uses the risk-consistent lower level with $\kappa = 1$. In Det and SAA the moves are screened by the deterministic and the sample objective respectively.

Evaluation. Every plan is evaluated out of sample by arc-level Monte Carlo with 20 000 scenarios (10 000 for $n \ge 50$), including the day factor, with common random numbers across methods. The reported *ex-post value* is the Monte Carlo estimate of $\mathbb{E}\ell_\lambda(\Phi)$. Paired comparisons use the Wilcoxon signed-rank test. All tables are generated from the raw results by the released scripts.

6.2 Accuracy and cost of the evaluator (E1, E2)

Table 1 compares PBOI with 50 000-sample arc-level Monte Carlo on 168 plans: the cost-efficient plans and random feasible plans whose routes are sequenced by the lower level, for $n \in \{15, 25, 35, 50\}$. The mean absolute relative error is below 0.11% for all five measures and at most 0.44% in the worst case, and 99–100% of the differences lie within three Monte Carlo standard errors. The mean z-scores are not exactly zero (up to 0.34 for the MAD): PBOI carries a systematic bias of a few hundredths of a percent, which we attribute to the moment-matched gamma marginal of each route (a sum of arc gammas with different scales is not exactly gamma). PBOI takes 0.44 ms per plan for $m = 5$, against 0.28 s for 20 000-scenario Monte Carlo. A 200-scenario SAA is about seven times faster than PBOI but is a noisy estimate. The advantage of PBOI is therefore that it is exact and noise-free at a cost compatible with thousands of evaluations per second, not raw speed.

On the ten $n = 10$ instances we enumerated all capacity- and budget-feasible partitions (4 to 145 per instance) with exact Held–Karp tours. The cost search found the optimal C^* on all ten. For both range and GMD, SORTIE-0 found the plan with the smallest exact ex-post value on nine of ten instances (mean gap 0.39%, maximum 3.9%). Det found the deterministic optimum on nine of ten; on the tenth instance, which has only 20 budget-feasible partitions, it stopped at a plan with twice the optimal deterministic range.

Table 2: E3: what each method believes its unfairness is, relative to the true ex-post value (Monte Carlo without the day factor, which is what each model represents), mean signed error over 50 runs per cell. Last column: largest absolute error of the PBOI prediction for any method's plan.

	Believed vs true (mean signed %)				PBOI
Measure	Det	SAA	SORTIE-0	SORTIE-1	max |err| %
Max	-7.97	-0.71	-0.01	-0.01	0.20
Range	-17.56	-0.99	+0.03	+0.02	0.70
GMD	-14.68	-0.91	+0.04	+0.05	0.61
MAD	-16.17	-0.88	+0.05	+0.07	0.68
Var	-27.23	-1.33	+0.00	+0.02	0.98




Figure 1: E3, $B = 1.10$: per-instance improvement of the ex-post value over Det.

6.3 The fairness illusion and the value of ex-post optimisation (E3)

Table 2 quantifies Proposition 5. A deterministic fair plan believes its unfairness is the ex-ante value, which understates the true ex-post value by 8% for the maximum, 15–18% for the range, GMD and MAD, and 27% for the variance. The in-sample SAA value is optimistic by about 1%, which is the optimiser's curse on 200 scenarios. The PBOI value believed by SORTIE has a mean signed error below 0.1%, a mean absolute error of 0.07–0.23% and a worst case of 1%.

Table 3 reports the out-of-sample ex-post value relative to Det. The signed-rank test favours SORTIE-0 and SORTIE-1 over Det in 9 of the 10 cells each. Pooled over all 250 runs, SORTIE-0 improves on Det by 0.89% on average (median 0.23%; better in 141 runs, tied in 96, worse in 13; $p < 10^{-18}$) and SORTIE-1 by 0.77% (better in 154, worse in 31). SAA improves by only 0.14% (median zero; better in 115, worse in 74) and is favoured significantly in 3 cells. Head to head, SORTIE-0 beats SAA in 145 runs and loses in 15, and SORTIE-1 beats it in 147 and loses in 27 ($p < 10^{-17}$). The largest gains are for the maximum at $B = 1.10$ (2.1%, 24 of 25 instances), where the extra budget lets the planner trade a slightly longer but less variable longest route.

The improvements are small, about one percent on average, and this is itself a finding. With homogeneous noise ($\sigma_k^2 = v^2 W_k$) the standard deviation of a route grows with its length, so the plans that balance lengths also nearly balance risks, and a deterministic fair plan is close to ex-post optimal even though its *assessment* of its own fairness is badly wrong. For context, the cost-optimal plan has an ex-post range of 2087 against about 1340 for the fair plans, at a cost 6.5% higher: fairness itself is worth optimising; the stochastic refinement adds about one percent on top. For range-optimised plans the $\mathrm{CVaR}_{0.9}$ of the realised range, which none of the methods optimises, is 2054 for Det, 2059 for SAA, 2037 for SORTIE-0 and 2032 for SORTIE-1; both SORTIE variants improve it at both budgets, SAA at neither.

Table 3: E3: improvement of the out-of-sample ex-post value over Det (mean %, positive is better; in brackets the number of instances on which the method is strictly better). *: Wilcoxon signed-rank $p < 0.05$ in favour of the method; the test is on ranks and can disagree in sign with the mean when one instance dominates (Var, $B = 1.10$, SORTIE-1).

Measure	B	SAA	SORTIE-0	SORTIE-1
GMD	1.05	-0.40 (6/25)	-0.09 (8/25)	-0.03 (9/25)
GMD	1.10	-0.32 (9/25)	+0.94* (14/25)	+0.34* (17/25)
MAD	1.05	+0.86* (11/25)	+1.10* (13/25)	+1.24* (12/25)
MAD	1.10	+0.70 (13/25)	+0.96* (14/25)	+1.53* (18/25)
Max	1.05	+0.71* (14/25)	+0.61* (15/25)	+0.77* (15/25)
Max	1.10	+1.70* (23/25)	+2.08* (24/25)	+1.93* (24/25)
Range	1.05	+0.12 (13/25)	+0.64* (15/25)	+1.56* (15/25)
Range	1.10	-0.21 (13/25)	+1.07* (17/25)	+0.39* (15/25)
Var	1.05	+0.48 (7/25)	+0.50* (9/25)	+0.48* (13/25)
Var	1.10	-2.25 (6/25)	+1.09* (12/25)	-0.49* (16/25)

An illustrative instance. Figure 2 shows the cost-optimal plan and the range-optimised Det and SORTIE-0 plans of X641-n30-b0 at $B = 1.10$, the instance whose SORTIE-0 gain is closest to the median of the 25 instances (+0.5%). The cost plan leaves one driver with a route of length 148 and another with 2610. Both fair plans balance the lengths to a deterministic range of about 940, and they differ only in how two route pairs are split. Figure 3 shows what the drivers experience. The realised workloads are wide and right-skewed, and the expected realised range is about 1225 for both fair plans: 30% above the deterministic range the Det planner reports. Figure 4 gives the distribution of the realised range of the two plans; in one day out of ten it exceeds 1750. The worst instance for SORTIE-0 in this cell is X641-n35-b0, where Det is 7.5% better (Figure 18 in Appendix A); this is consistent with the search noise documented in E7b rather than a systematic effect, since on the same instance SORTIE-0 is better or tied in eight of the other nine cells.

How tight is the illusion bound? Figure 5 plots the gap between the true ex-post value and the ex-ante value of every Det plan against the bound of Proposition 5. No plan violates the bound. The bound is conservative: the realised gap is on average 33% of the bound for the maximum, 18% for the range and 11–12% for the GMD and MAD, because it charges every route its full standard deviation while only the routes near the extreme ranks matter. Its value is that it is computable from the quadratic lengths alone and grows like $v\sqrt{\sum_k W_k}$, which is the scaling observed in E6.

6.4 Route consistency (E4)

E4 removes the lower level: the planner may sequence each route freely (*free*) instead of using the TSP-optimal tour (*lower*). For the maximum, free sequencing changes almost nothing, as monotonicity predicts in the deterministic case. For the range and the GMD the free planner lengthens the short routes: 23 of 25 Det plans and 24 of 25 SORTIE plans for the range contain a route that is not a shortest tour of its customers; the excess length is 95 (Det) and 113 (SORTIE) distance units on average over the runs that have one. This lowers the ex-post value by 1–3% for Det on both measures and for SORTIE on the range (Table 4); for SORTIE on the GMD it is 0.9% worse, because the larger space slows the search. Wherever it helps, the reduction comes from making some drivers drive detours. This is exactly the behaviour signalled by negative rank prices and excluded by the bilevel constraint. The deterministic belief error persists with free tours




Figure 2: X641-n30-b0, $B = 1.10$: cost-optimal plan and the range-optimised Det and SORTIE-0 plans. L_k: tour lengths.

(-24% for the range); the PBOI belief remains exact.

6.5 Rank-price screening (E5)

Replacing the price screen by $K = 8$ random feasible moves, with everything else unchanged, gives slightly more iterations (3200 against 2700) but worse plans: price screening lowers the ex-post value by 5.6% for the range and 8.5% for the GMD on average (medians 0% and 5.4%; better on 12 and 17 of 25 instances, worse on 3 and 2, ties otherwise; $p = 0.005$ and $p < 10^{-3}$). For the range the mean is driven by a few instances on which random screening stalls far from a good plan; for the GMD the advantage is broad. The rank prices are thus an effective first-order guide for the search, and not only an interpretive device.

6.6 Noise level (E6)

Table 5 varies v for $n = 25$. The illusion grows monotonically with v, as the bound of Proposition 5 suggests: at $v = 0.5$ a deterministic plan understates its range by 43%. The gain of SORTIE over Det also grows with v for the range, from -0.5% at $v = 0.1$ (where the illusion is small and the differences are search noise) to 2.1% at $v = 0.5$ (better on all five instances); for the GMD it is 1.1%, 0.7% and 1.8%.

6.7 Larger instances (E7, E7b)

E7 runs $n \in \{50, 75, 100\}$ on six blocks each, with a 120 s limit, for the range, GMD and maximum (18 instance–measure pairs per size). With the screening lower level the search runs about 9000 iterations per run at $n = 50$ and 3500 at $n = 100$ for SORTIE-0 (Det: 15 000 and 4400). PBOI remains accurate (mean absolute error 0.15%, at most 0.9%) and the deterministic belief error is -17% on average, as on the small instances.




Figure 3: Simulated realised workloads $\Phi_k = YX_k$ of the three plans of Figure 2 (20 000 days); diamonds mark the tour lengths L_k that a deterministic model sees.

Table 6 pools the 18 pairs per size. SORTIE-0 improves on Det by about 2% at every size and is better on 12–13 of 18 pairs; the improvement is significant at $n = 50$ ($p = 0.035$) but not at $n = 75$ or $n = 100$ ($p = 0.17$). SORTIE-1 is significant at $n = 50$ but not beyond: its lower level runs a dichotomic search per route and completes only about 850 iterations at $n = 100$, and 56% of its runs there (39% at $n = 75$) still improved by more than 0.5% in the second half of the time limit. Single runs at $n \geq 75$ are thus dominated by search noise, especially for the range and the GMD, whose value depends on a few routes.

E7b separates the method from the search noise at $n = 100$: the six instances and three measures are solved by Det, SAA and SORTIE-0 with three solver seeds each and a 240 s limit, and each method is represented by the mean ex-post value over its seeds (Table 7). Averaged over seeds, SORTIE-0 improves on Det by 4.2% (median 2.5%, better on 13 of 18 pairs, $p = 0.021$); SAA improves by 3.2% ($p = 0.10$). The difference between SORTIE-0 and SAA (1.1%, 12 of 18) is not significant ($p = 0.25$), and no single measure is significant on six instances. The seed-to-seed variation is itself informative: for the range, the ex-post value of the Det plan varies by 11% across seeds, that of SORTIE-0 by 3%. The deterministic range is flat over many plans whose realised fairness differs, and the search picks among them arbitrarily; the ex-post objective distinguishes them. Even at 240 s, 19% of the SORTIE-0 runs improved their in-model value by more than 0.5% in the second half of the run, so the absolute values at $n = 100$ are not fully converged.

Figure 8 shows the anytime behaviour behind these numbers, and Figure 9 the seed-to-seed spread at $n = 100$.

6.8 Heterogeneous noise: congestion (E8)

E3 suggests that the stochastic refinement matters little when a route's variance is proportional to its length. E8 breaks that proportionality: arcs near a congestion centre have up to twice the base coefficient of variation and arcs far from it a third. We rerun the main grid ($B = 1.10$; range, GMD




Figure 4: Distribution of the realised range of the Det and SORTIE-0 plans of Figure 2; dashed: deterministic range of the tour lengths, solid: expected realised range.

and max) on the congested versions of the same 25 instances, starting from the same cost plans.

The illusion becomes somewhat larger: a deterministic plan understates its range by 26%, its GMD by 22% and its maximum by 10%, against 23%, 19% and 9% on the same cells under homogeneous noise. The PBOI belief has a mean signed error below 0.2% (mean absolute error at most 0.26%). The gain of optimising ex-post fairness, however, grows only slightly (Table 8). Pooled over 75 runs, SORTIE-0 improves on Det by 1.51% (median 1.24%; better in 53, worse in 3; $p < 10^{-8}$) and SORTIE-1 by 1.66% (better in 57, worse in 10), against 1.36% for SORTIE-0 on the same cells without congestion. By measure the change is 0.94% $\rightarrow$ 1.40% for the GMD, 2.08% $\rightarrow$ 2.15% for the maximum and 1.07% $\rightarrow$ 0.96% for the range. SAA gains 1.28%, but with a median of only 0.47%, and SORTIE-0 beats it head to head in 49 of 75 runs (worse in 6; $p < 10^{-6}$). The $\mathrm{CVaR}_{0.9}$ of the range is 1919 for Det, 1905 for SAA, 1912 for SORTIE-0 and 1907 for SORTIE-1. Heterogeneous noise thus makes a deterministic plan's self-assessment much worse, but in our instances it changes the best plan only a little more than homogeneous noise does: the congested arcs are concentrated in one part of the network, and the assignment that balances lengths still roughly balances risk there.

Figure 10 shows the congestion multiplier on the illustrative instance, and Figure 11 pairs every instance and measure with and without congestion: the points scatter around the diagonal, with the GMD above it and the range on both sides.

6.9 The budget frontier (E9)

E9 repeats the comparison of Det and SORTIE-0 on the 10 instances with $n \in \{25, 35\}$ for budget factors B from 1.02 to 1.30 (Table 9, Figure 12). Two effects stand out. First, the gain of SORTIE-0 grows with the budget: from about 1% for $B \leq 1.10$ to 3.2% at $B = 1.15$ and 6.8% at $B = 1.30$, where SORTIE-0 is better on all 30 instance–measure pairs. With little slack every fair plan is close to the cost-optimal one and there is little room to choose; with more slack there are many plans whose deterministic range is about equal but whose realised range differs, and only the ex-post objective can tell them apart. Second, SORTIE-0 uses *less* of the budget. At $B = 1.30$ the Det




Figure 5: Fairness illusion of the Det plans of E3 (true ex-post minus ex-ante value) against the bound $\|\lambda\|_1 v(\sum_k W_k)^{1/2}$ of Proposition 5.

plans for the maximum have a total length 20.5% above C^* and an expected realised maximum 5.2% below that of the cost plan; the SORTIE-0 plans use 13.5% and achieve 8.7%. For the maximum the deterministic objective even deteriorates from $B = 1.20$ to $B = 1.30$: the extra length it buys goes into routes whose deterministic length is irrelevant to the maximum but whose variance raises the realised one. The price of fairness is therefore lower under the ex-post objective, at every budget level we tested.

The same cells at $B = 1.10$ appear in E3, and the per-cell means differ (for example -0.7% for the range in E9 against $+0.9\%$ in E3 on the same ten instances). The runs are identical except for wall-clock timing: the search stops at a time limit, so the iteration at which it stops, and hence the plan, can differ between two executions on a shared machine. This is another manifestation of the search noise analysed in E7b, and the reason we draw conclusions only from pooled comparisons.

6.10 Fleet size (E10)

E10 varies the number of vehicles on the $n = 35$ instances, with capacity $\lfloor \sum_i q_i/(m-1) \rfloor$ so that one vehicle's worth of slack remains (Table 10, Figure 13). The illusion grows with m: a deterministic plan understates its range by 10% with three vehicles, 24% with five and 27% with seven. More routes compete for the extreme ranks, so more of the realised unfairness comes from which route happens to be longest on the day, which a deterministic model cannot see. The gain of SORTIE-0 grows accordingly, from 0.5% ($m = 3$) and 0.6% ($m = 5$) to 3.2% with seven vehicles (better on 13 of 15 pairs). With $m = 3$ the GMD is proportional to the range (4/9 of it), so the two rows coincide.




Figure 6: E3, range-optimised plans: total length relative to C^* against the expected realised range relative to that of the cost-optimal plan.

6.11 Risk attitude of the drivers (E13)

E13 solves the $n \in \{25, 35\}$ instances with the risk-consistent lower level for $\kappa \in \{0, 0.5, 1, 2, 4\}$ (Table 11, Figure 14). Under homogeneous noise the risk-consistent tour rarely differs from the shortest one: even at $\kappa = 4$ only 14–16% of the routes change, and the extra length is about 13 distance units per plan, under 0.2% of the total. The ex-post value changes by less than the search noise (median ratio to $\kappa = 0$ equal to one, interquartile range within $\pm 1\%$). The reason is that, for a fixed customer set, a tour's length and its quadratic length are nearly comonotone: the shortest tour is also nearly the least variable one. Proposition 6 shows that this need not be so, and with heterogeneous noise the two can separate, but on these instances the choice of κ is a matter of the drivers' preferences rather than of fairness.

6.12 Why the rank prices work (E14)

E14 checks the first-order model behind the screen directly. For the cost-optimal start plan and the final SORTIE-0 plan of the ten $n \in \{25, 35\}$ instances, we evaluated up to 400 relocate and swap moves both by the rank-price prediction and exactly (exact lower level and PBOI); 5577 moves in total (Table 12, Figures 15 and 16). The prediction ranks the moves almost as the exact evaluation does: Spearman's ρ is 0.89–0.94 on the start plans and 0.78–0.89 on the final plans. On the start plans the exact-best improving move is among the 8 best predicted moves in 80% of the cases (random ordering: 7%) and always among the 21 best. On the final plans of SORTIE-0 there is

Table 4: E4: free tours against lower-level tours, $B = 1.10$, 25 instances per row. [a]Runs with at least one route that is not a shortest tour of its customers. [b]Mean number of such routes. Excess: total length above the shortest tours of the same customer sets. Belief err: believed minus true ex-post value. Ex-post: out-of-sample ex-post value.

Measure	Method	Runs[a]	Routes[b]	Excess	Belief err (%)	Ex-post
Max	Det-lower	0/25	0.00	0.0	-8.94	2221.0
Max	Det-free	0/25	0.00	0.0	-9.01	2232.4
Max	SORTIE-lower	0/25	0.00	0.0	-0.00	2174.8
Max	SORTIE-free	7/25	0.36	2.7	+0.00	2177.1
Range	Det-lower	0/25	0.00	0.0	-23.11	1099.7
Range	Det-free	23/25	1.00	87.8	-24.19	1088.5
Range	SORTIE-lower	0/25	0.00	0.0	+0.04	1098.4
Range	SORTIE-free	24/25	1.64	108.4	+0.06	1072.6
GMD	Det-lower	0/25	0.00	0.0	-19.46	437.7
GMD	Det-free	21/25	1.12	90.0	-20.02	425.9
GMD	SORTIE-lower	0/25	0.00	0.0	+0.07	434.5
GMD	SORTIE-free	23/25	1.52	82.4	+0.10	438.6

Table 5: E6: arc noise v, $n = 25$, $B = 1.10$, five instances per row.

v	Measure	Det belief err (%)	SORTIE-0 vs Det
0.1	GMD	-6.4	+1.05 (2/5)
0.1	Range	-8.7	-0.54 (1/5)
0.3	GMD	-22.0	+0.65 (2/5)
0.3	Range	-28.7	+0.96 (3/5)
0.5	GMD	-35.6	+1.84 (4/5)
0.5	Range	-42.7	+2.14 (5/5)

no improving relocate or swap move in 28 of 30 cases: the search ends in genuine local optima of its neighbourhood. Consistently, the number of candidates K evaluated per step barely matters (Table 13): the median ex-post value is the same for K from 1 to 32.

6.13 Is a larger sample enough? (E15)

A natural objection is that SAA with 200 scenarios is simply too small. E15 reruns SAA with $S \in \{50, 200, 1000, 5000\}$ scenarios on the $n \in \{25, 35\}$ instances (Table 14, Figure 17). SORTIE-0 beats every SAA variant: by 1.4% ($S = 50$), 1.6% ($S = 200$), 1.1% ($S = 1000$) and 0.9% ($S = 5000$). The gap shrinks as the sample grows, as it should, since SAA converges to the objective that PBOI evaluates exactly; but with 5000 scenarios SAA still loses (better on 2, worse on 9 of 30 pairs, the rest tied). The sample average is cached per route, so the number of iterations drops only from about 5300 to 4200, and the remaining gap is the sampling error of the objective, not a lack of search time.

6.14 Simpler stochastic baselines (E16)

Two simpler ways of taking uncertainty into account are tested on the full E3 grid at $B = 1.10$ (Table 15). *Det-σ* applies the deterministic ordered measure to risk-adjusted lengths $L_k + \sigma_k$, the fairness analogue of the robust-length heuristics common in stochastic routing. It is *worse* than Det




Figure 7: Left: range believed by Det (ex-ante) and by SORTIE (PBOI) against the true expected range, E3. Right: share of the true ex-post unfairness that a deterministic plan does not see, against v (E6).

Table 6: E7: improvement of the ex-post value over Det, $B = 1.10$, 120 s, pooled over six instances and three measures (range, GMD, max); format as Table 3.

n	SAA	SORTIE-0	SORTIE-1
50	+1.56 (12/18)	+2.04* (13/18)	+1.20* (15/18)
75	+1.96 (11/18)	+2.04 (13/18)	-0.29 (11/18)
100	+1.52 (10/18)	+2.21 (12/18)	-0.04 (8/18)

(-0.8%, better on 37 and worse on 61 of 125 pairs, $p = 0.02$), and SORTIE-0 beats it by 1.9% (97 of 125, $p < 10^{-14}$). Adding a standard deviation to every route treats each route's risk separately, whereas the expected ordered measure depends on how likely each route is to occupy each rank, which is what the Poisson–binomial integral and the rank prices capture. *PBOI-N* replaces the gamma marginals by normal ones; it performs like SORTIE-0 (no significant difference, $p = 0.65$), so the method is insensitive to the marginal family once the first two moments are right.

6.15 Managerial insights

The experiments support five statements a planner can act on.

(i) Report realised, not planned, fairness. A deterministic fair plan understates the unfairness its drivers will experience by 8–27% at a moderate noise level, by more with more vehicles (E10) and more noise (E6), and the understatement is not uniform across plans. PBOI computes the realised value in under a millisecond, so it can be reported for any plan, whatever method produced it.

(ii) Do not spend the budget on deterministic balance. With generous budgets a deterministic fair model spends the slack to equalise route lengths, which does little for realised fairness and can make it worse (E9). The ex-post objective uses less of the budget and achieves more: up to 7% lower expected range with a total length 3.4 percentage points closer to C^* at $B = 1.30$.

(iii) Risk adjustment is not a substitute. Adding a safety margin to each route and then balancing (Det-σ) is worse than doing nothing (E16). What matters is the probability that each

Table 7: E7b, $n = 100$, $B = 1.10$, 240 s, three solver seeds: improvement of the seed-averaged ex-post value over Det (six instances per measure; format as Table 3) and coefficient of variation of the ex-post value across seeds, averaged over instances.

	vs Det, mean of 3 seeds		Seed CV (%)		
Measure	SAA	SORTIE-0	Det	SAA	SORTIE-0
Range	+6.45 (4/6)	+8.68 (5/6)	11.3	4.7	3.0
GMD	+2.49 (3/6)	+3.18 (4/6)	6.4	2.7	4.5
Max	+0.78 (4/6)	+0.86 (4/6)	1.4	0.8	1.4
All	+3.24 (11/18)	+4.24* (13/18)	SORTIE-0 vs SAA: +1.09 (12/18)		




Figure 8: E7: best in-model value relative to the final value over the run (median and interquartile range over the 18 runs per method and size).

route ends up the longest or the shortest, which is what the rank prices measure.

(iv) More scenarios are not a substitute either. Sample average approximation needs thousands of scenarios to approach the exact objective and still loses at 5000 (E15). The exact integral takes 0.4 ms per plan, whereas the 200-scenario average takes 0.06 ms, so a 5000-scenario average would, extrapolating linearly, cost several times more than PBOI.

(v) Keep the shortest-tour requirement. Without it, a fairness objective buys balance by sending drivers on detours (E4), which is exactly the behaviour the bilevel formulation of Ljubić et al. [2026] was designed to exclude; under uncertainty it must be imposed explicitly, and the risk-consistent variant changes few tours in practice (E13).

7 Conclusions

Fair routing under random travel times has to say whose fairness it measures. Drivers experience realised durations, so the relevant quantity is the expected unfairness of the realised day. A deterministic fair plan cannot see this quantity, and in our experiments it understated it by 8% (maximum) to 27% (variance) at $v = 0.3$, and by up to 43% at $v = 0.5$. A multiplicative common




Figure 9: E7b, $n = 100$: ex-post value of each solver seed, relative to the mean Det value of the instance; larger markers: seed means.

Table 8: E8: congested arc noise, $B = 1.10$, 25 instances per row; improvement of the ex-post value over Det (format as Table 3).

Measure	SAA	SORTIE-0	SORTIE-1
GMD	+0.76 (11/25)	+1.40* (13/25)	+1.82* (15/25)
Max	+1.87* (23/25)	+2.15* (23/25)	+2.21* (24/25)
Range	+1.22 (13/25)	+0.96* (17/25)	+0.94* (18/25)

shock, by contrast, leaves the expected value of every ordered measure unchanged (Proposition 2), although it does inflate the variance measure and the tail of unfairness.

The Poisson–binomial order-statistic integral evaluates any ordered measure of independent route durations exactly, in under a millisecond for five vehicles, and its derivatives, the rank prices, are both interpretable and useful: they identify the routes a fairness objective would like to lengthen, and screening moves with them gave plans 6–9% better on average than random screening. Under uncertainty, route consistency no longer follows from monotonicity, so it has to be part of the model; the mean–standard-deviation lower level makes it computable by parametric Held–Karp. Without a lower level, both the deterministic and the stochastic planners produced detours in almost every range-optimised plan.

Optimising ex-post fairness improved the realised fairness by about one percent over a deterministic fair plan under homogeneous noise (up to 2% for the maximum), consistently but modestly, because balancing route lengths already roughly balances route risks when variance grows with length. When noise is heterogeneous, as with congestion concentrated in part of the network, the illusion grows to 26% for the range (23% on the same cells without congestion), and the gain of SORTIE grows only to about 1.5%. The main practical value of the stochastic model in our experiments is therefore that it tells the planner, accurately, how unfair a plan will be, and that it guarantees consistent tours; the improvement of the plan itself is significant in the pooled compar-




Figure 10: X641-n30-b0 with congestion: customers coloured by the mean noise multiplier H of their arcs, with the range-optimised Det and SORTIE-0 plans of E8.

Table 9: E9: improvement of the ex-post value of SORTIE-0 over Det by budget factor, $n \in \{25, 35\}$, ten instances per measure; cost ratio: mean $\sum_k L_k/C^*$ of the SORTIE-0 plans (format as Table 3).

B	Cost ratio	Pooled	Range	GMD	Max
1.02	1.014	+1.43* (17/30)	+1.40 (4/10)	+2.07 (5/10)	+0.82* (8/10)
1.05	1.041	+0.73* (19/30)	+0.81 (5/10)	+0.66 (5/10)	+0.73* (9/10)
1.10	1.094	+0.67* (20/30)	-0.72 (7/10)	+0.30 (3/10)	+2.45* (10/10)
1.15	1.134	+3.20* (27/30)	+5.92* (8/10)	+1.31 (9/10)	+2.37* (10/10)
1.20	1.170	+2.46* (26/30)	+2.59 (9/10)	+2.34* (7/10)	+2.46* (10/10)
1.30	1.222	+6.79* (30/30)	+7.64* (10/10)	+9.11* (10/10)	+3.63* (10/10)

isons but small. On instances with 100 customers the faster search makes the improvement over the deterministic plan statistically significant when search noise is averaged over three seeds and the 18 instance–measure pairs are pooled (4%, $p = 0.02$), although no single measure is significant on six instances, and it does not separate SORTIE from a 200-scenario SAA at that size.

Limitations. The lower level is exact only for routes of up to 15 customers; beyond that it relies on multi-start local search (warm-started from the screening tour), and the bilevel constraint is then satisfied only approximately. PBOI assumes conditionally independent route durations; dependence beyond a common multiplicative factor (for example spatially correlated congestion affecting several routes) would need the copula or simulation layer we did not develop. At $n = 100$ single runs are still noisy and some are not converged after 240 s; the risk-consistent variant ($\kappa = 1$) is too slow to converge at $n \geq 75$; and at that size SORTIE could not be separated statistically from a 200-scenario SAA. The time limits are per run and are slightly exceeded by the final iteration. All instances are slices of one benchmark file with five vehicles.




Figure 11: Improvement of SORTIE-0 over Det per instance and measure, without (E3) and with (E8) congestion, $B = 1.10$.

Future work. Exact methods for the stochastic bilevel problem (branch-and-cut with PBOI-based bounds), correlated route durations, multi-day fairness, in which unfairness can be compensated across days, and time-dependent travel times.

References

Yossiri Adulyasak and Patrick Jaillet. Models and Algorithms for Stochastic and Robust Vehicle Routing with Deadlines. *Transportation Science*, 50(2):608–626, 2016. doi: 10.1287/trsc.2014.0581.

Maxime Agius, Nabil Absi, Dominique Feillet, and Thierry Garaix. Solution algorithms for a vehicle routing problem with route-cost equity constraints. *Annals of Operations Research*, 359 (3):2309–2349, 2026. doi: 10.1007/s10479-026-07145-6.

Douglas Alem, Aakil M. Caunhye, and Alfredo Moreno. Revisiting Gini for equitable humanitarian logistics. *Socio-Economic Planning Sciences*, 82:101312, 2022. doi: 10.1016/j.seps.2022.101312.

Aldair Alvarez, Jean-François Cordeau, and Raf Jans. The consistent vehicle routing problem with stochastic customers and demands. *Transportation Research Part B: Methodological*, 186:102968, 2024. doi: 10.1016/j.trb.2024.102968.

Yash P. Aneja and Kunhiraman P. K. Nair. Bicriteria Transportation Problem. *Management Science*, 25(1):73–78, 1979. doi: 10.1287/mnsc.25.1.73.




Figure 12: E9: price of fairness. Mean total length relative to C^* against the mean expected realised unfairness relative to the cost-optimal plan, for budget factors B from 1.02 to 1.30 (labels).

Table 10: E10: fleet size, $n = 35$, five instances per measure, $B = 1.10$ (format as Table 3).

m	Measure	Det belief err. (%)	SAA vs Det	SORTIE-0 vs Det
3	Pooled	-8.2	+2.83* (7/15)	+0.48* (8/15)
3	Range	-9.9	+3.66 (2/5)	+0.14 (2/5)
3	GMD	-9.9	+3.66 (2/5)	+0.14 (2/5)
3	Max	-4.6	+1.17 (3/5)	+1.17 (4/5)
5	Pooled	-18.6	-1.62 (8/15)	+0.64 (9/15)
5	Range	-24.1	-3.11 (2/5)	+2.78 (3/5)
5	GMD	-23.2	-4.01 (1/5)	-3.58 (1/5)
5	Max	-8.5	+2.26 (5/5)	+2.71 (5/5)
7	Pooled	-20.3	+2.34* (10/15)	+3.16* (13/15)
7	Range	-26.7	+1.36 (3/5)	+4.58 (5/5)
7	GMD	-23.3	+3.34 (2/5)	+2.14 (3/5)
7	Max	-10.9	+2.31 (5/5)	+2.76 (5/5)

Nguyen Duy Anh. Workload Balancing Among Heathcare Workers Under Uncertain Service Time Using Distributionally Robust Optimization. *Asia-Pacific Journal of Operational Research*, 39 (05):2150045, 2022. doi: 10.1142/s0217595921500457.

David Applegate, William Cook, Sanjeeb Dash, and André Rohe. Solution of a Min-Max Vehicle Routing Problem. *INFORMS Journal on Computing*, 14(2):132–143, 2002. doi: 10.1287/ijoc.14.2.132.118.

Iurii Bakach, Ann Melissa Campbell, Jan Fabian Ehmke, and Timothy L. Urban. Solving vehicle routing problems with stochastic and correlated travel times and makespan objectives. *EURO Journal on Transportation and Logistics*, 10:100029, 2021. doi: 10.1016/j.ejtl.2021.100029.

N. Balakrishnan. Permanents, Order Statistics, Outliers, and Robustness. *Revista Matemática Complutense*, 20(1):7–107, 2007. doi: 10.5209/rev_rema.2007.v20.n1.16528.

R. B. Bapat and M. I. Beg. Order statistics for nonidentically distributed variables and perma-




Figure 13: E10: belief error of Det and improvement over Det against the number of vehicles.

Table 11: E13: risk-consistent lower level, $n \in \{25, 35\}$, $B = 1.10$, ten instances per row. Routes changed: share of routes whose tour differs from the shortest tour of the same customers; extra length: total, per plan; mean risk: mean of $L_k + v\sqrt{W_k}$.

Measure	κ	Routes changed (%)	Extra length	Mean risk	Ex-post vs κ=0 [IQR]	Cost ratio
Range	0	0.0	0.0	1913.9	1.000 [1.000, 1.000]	1.098
Range	0.5	6.0	1.0	1912.9	1.000 [0.998, 1.000]	1.098
Range	1	4.0	0.9	1915.6	1.000 [0.995, 1.000]	1.098
Range	2	14.0	9.2	1914.0	1.000 [0.989, 1.004]	1.099
Range	4	14.0	14.1	1911.4	0.997 [0.989, 1.004]	1.099
Max	0	0.0	0.0	1890.7	1.000 [1.000, 1.000]	1.086
Max	0.5	2.0	0.5	1885.9	1.000 [1.000, 1.000]	1.084
Max	1	6.0	2.6	1891.3	1.000 [1.000, 1.000]	1.087
Max	2	6.0	3.9	1887.0	1.000 [1.000, 1.000]	1.085
Max	4	16.0	12.4	1882.6	1.000 [1.000, 1.000]	1.083

nents. *Sankhyā: The Indian Journal of Statistics, Series A*, 51(1):79–93, 1989. No DOI; JSTOR 25050725 (id as listed in search results; indexed in IAS repository).

Zohar Barak and Inbal Talgam-Cohen. Ex-ante versus Ex-post: Egalitarian Facility Location Mechanism Design. arXiv preprint, 2026.

Fuat Basik, Bugra Gedik, Hakan Ferhatosmanoglu, and Kun-Lung Wu. Fair Task Allocation in Crowdsourced Delivery. *IEEE Transactions on Services Computing*, 14(4):1040–1053, 2021. doi: 10.1109/tsc.2018.2854866.

Tolga Bektaş and Adam N. Letchford. Using ℓ_p-norms for fairness in combinatorial optimisation. *Computers & Operations Research*, 120:104975, 2020. doi: 10.1016/j.cor.2020.104975.

Richard Bellman. Dynamic Programming Treatment of the Travelling Salesman Problem. *Journal of the ACM*, 9(1):61–63, 1962. doi: 10.1145/321105.321111.




Figure 14: E13: effect of the drivers' risk weight κ.

Table 12: E14: rank-price prediction against exact evaluation. Groups: (instance, plan) pairs with at least one exactly improving move; top-8 hit: share of those groups in which the exact-best improving move is among the 8 best predicted moves (in brackets: expected share under a random order).

Measure	Plans	Moves	Groups with improving move	Spearman ρ	Top-8 hit (random)
Range	Cost plan	1448	10/10	0.93	0.80 (0.07)
Range	SORTIE-0 plan	346	1/10	0.80	0.00 (0.19)
Range	All plans	1794	11/20	0.92	0.73 (0.08)
GMD	Cost plan	1448	10/10	0.94	0.80 (0.07)
GMD	SORTIE-0 plan	333	0/10	0.78	– (–)
GMD	All plans	1781	10/20	0.94	0.80 (0.07)
Max	Cost plan	1448	10/10	0.89	0.80 (0.07)
Max	SORTIE-0 plan	554	1/10	0.89	0.00 (0.21)
Max	All plans	2002	11/20	0.89	0.73 (0.08)

Luca Bertazzi, Bruce Golden, and Xingyin Wang. Min–Max vs. Min–Sum Vehicle Routing: A worst-case analysis. *European Journal of Operational Research*, 240(2):372–381, 2015. doi: 10.1016/j.ejor.2014.07.025.

Dimitris Bertsimas, Vivek F. Farias, and Nikolaos Trichakis. The Price of Fairness. *Operations Research*, 59(1):17–31, 2011. doi: 10.1287/opre.1100.0865.

Abhay Singh Bhadoriya, Deepjyoti Deka, and Kaarthik Sundar. Equitable routing–rethinking the Multiple Traveling Salesman Problem. *Journal of the Operational Research Society*, pages 1–20, 2026. doi: 10.1080/01605682.2026.2708027. Published online 2026; first posted on arXiv April 2024.

William Biscarri, Sihai Dave Zhao, and Robert J. Brunner. A simple and fast method for computing the Poisson binomial distribution function. *Computational Statistics & Data Analysis*, 122:92–100, 2018. doi: 10.1016/j.csda.2018.01.007.

Table 13: E14: number of candidates K evaluated per step; ex-post value relative to $K = 8$, median and interquartile range over 20 instance–measure pairs.

K	Range median [IQR]	GMD median [IQR]
1	1.000 [1.000, 1.000]	1.000 [1.000, 1.000]
2	1.000 [1.000, 1.000]	1.000 [1.000, 1.000]
4	1.000 [1.000, 1.000]	1.000 [1.000, 1.000]
8	1.000 [1.000, 1.000]	1.000 [1.000, 1.000]
16	1.000 [1.000, 1.000]	1.000 [0.996, 1.000]
32	1.000 [0.994, 1.000]	1.000 [1.000, 1.000]




Figure 15: E14: predicted (rank prices) against exact change of the objective for 5577 moves.

Víctor Blanco, Alfredo Marín, and Justo Puerto. Intra-facility equity in discrete and continuous p-facility location problems. *Computers & Operations Research*, 162:106487, 2024. doi: 10.1016/j.cor.2023.106487.

Víctor Blanco, Miguel A. Pozo, Justo Puerto, and Alberto Torrejon. Optimization-based computation and integration of ordered measures. *Annals of Operations Research*, 364(3):2799–2834, 2026. doi: 10.1007/s10479-026-07314-7.

Federica Bomboi, Christoph Buchheim, and Jonas Pruente. On the stochastic vehicle routing problem with time windows, correlated travel times, and time dependency. *4OR*, 20(2):217–239, 2021. doi: 10.1007/s10288-021-00476-z.

Tom Bruinink, Lotte Berghman, and Twan Dollevoet. Beyond efficiency: Exploring the cost effects of prioritizing driver satisfaction in vehicle routing. *Operations Research Letters*, 63:107344, 2025. doi: 10.1016/j.orl.2025.107344.

H. A. David and H. N. Nagaraja. *Order Statistics*. Wiley Series in Probability and Statistics. Wiley, 3 edition, 2003. doi: 10.1002/0471722162.

Jan Fabian Ehmke, Ann Melissa Campbell, and Timothy L. Urban. Ensuring service levels in routing problems with time windows and stochastic travel times. *European Journal of Operational Research*, 240(2):539–550, 2015. doi: 10.1016/j.ejor.2014.06.045.




Figure 16: E14: share of cases in which the exact-best improving move is among the K best predicted moves.

Table 14: E15: improvement of SORTIE-0 over SAA with S scenarios, $n \in \{25, 35\}$, $B = 1.10$, ten instances per measure (format as Table 3; positive: SORTIE-0 better).

S	Range	GMD	Max	Pooled	p (pooled)
50	+0.17 (6/10)	+2.88* (8/10)	+0.99* (8/10)	+1.35* (22/30)	0.001
200	+2.11* (7/10)	+2.15* (7/10)	+0.46* (10/10)	+1.57* (24/30)	0.000
1000	+1.45 (5/10)	+1.64 (4/10)	+0.19 (6/10)	+1.09* (15/30)	0.002
5000	+1.34 (3/10)	+1.38 (2/10)	+0.02 (4/10)	+0.91 (9/30)	0.050

Matthias Ehrgott. *Multicriteria Optimization*. Springer, Berlin, Heidelberg, 2 edition, 2005. doi: 10.1007/3-540-27659-9. Author field absent in DOI metadata; ISBN 3540213988.

Matthias Ehrgott and Xavier Gandibleux. A survey and annotated bibliography of multiobjective combinatorial optimization. *OR Spectrum*, 22(4):425–460, 2000. doi: 10.1007/s002910000046.

F. Errico, G. Desaulniers, M. Gendreau, W. Rei, and L.-M. Rousseau. A priori optimization with recourse for the vehicle routing problem with hard time windows and stochastic service times. *European Journal of Operational Research*, 249(1):55–66, 2016. doi: 10.1016/j.ejor.2015.07.027.

F. Errico, G. Desaulniers, M. Gendreau, W. Rei, and L.-M. Rousseau. The vehicle routing problem with hard time windows and stochastic service times. *EURO Journal on Transportation and Logistics*, 7(3):223–251, 2018. doi: 10.1007/s13676-016-0101-4.

Manuel Fernandez and Stuart Williams. Closed-Form Expression for the Poisson-Binomial Probability Density Function. *IEEE Transactions on Aerospace and Electronic Systems*, 46(2):803–817, 2010. doi: 10.1109/taes.2010.5461658.

Marc Fleurbaey and Vito Peragine. Ex Ante Versus Ex Post Equality of Opportunity. *Economica*, 80(317):118–130, 2012. doi: 10.1111/j.1468-0335.2012.00941.x.




Figure 17: E15: ex-post value of SAA relative to SORTIE-0 against the sample size S.

Table 15: E16: risk-adjusted deterministic (Det-σ) and normal-marginal (PBOI-N) objectives, E3 grid, $B = 1.10$ (format as Table 3; the variance is omitted for PBOI-N because the expected empirical variance does not depend on the marginal family).

Measure	Det-σ vs Det	PBOI-N vs Det	SORTIE-0 vs Det	SORTIE-0 vs Det-σ
Range	-0.49 (8/25)	+0.48* (18/25)	+1.07* (17/25)	+1.51* (19/25)
GMD	-0.69 (8/25)	+0.83* (14/25)	+0.94* (14/25)	+1.50* (18/25)
Max	-0.08 (8/25)	+2.08* (24/25)	+2.08* (24/25)	+2.13* (25/25)
Pooled	-0.81† (37/125)	+1.10* (71/100)	+1.23* (81/125)	+1.86* (97/125)

Haiqing Gao, Seyed M. R. Iravani, and Sean R. Sinclair. Sequential Fair Allocation and Routing in Nonprofit Operations. arXiv preprint, 2026.

Michel Gendreau, Ola Jabali, and Walter Rei. 50th Anniversary Invited Article—Future Research Directions in Stochastic Vehicle Routing. *Transportation Science*, 50(4):1163–1173, 2016. doi: 10.1287/trsc.2016.0709.

Shubhechyya Ghosal and Wolfram Wiesemann. The Distributionally Robust Chance-Constrained Vehicle Routing Problem. *Operations Research*, 68(3):716–732, 2020. doi: 10.1287/opre.2019.1924.

D. H. Glueck, A. Karimpour-Fard, J. Mandel, L. Hunter, and K. E. Muller. Fast Computation by Block Permanents of Cumulative Distribution Functions of Order Statistics from Several Populations. *Communications in Statistics - Theory and Methods*, 37(18):2815–2824, 2008. doi: 10.1080/03610920802001896.

Chris Groër, Bruce Golden, and Edward Wasil. The Consistent Vehicle Routing Problem. *Manufacturing & Service Operations Management*, 11(4):630–643, 2009. doi: 10.1287/msom.1080.0243.

Walter J. Gutjahr. Inequity-averse stochastic decision processes. *European Journal of Operational Research*, 288(1):258–270, 2021. doi: 10.1016/j.ejor.2020.05.045.

Andrés Gómez, Ricardo Mariño, Raha Akhavan-Tabatabaei, Andrés L. Medaglia, and Jorge E. Mendoza. On Modeling Stochastic Travel and Service Times in Vehicle Routing. *Transportation Science*, 50(2):627–641, 2016. doi: 10.1287/trsc.2015.0601.

Elin E. Halvorsen-Weare and Martin W.P. Savelsbergh. The bi-objective mixed capacitated general routing problem with different route balance criteria. *European Journal of Operational Research*, 251(2):451–465, 2016. doi: 10.1016/j.ejor.2015.11.024.

Jinil Han, Chungmok Lee, and Sungsoo Park. A Robust Scenario Approach for the Vehicle Routing Problem with Uncertain Travel Times. *Transportation Science*, 48(3):373–390, 2014. doi: 10.1287/trsc.2013.0476.

Michael Held and Richard M. Karp. A Dynamic Programming Approach to Sequencing Problems. *Journal of the Society for Industrial and Applied Mathematics*, 10(1):196–210, 1962. doi: 10.1137/0110015.

Wassily Hoeffding. On the Distribution of the Number of Successes in Independent Trials. *The Annals of Mathematical Statistics*, 27(3):713–721, 1956. doi: 10.1214/aoms/1177728178.

Yili Hong. On computing the distribution function for the Poisson binomial distribution. *Computational Statistics & Data Analysis*, 59:41–51, 2013. doi: 10.1016/j.csda.2012.10.006.

Hadi Hosseini and Šimon Schierreich. The Algorithmic Landscape of Fair and Efficient Distribution of Delivery Orders in the Gig Economy. arXiv preprint, 2025.

Patrick Jaillet, Jin Qi, and Melvyn Sim. Routing Optimization Under Uncertainty. *Operations Research*, 64(1):186–200, 2016. doi: 10.1287/opre.2015.1462.

Özlem Karsu and Alec Morton. Inequity averse optimization in operational research. *European Journal of Operational Research*, 245(2):343–359, 2015. doi: 10.1016/j.ejor.2015.02.035.

Astrid S. Kenyon and David P. Morton. Stochastic Vehicle Routing with Random Travel Times. *Transportation Science*, 37(1):69–82, 2003. doi: 10.1287/trsc.37.1.69.12820.

Joris Kinable, Natasja Sluijk, Michel Gendreau, Walter Rei, and Tom Van Woensel. Fair Stochastic Vehicle Routing with Partial Deliveries. *Transportation Science*, 60(2):264–283, 2026. doi: 10.1287/trsc.2024.0556.

Attila A. Kovacs, Bruce L. Golden, Richard F. Hartl, and Sophie N. Parragh. The Generalized Consistent Vehicle Routing Problem. *Transportation Science*, 49(4):796–816, 2015. doi: 10.1287/trsc.2014.0529.

Gilbert Laporte, François Louveaux, and Hélène Mercure. The Vehicle Routing Problem with Stochastic Travel Times. *Transportation Science*, 26(3):161–170, 1992. doi: 10.1287/trsc.26.3.161.

Shaochong Lin, Monique Bakker, and Eman Leung. Robust meal delivery service for the elderly: a case study in Hong Kong. *Scientific Reports*, 12(1):21546, 2022. doi: 10.1038/s41598-022-25924-6.

Ivana Ljubić, Justo Puerto, and Alberto Torrejón. Fair Vehicle Routing via Bilevel Optimization. Optimization Online preprint, submitted 5 March 2026, 2026. URL https://optimization-online.org/2026/03/fair-vehicle-routing-via-bilevel-optimization/.

Ivana Ljubić, Miguel A. Pozo, Justo Puerto, and Alberto Torrejón. Benders decomposition for the discrete ordered median problem. *European Journal of Operational Research*, 317(3):858–874, 2024. doi: 10.1016/j.ejor.2024.04.030.

Ivana Ljubić, Alfredo Marín, Justo Puerto, and Francisco Temprano. Ordered Median Traveling Salesman Problem. *Networks*, 88(1):59–84, 2026. doi: 10.1002/net.70036.

Aitor López Sánchez, Marin Lujak, Frederic Semet, and Holger Billhardt. How to achieve fair and efficient cooperative vehicle routing? *AI Communications*, 37(2):223–245, 2024. doi: 10.3233/aic-220315.

Vahid Mahmoodian, Hadi Charkhgard, and Iman Dayarian. Equitable Workload Allocation in Vehicle Routing Problem With Heterogeneous Drivers. *Production and Operations Management*, 34(7):1875–1900, 2025. doi: 10.1177/10591478241305873.

P. Matl, R. F. Hartl, and T. Vidal. Workload Equity in Vehicle Routing Problems: A Survey and Analysis. *Transportation Science*, 52(2):239–260, 2018. doi: 10.1287/trsc.2017.0744.

P. Matl, R.F. Hartl, and T. Vidal. Workload equity in vehicle routing: The impact of alternative workload resources. *Computers & Operations Research*, 110:116–129, 2019a. doi: 10.1016/j.cor.2019.05.016.

Piotr Matl, Richard F. Hartl, and Thibaut Vidal. Leveraging single-objective heuristics to solve bi-objective problems: Heuristic box splitting and its application to vehicle routing. *Networks*, 73(4):382–400, 2019b. doi: 10.1002/net.21876.

J. Mo and J. Walrand. Fair end-to-end window-based congestion control. *IEEE/ACM Transactions on Networking*, 8(5):556–567, 2000. doi: 10.1109/90.879343.

Mahdi Mostajabdaveh, Walter J. Gutjahr, and F. Sibel Salman. Inequity-averse shelter location for disaster preparedness. *IISE Transactions*, 51(8):809–829, 2019. doi: 10.1080/24725854.2018.1496372.

Mahdi Mostajabdaveh, F. Sibel Salman, and Walter J. Gutjahr. A branch-and-price algorithm for fast and equitable last-mile relief aid distribution. *European Journal of Operational Research*, 324(2):522–537, 2025. doi: 10.1016/j.ejor.2025.01.032.

Vedant Nanda, Pan Xu, Karthik Abhinav Sankararaman, John Dickerson, and Aravind Srinivasan. Balancing the Tradeoff between Profit and Fairness in Rideshare Platforms during High-Demand Hours. In *Proceedings of the AAAI Conference on Artificial Intelligence*, pages 2210–2217. Association for the Advancement of Artificial Intelligence (AAAI), 2020. doi: 10.1609/aaai.v34i02.5597.

Najmeh Nekooghadirli, Michel Gendreau, Jean-Yves Potvin, and Thibaut Vidal. Workload equity in multiperiod vehicle routing problems. *International Transactions in Operational Research*, 33 (1):245–267, 2026. doi: 10.1111/itor.70012. Published online 2025; volume 33 (2026).

Stefan Nickel and Justo Puerto. *Location Theory: A Unified Approach*. Springer, Berlin, Heidelberg, 2005. doi: 10.1007/3-540-27640-8. Hardcover 2005; eBook 2006 (cited as 2006 in Ljubić et al.).

E. Nikolova and N. E. Stier-Moses. A Mean-Risk Model for the Traffic Assignment Problem with Stochastic Travel Times. *Operations Research*, 62(2):366–382, 2014. doi: 10.1287/opre.2013.1246.

Wlodzimierz Ogryczak and Arie Tamir. Minimizing the sum of the k largest functions in linear time. *Information Processing Letters*, 85(3):117–122, 2003. doi: 10.1016/s0020-0190(02)00370-8.

Włodzimierz Ogryczak. Inequality measures and equitable approaches to location problems. *European Journal of Operational Research*, 122(2):374–391, 2000. doi: 10.1016/s0377-2217(99)00240-4.

Włodzimierz Ogryczak and Tomasz Śliwiński. On solving linear programs with the ordered weighted averaging objective. *European Journal of Operational Research*, 148(1):80–91, 2003. doi: 10.1016/s0377-2217(02)00399-5.

Alan Osorio-Mora, Francisco Saldanha-da Gama, and Paolo Toth. A risk-averse latency location-routing problem with stochastic travel times. *European Journal of Operational Research*, 321(3):837–850, 2025. doi: 10.1016/j.ejor.2024.10.041.

Jorge Oyola, Halvard Arntzen, and David L. Woodruff. The stochastic vehicle routing problem, a literature review, part I: models. *EURO Journal on Transportation and Logistics*, 7(3):193–221, 2018. doi: 10.1007/s13676-016-0100-5.

Mojtaba Rajabi-Bahaabadi, Afshin Shariat-Mohaymany, Mohsen Babaei, and Daniele Vigo. Reliable vehicle routing problem in stochastic networks with correlated travel times. *Operational Research*, 21(1):299–330, 2019. doi: 10.1007/s12351-019-00452-w.

Ted K. Ralphs, Matthew J. Saltzman, and Margaret M. Wiecek. An improved algorithm for solving biobjective integer programs. *Annals of Operations Research*, 147(1):43–70, 2006. doi: 10.1007/s10479-006-0058-z.

R. Tyrrell Rockafellar and Stanislav Uryasev. Optimization of conditional value-at-risk. *The Journal of Risk*, 2(3):21–41, 2000. doi: 10.21314/jor.2000.038.

R.Tyrrell Rockafellar and Stanislav Uryasev. Conditional value-at-risk for general loss distributions. *Journal of Banking & Finance*, 26(7):1443–1471, 2002. doi: 10.1016/s0378-4266(02)00271-6.

Borzou Rostami, Guy Desaulniers, Fausto Errico, and Andrea Lodi. Branch-Price-and-Cut Algorithms for the Vehicle Routing Problem with Stochastic and Correlated Travel Times. *Operations Research*, 69(2):436–455, 2021. doi: 10.1287/opre.2020.2037.

Pranab Kumar Sen. A Note on Order Statistics for Heterogeneous Distributions. *The Annals of Mathematical Statistics*, 41(6):2137–2139, 1970. doi: 10.1214/aoms/1177696715.

Suvrajeet Sen, Rekha Pillai, Shirish Joshi, and Ajay K. Rathi. A Mean-Variance Model for Route Guidance in Advanced Traveler Information Systems. *Transportation Science*, 35(1):37–49, 2001. doi: 10.1287/trsc.35.1.37.10141.

Karmel S. Shehadeh. Reducing disparities in transportation distance in a stochastic facility location problem. *Transportation Research Part C: Emerging Technologies*, 153:104199, 2023. doi: 10.1016/j.trc.2023.104199.

Sean R. Sinclair, Gauri Jain, Siddhartha Banerjee, and Christina Lee Yu. Sequential Fair Allocation: Achieving the Optimal Envy-Efficiency Trade-off Curve. *Operations Research*, 71(5):1689–1705, 2023. doi: 10.1287/opre.2022.2397.

Daman Deep Singh, Syamantak Das, and Abhijnan Chakraborty. FairAssign: Stochastically Fair Driver Assignment in Gig Delivery Platforms. In *Proceedings of the 2023 ACM Conference on Fairness, Accountability, and Transparency (FAccT)*, pages 753–763. ACM, 2023. doi: 10.1145/3593013.3594040.

Abhay Sobhanan, Hadi Charkhgard, and Iman Dayarian. Equity-Driven Workload Allocation for Crowdsourced Last-Mile Delivery. *Production and Operations Management*, 35(8):3180–3203, 2026. doi: 10.1177/10591478261425875.

Man Yiu Tsang and Karmel S. Shehadeh. A Unified Framework for Analyzing and Optimizing a Class of Convex Fairness Measures. *Operations Research*, 74(2):1087–1103, 2026. doi: 10.1287/opre.2023.0301.

Eduardo Uchoa, Diego Pecin, Artur Pessoa, Marcus Poggi, Thibaut Vidal, and Anand Subramanian. New benchmark instances for the Capacitated Vehicle Routing Problem. *European Journal of Operational Research*, 257(3):845–858, 2017. doi: 10.1016/j.ejor.2016.08.012.

Bart van Rossum, Rui Chen, and Andrea Lodi. A New Branching Rule for Range Minimization Problems. In *Integer Programming and Combinatorial Optimization (IPCO 2024)*, Lecture Notes in Computer Science, pages 433–445. Springer Nature Switzerland, 2024. doi: 10.1007/978-3-031-59835-7_32.

Bart van Rossum, Rui Chen, and Andrea Lodi. Efficient branching rules for optimizing range and order-based objective functions. *Mathematical Programming*, 216(1-2):539–572, 2026a. doi: 10.1007/s10107-025-02306-0. Online 2025; volume 216 (2026).

Bart van Rossum, Rui Chen, and Andrea Lodi. Enforcing TSP-optimality in fair vehicle routing by cutting planes. *Operations Research Letters*, 70:107539, 2026b. doi: 10.1016/j.orl.2026.107539. Published online 2026; Crossref dates volume 70 as January 2027.

R. J. Vaughan and W. N. Venables. Permanent Expressions for Order Statistic Densities. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 34(2):308–310, 1972. doi: 10.1111/j.2517-6161.1972.tb00909.x.

Bram Verweij, Shabbir Ahmed, Anton J. Kleywegt, George Nemhauser, and Alexander Shapiro. The Sample Average Approximation Method Applied to Stochastic Routing Problems: A Computational Study. *Computational Optimization and Applications*, 24(2-3):289–333, 2003. doi: 10.1023/a:1021814225969.

Shiping Wu, Chun Jin, and Hongguang Bo. Exact solution of workload consistent vehicle routing problem with priority distribution and demand uncertainty. *Computers & Industrial Engineering*, 202:110940, 2025. doi: 10.1016/j.cie.2025.110940.

R.R. Yager. On ordered weighted averaging aggregation operators in multicriteria decisionmaking. *IEEE Transactions on Systems, Man, and Cybernetics*, 18(1):183–190, 1988. doi: 10.1109/21.87068.

Qing Ye, Grani A. Hanasusanto, and Weijun Xie. Distributionally Fair Stochastic Optimization using Wasserstein Distance. arXiv preprint, 2024.

Xiao-Pu Yu, Yong-Shi Hu, and Peng Wu. The consistent vehicle routing problem considering driver equity and flexible route consistency. *Computers & Industrial Engineering*, 187:109803, 2024. doi: 10.1016/j.cie.2023.109803.

Dali Zhang, Dong Li, Hailin Sun, and Liwen Hou. A vehicle routing problem with distribution uncertainty in deadlines. *European Journal of Operational Research*, 292(1):311–326, 2021. doi: 10.1016/j.ejor.2020.10.026.

Junlong Zhang, William H. K. Lam, and Bi Yu Chen. A Stochastic Vehicle Routing Problem with Travel Time Uncertainty: Trade-Off Between Cost and Customer Service. *Networks and Spatial Economics*, 13(4):471–496, 2013. doi: 10.1007/s11067-013-9190-x.

A Per-instance results




Figure 18: E3: improvement of the ex-post value of SORTIE-0 over Det (%) for every instance, measure and budget.

Tables 16 and 17 list the out-of-sample ex-post value of every E3 plan, Table 18 the seed means and standard deviations of E7b. Every plan is stored in the released plan cache and can be re-evaluated with the released simulator.

Table 16: E3, $B = 1.10$: out-of-sample ex-post value of every plan (bold: best); Cost: the cost-optimal plan; S-0/S-1: SORTIE-0/1.

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
n15-b0	Max	2,640	2,547	2,561	2,547	**2,545**	+0.00	1.026
	Range	2,412	1,575	**1,568**	**1,568**	1,573	+0.47	1.098
	GMD	907	625	625	625	**623**	+0.06	1.100
	MAD	652	468	**467**	**467**	467	+0.20	1.100
	Var	730,909	357,317	357,317	357,238	**354,022**	+0.02	1.100
n15-b1	Max	2,165	2,091	2,095	**2,091**	**2,091**	+0.00	1.098
	Range	1,875	1,295	**1,285**	**1,285**	**1,285**	+0.79	1.095
	GMD	729	**530**	**530**	**530**	**530**	+0.00	1.095
	MAD	537	**405**	**405**	**405**	**405**	+0.00	1.095

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
	Var	461,317	**258,741**	261,374	**258,741**	**258,741**	+0.00	1.095
n15-b2	Max	2,419	2,419	2,376	**2,376**	**2,376**	+1.78	1.070
	Range	2,187	**1,231**	1,234	**1,231**	**1,231**	+0.00	1.081
	GMD	804	**478**	482	**478**	479	+0.00	1.099
	MAD	568	**347**	**347**	**347**	348	+0.00	1.099
	Var	604,666	**222,814**	223,451	**222,814**	223,451	+0.00	1.099
n15-b3	Max	1,926	1,826	1,826	1,805	**1,803**	+1.13	1.070
	Range	1,899	784	806	779	**773**	+0.64	1.070
	GMD	665	305	305	303	**300**	+0.59	1.070
	MAD	452	226	224	223	**221**	+1.15	1.070
	Var	471,206	90,921	95,342	90,921	**89,222**	+0.00	1.095
n15-b4	Max	2,412	1,940	**1,934**	1,935	1,935	+0.25	1.056
	Range	2,170	**1,024**	**1,024**	**1,024**	**1,024**	+0.00	1.098
	GMD	772	**417**	**417**	**417**	**417**	+0.00	1.098
	MAD	531	**316**	**316**	**316**	**316**	+0.00	1.098
	Var	543,628	**164,264**	**164,264**	**164,264**	**164,264**	+0.00	1.098
n20-b0	Max	2,656	2,651	2,647	**2,628**	**2,628**	+0.90	1.081
	Range	2,428	1,549	1,537	1,526	**1,522**	+1.48	1.097
	GMD	926	628	631	628	**626**	+0.00	1.099
	MAD	672	477	481	477	**477**	+0.00	1.099
	Var	782,670	358,873	361,706	358,394	**356,722**	+0.13	1.097
n20-b1	Max	1,756	1,743	**1,668**	1,669	1,669	+4.28	1.037
	Range	1,332	563	562	550	**550**	+2.42	1.099
	GMD	472	217	217	214	**214**	+1.40	1.077
	MAD	323	157	159	**157**	**157**	+0.04	1.077
	Var	230,329	47,975	47,125	**46,117**	**46,117**	+3.87	1.077
n20-b2	Max	2,393	2,224	2,177	**2,158**	**2,158**	+2.97	1.096
	Range	2,309	**1,283**	**1,283**	1,299	1,293	-1.24	1.090
	GMD	886	558	**552**	553	553	+0.92	1.099
	MAD	645	387	**383**	383	383	+1.02	1.099
	Var	692,985	257,489	**252,913**	**252,913**	**252,913**	+1.78	1.098
n20-b3	Max	2,668	2,472	2,349	**2,348**	**2,348**	+5.02	1.084
	Range	2,518	**1,272**	1,278	**1,272**	**1,272**	+0.00	1.100
	GMD	918	523	517	**516**	523	+1.42	1.099
	MAD	644	373	**372**	**372**	374	+0.17	1.099
	Var	757,430	250,084	247,116	247,116	**245,946**	+1.19	1.098
n20-b4	Max	1,981	1,823	1,782	1,778	**1,776**	+2.48	1.068
	Range	1,472	1,085	1,066	1,062	**1,060**	+2.13	1.068
	GMD	617	399	398	397	**396**	+0.52	1.067
	MAD	477	282	**282**	**282**	**282**	+0.14	1.098
	Var	332,502	147,358	148,357	147,358	**146,846**	+0.00	1.067
n25-b0	Max	2,607	2,622	2,613	**2,588**	**2,588**	+1.30	1.095
	Range	2,379	1,253	1,260	**1,252**	1,253	+0.11	1.100
	GMD	858	**506**	508	**506**	507	+0.00	1.100
	MAD	597	381	383	381	**381**	+0.00	1.100
	Var	718,676	245,392	248,480	**243,295**	**243,295**	+0.85	1.099
n25-b1	Max	2,145	1,971	1,939	**1,931**	**1,931**	+2.05	1.027
	Range	1,257	618	615	615	**611**	+0.54	1.098
	GMD	535	247	238	239	**236**	+3.19	1.098
	MAD	417	179	176	175	**173**	+2.08	1.098
	Var	255,701	58,275	60,272	58,275	**57,003**	+0.00	1.098
n25-b2	Max	2,148	2,065	2,035	**2,033**	**2,033**	+1.56	1.097

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
	Range	2,036	**1,312**	1,312	1,312	1,312	-0.01	1.099
	GMD	763	539	539	539	**537**	+0.00	1.092
	MAD	546	412	409	409	**408**	+0.62	1.092
	Var	516,255	265,863	267,817	266,385	**264,172**	-0.20	1.099
n25-b3	Max	1,984	2,025	1,958	**1,946**	**1,946**	+3.90	1.097
	Range	1,474	874	840	838	**836**	+4.13	1.099
	GMD	560	331	333	**328**	331	+1.06	1.098
	MAD	405	248	244	**242**	**242**	+2.15	1.098
	Var	283,090	**104,507**	107,279	**104,507**	105,353	+0.00	1.098
n25-b4	Max	2,151	2,197	2,143	2,137	**2,137**	+2.74	1.099
	Range	1,931	1,002	1,004	999	**994**	+0.21	1.099
	GMD	683	393	394	393	**391**	+0.00	1.100
	MAD	468	291	292	291	**290**	+0.00	1.100
	Var	459,013	152,985	153,860	150,758	**149,397**	+1.46	1.099
n30-b0	Max	2,773	2,615	2,587	**2,585**	**2,585**	+1.14	1.050
	Range	2,626	1,229	1,228	1,223	**1,223**	+0.47	1.099
	GMD	1,000	497	497	**493**	493	+0.79	1.099
	MAD	724	392	372	**371**	372	+5.22	1.099
	Var	911,746	235,555	233,126	**231,861**	**231,861**	+1.57	1.099
n30-b1	Max	2,657	2,574	2,540	**2,516**	2,521	+2.23	1.075
	Range	2,287	1,537	1,421	**1,420**	**1,420**	+7.63	1.095
	GMD	913	568	551	551	**550**	+3.06	1.100
	MAD	684	411	**409**	411	412	+0.00	1.098
	Var	730,545	297,591	298,366	277,261	**275,214**	+6.83	1.095
n30-b2	Max	1,957	1,838	1,784	1,784	**1,783**	+2.94	1.078
	Range	1,838	1,002	988	988	**961**	+1.44	1.091
	GMD	667	365	372	365	**365**	+0.00	1.091
	MAD	466	259	263	259	**259**	+0.00	1.091
	Var	419,940	126,369	126,369	126,369	**126,081**	+0.00	1.091
n30-b3	Max	2,401	2,350	2,307	**2,300**	**2,300**	+2.13	1.097
	Range	2,371	1,264	**1,223**	**1,223**	1,284	+3.25	1.100
	GMD	853	532	502	**496**	500	+6.75	1.100
	MAD	592	396	**369**	375	**369**	+5.12	1.100
	Var	717,888	**225,491**	229,750	**225,491**	228,530	+0.00	1.100
n30-b4	Max	2,325	2,231	2,196	2,193	**2,191**	+1.69	1.068
	Range	2,100	**1,217**	1,237	1,232	1,231	-1.21	1.093
	GMD	762	494	502	493	**492**	+0.27	1.098
	MAD	533	347	351	344	**344**	+0.90	1.100
	Var	536,206	225,051	219,695	**216,280**	225,293	+3.90	1.099
n35-b0	Max	2,892	2,717	2,704	**2,682**	**2,682**	+1.27	1.098
	Range	2,744	**1,376**	1,741	1,479	1,724	-7.46	1.096
	GMD	1,013	**529**	609	**529**	609	+0.00	1.100
	MAD	718	421	429	428	**385**	-1.72	1.098
	Var	964,248	**259,530**	372,532	**259,530**	368,171	+0.00	1.100
n35-b1	Max	2,543	2,127	2,090	**2,087**	**2,087**	+1.91	1.096
	Range	2,429	**825**	**825**	**825**	**825**	+0.00	1.098
	GMD	946	**323**	324	**323**	**323**	+0.00	1.098
	MAD	696	**238**	239	**238**	**238**	+0.00	1.098
	Var	758,306	**102,506**	102,787	**102,506**	**102,506**	+0.00	1.098
n35-b2	Max	1,922	1,794	1,742	**1,741**	1,814	+2.91	1.061
	Range	1,526	553	546	**538**	**538**	+2.61	1.093
	GMD	562	209	**208**	**208**	**208**	+0.48	1.093

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
	MAD	398	**153**	153	153	153	-0.06	1.093
	Var	296,372	43,908	43,424	**42,644**	**42,644**	+2.88	1.095
n35-b3	Max	2,527	2,452	2,398	2,379	**2,379**	+2.98	1.100
	Range	2,137	1,066	1,012	**998**	**998**	+6.37	1.100
	GMD	780	**377**	381	**377**	**377**	+0.00	1.100
	MAD	548	291	273	273	**273**	+6.35	1.100
	Var	555,805	**142,628**	144,945	**142,628**	**142,628**	+0.00	1.100
n35-b4	Max	2,510	2,186	2,142	**2,133**	**2,133**	+2.43	1.085
	Range	2,436	702	702	**687**	**687**	+2.03	1.097
	GMD	902	272	275	**264**	**264**	+2.92	1.097
	MAD	640	194	**193**	**193**	**193**	+0.58	1.097
	Var	772,452	71,590	72,810	**69,523**	**69,523**	+2.89	1.097

Table 17: E3, $B = 1.05$: out-of-sample ex-post value of every plan (bold: best); Cost: the cost-optimal plan; S-0/S-1: SORTIE-0/1.

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
n15-b0	Max	2,640	**2,547**	2,559	**2,547**	**2,547**	+0.00	1.026
	Range	2,412	**2,319**	2,331	**2,319**	**2,319**	+0.00	1.026
	GMD	907	**868**	873	**868**	**868**	+0.00	1.026
	MAD	652	**621**	**621**	**621**	**621**	+0.00	1.026
	Var	730,909	**691,734**	699,905	**691,734**	**691,734**	+0.00	1.026
n15-b1	Max	2,165	**2,091**	**2,091**	**2,091**	**2,091**	+0.00	1.041
	Range	1,875	**1,503**	**1,503**	**1,503**	1,505	+0.00	1.040
	GMD	729	**616**	**616**	**616**	617	+0.00	1.040
	MAD	537	469	469	**467**	468	+0.50	1.048
	Var	461,317	337,238	339,053	**336,907**	337,815	+0.10	1.048
n15-b2	Max	2,419	**2,419**	**2,419**	**2,419**	**2,419**	+0.00	1.000
	Range	2,187	**2,187**	**2,187**	**2,187**	**2,187**	+0.00	1.000
	GMD	804	**804**	**804**	**804**	**804**	+0.00	1.000
	MAD	568	**568**	**568**	**568**	**568**	+0.00	1.000
	Var	604,666	**604,666**	**604,666**	**604,666**	**604,666**	+0.00	1.000
n15-b3	Max	1,926	**1,926**	**1,926**	**1,926**	**1,926**	+0.00	1.000
	Range	1,899	1,181	**1,177**	**1,177**	**1,177**	+0.37	1.043
	GMD	665	468	**468**	468	468	+0.00	1.041
	MAD	452	**349**	349	**349**	**349**	+0.00	1.041
	Var	471,206	200,220	**198,775**	**198,775**	**198,775**	+0.72	1.043
n15-b4	Max	2,412	**1,940**	**1,940**	**1,940**	**1,940**	+0.00	1.000
	Range	2,170	**1,698**	**1,698**	**1,698**	**1,698**	+0.00	1.000
	GMD	772	**648**	**648**	**648**	**648**	+0.00	1.022
	MAD	531	**466**	**466**	**466**	**466**	+0.00	1.022
	Var	543,628	**375,246**	**375,246**	**375,246**	**375,246**	+0.00	1.000
n20-b0	Max	2,656	**2,639**	2,647	**2,639**	**2,639**	+0.00	1.021
	Range	2,428	2,411	2,418	**2,405**	**2,405**	+0.25	1.022
	GMD	926	**863**	**863**	**863**	**863**	+0.00	1.049
	MAD	672	**595**	**595**	**595**	**595**	+0.00	1.049
	Var	782,670	748,918	751,999	**748,239**	**748,239**	+0.09	1.022
n20-b1	Max	1,756	1,751	**1,668**	1,669	1,669	+4.72	1.037
	Range	1,332	666	**652**	**652**	**652**	+2.20	1.036
	GMD	472	251	**251**	251	251	+0.00	1.049

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
	MAD	323	185	184	**183**	**183**	+0.99	1.049
	Var	230,329	62,141	**61,759**	**61,759**	**61,759**	+0.62	1.036
n20-b2	Max	2,393	**2,191**	2,202	2,279	2,279	-4.03	1.039
	Range	2,309	**1,652**	1,988	1,988	1,988	-20.38	1.039
	GMD	886	**656**	**656**	**656**	**656**	+0.00	1.049
	MAD	645	505	**490**	**490**	**490**	+2.92	1.049
	Var	692,985	**379,751**	**379,751**	**379,751**	**379,751**	+0.00	1.049
n20-b3	Max	2,668	2,398	**2,381**	**2,381**	**2,381**	+0.71	1.046
	Range	2,518	**1,640**	1,642	1,642	1,642	-0.10	1.044
	GMD	918	**673**	**673**	**673**	**673**	+0.00	1.046
	MAD	644	**490**	**490**	**490**	**490**	+0.00	1.046
	Var	757,430	**408,393**	**408,393**	**408,393**	**408,393**	+0.00	1.046
n20-b4	Max	1,981	1,819	**1,814**	**1,814**	1,815	+0.30	1.050
	Range	1,472	1,235	**1,216**	**1,216**	**1,216**	+1.51	1.047
	GMD	617	**485**	**485**	**485**	**485**	+0.00	1.048
	MAD	477	**341**	**341**	**341**	**341**	+0.00	1.048
	Var	332,502	208,876	208,876	208,876	**208,721**	+0.00	1.045
n25-b0	Max	2,607	2,610	2,613	**2,589**	**2,589**	+0.81	1.019
	Range	2,379	**1,755**	**1,755**	**1,755**	**1,755**	+0.00	1.050
	GMD	858	**683**	**683**	**683**	**683**	+0.00	1.050
	MAD	597	**503**	**503**	**503**	**503**	+0.00	1.050
	Var	718,676	**416,021**	**416,021**	**416,021**	**416,021**	+0.00	1.050
n25-b1	Max	2,145	1,971	1,938	**1,931**	**1,931**	+2.05	1.027
	Range	1,257	774	745	**738**	**738**	+4.66	1.027
	GMD	535	290	288	**285**	**285**	+1.69	1.027
	MAD	417	212	211	**209**	**209**	+1.22	1.027
	Var	255,701	82,851	81,533	**80,430**	**80,430**	+2.92	1.027
n25-b2	Max	2,148	2,148	2,147	**2,145**	**2,145**	+0.13	1.029
	Range	2,036	1,815	**1,693**	**1,693**	**1,693**	+6.71	1.046
	GMD	763	**654**	**654**	**654**	689	+0.00	1.046
	MAD	546	**479**	504	498	**479**	-3.99	1.040
	Var	516,255	418,916	**372,661**	418,916	417,741	+0.00	1.036
n25-b3	Max	1,984	1,968	1,958	**1,953**	**1,953**	+0.72	1.045
	Range	1,474	1,162	1,058	**1,056**	**1,056**	+9.13	1.045
	GMD	560	**419**	423	**419**	**419**	+0.00	1.046
	MAD	405	314	312	**311**	**311**	+1.10	1.050
	Var	283,090	169,897	**160,109**	**160,109**	**160,109**	+5.76	1.046
n25-b4	Max	2,151	2,151	2,151	**2,150**	**2,150**	+0.03	1.044
	Range	1,931	**1,433**	1,544	**1,433**	1,544	+0.00	1.034
	GMD	683	**565**	**565**	**565**	**565**	+0.00	1.044
	MAD	468	**397**	**397**	**397**	**397**	+0.00	1.044
	Var	459,013	**294,618**	**294,618**	**294,618**	**294,618**	+0.00	1.044
n30-b0	Max	2,773	2,606	2,602	**2,597**	2,597	+0.36	1.044
	Range	2,626	2,388	2,393	2,368	**2,357**	+0.86	1.044
	GMD	1,000	**760**	831	820	821	-7.91	1.044
	MAD	724	566	551	552	**548**	+2.54	1.044
	Var	911,746	516,782	516,782	516,782	**515,564**	+0.00	1.048
n30-b1	Max	2,657	2,552	2,530	**2,524**	2,526	+1.13	1.037
	Range	2,287	1,903	1,893	**1,892**	1,893	+0.61	1.041
	GMD	913	759	761	758	**706**	+0.16	1.044
	MAD	684	535	516	**516**	517	+3.54	1.049
	Var	730,545	**460,952**	**460,952**	**460,952**	**460,952**	+0.00	1.044

Instance	Measure	Cost	Det	SAA	S-0	S-1	S-0 vs Det (%)	Cost ratio S-0
n30-b2	Max	1,957	**1,942**	1,958	1,950	1,949	-0.39	1.034
	Range	1,838	1,224	1,217	1,205	**1,204**	+1.53	1.049
	GMD	667	508	514	508	**508**	+0.00	1.048
	MAD	466	393	393	**392**	394	+0.28	1.049
	Var	419,940	225,689	230,414	225,689	**225,408**	+0.00	1.048
n30-b3	Max	2,401	2,387	2,306	**2,300**	**2,300**	+3.67	1.042
	Range	2,371	1,946	**1,878**	**1,878**	**1,878**	+3.51	1.050
	GMD	853	664	**661**	**661**	**661**	+0.39	1.050
	MAD	592	453	453	**451**	**451**	+0.32	1.050
	Var	717,888	449,606	**447,388**	**447,388**	**447,388**	+0.49	1.050
n30-b4	Max	2,325	2,244	2,200	**2,193**	**2,193**	+2.28	1.045
	Range	2,100	1,739	1,701	**1,695**	**1,695**	+2.53	1.043
	GMD	762	632	632	**629**	630	+0.51	1.050
	MAD	533	457	**433**	**433**	**433**	+5.16	1.048
	Var	536,206	354,473	364,184	**352,875**	**352,875**	+0.45	1.046
n35-b0	Max	2,892	2,717	2,703	2,697	**2,694**	+0.75	1.035
	Range	2,744	2,488	2,486	2,468	**2,447**	+0.80	1.035
	GMD	1,013	846	860	844	**841**	+0.24	1.028
	MAD	718	580	574	564	**562**	+2.88	1.028
	Var	964,248	809,215	800,421	798,285	**793,815**	+1.35	1.022
n35-b1	Max	2,543	2,245	**2,137**	2,233	**2,137**	+0.52	1.045
	Range	2,429	1,793	1,768	1,767	**1,239**	+1.48	1.049
	GMD	946	502	502	501	**499**	+0.22	1.050
	MAD	696	**374**	**374**	**374**	**374**	+0.00	1.050
	Var	758,306	228,342	230,146	228,342	**228,317**	+0.00	1.050
n35-b2	Max	1,922	**1,748**	**1,748**	**1,748**	**1,748**	+0.00	1.047
	Range	1,526	**620**	**620**	**620**	**620**	+0.00	1.047
	GMD	562	**246**	**246**	**246**	**246**	+0.00	1.047
	MAD	398	**184**	**184**	**184**	**184**	+0.00	1.047
	Var	296,372	**58,746**	**58,746**	**58,746**	**58,746**	+0.00	1.047
n35-b3	Max	2,527	2,453	2,420	**2,420**	2,423	+1.36	1.037
	Range	2,137	**1,292**	**1,292**	**1,292**	**1,292**	+0.00	1.050
	GMD	780	532	**524**	**524**	**524**	+1.42	1.050
	MAD	548	440	**397**	**397**	**397**	+9.77	1.050
	Var	555,805	**247,798**	**247,798**	**247,798**	**247,798**	+0.00	1.050
n35-b4	Max	2,510	**2,153**	2,174	**2,153**	**2,153**	+0.00	1.025
	Range	2,436	1,026	**1,022**	**1,022**	**1,022**	+0.38	1.050
	GMD	902	415	**411**	**411**	**411**	+0.96	1.050
	MAD	640	290	**290**	**290**	**290**	+0.24	1.048
	Var	772,452	**156,368**	158,633	**156,368**	158,899	+0.00	1.050

Table 18: E7b, $n = 100$: mean (standard deviation) over three solver seeds of the ex-post value.

Instance	Measure	Det	SAA	S-0	S-0 vs Det (%)
n100-b0	GMD	378 (53.9)	349 (11.7)	337 (3.7)	+10.77
n100-b0	Max	3,016 (137)	2,930 (6.2)	2,924 (6.3)	+3.06
n100-b0	Range	870 (1.0)	861 (12.9)	840 (0.3)	+3.44
n100-b1	GMD	249 (30.4)	231 (1.6)	235 (4.6)	+5.71
n100-b1	Max	2,727 (9.7)	2,665 (2.7)	2,667 (0.3)	+2.20

Instance	Measure	Det	SAA	S-0	S-0 vs Det (%)
n100-b1	Range	604 (3.5)	648 (27.8)	610 (8.6)	-1.05
n100-b2	GMD	346 (17.6)	351 (6.7)	401 (39.8)	-15.87
n100-b2	Max	2,444 (8.5)	2,493 (32.0)	2,484 (62.5)	-1.65
n100-b2	Range	1,081 (318)	929 (44.4)	970 (5.4)	+10.26
n100-b3	GMD	304 (10.3)	313 (14.9)	305 (7.8)	-0.20
n100-b3	Max	2,583 (51.7)	2,535 (0.2)	2,602 (117)	-0.72
n100-b3	Range	881 (10.6)	785 (7.3)	799 (9.8)	+9.35
n100-b4	GMD	516 (5.5)	523 (19.8)	502 (11.3)	+2.79
n100-b4	Max	2,525 (8.6)	2,497 (12.4)	2,475 (15.8)	+2.02
n100-b4	Range	1,346 (64.1)	1,365 (63.8)	1,335 (50.6)	+0.83
n100-b5	GMD	289 (6.6)	273 (4.1)	243 (22.8)	+15.89
n100-b5	Max	2,396 (12.8)	2,431 (62.3)	2,391 (8.5)	+0.23
n100-b5	Range	890 (282)	700 (85.3)	630 (69.9)	+29.24