

Problem B. Good Sequence

Time Limit 2000 ms

Problem Statement

You are given a sequence of positive integers of length N , $a = (a_1, a_2, \dots, a_N)$. Your objective is to remove some of the elements in a so that a will be a **good sequence**.

Here, an sequence b is a **good sequence** when the following condition holds true:

- For each element x in b , the value x occurs exactly x times in b .

For example, $(3, 3, 3)$, $(4, 2, 4, 1, 4, 2, 4)$ and $()$ (an empty sequence) are good sequences, while $(3, 3, 3, 3)$ and $(2, 4, 1, 4, 2)$ are not.

Find the minimum number of elements that needs to be removed so that a will be a good sequence.

Constraints

- $1 \leq N \leq 10^5$
- a_i is an integer.
- $1 \leq a_i \leq 10^9$

Input

Input is given from Standard Input in the following format:

```
N
a1 a2 ... aN
```

Output

Print the minimum number of elements that needs to be removed so that a will be a good sequence.

Sample 1

Input	Output
4 3 3 3 3	1

We can, for example, remove one occurrence of 3. Then, (3, 3, 3) is a good sequence.

Sample 2

Input	Output
5 2 4 1 4 2	2

We can, for example, remove two occurrences of 4. Then, (2, 1, 2) is a good sequence.

Sample 3

Input	Output
6 1 2 2 3 3 3	0

Sample 4

Input	Output
1 1000000000	1

Remove one occurrence of 10^9 . Then, () is a good sequence.

Sample 5

Input	Output
8 2 7 1 8 2 8 1 8	5