

**Unexpected Turnaround: Cushing's Syndrome Causing New Weight Gain After
Bariatric Surgery**

Tatiana Tatranska, MD^a

Pia Strele-Trieb, MD^b

Clemens G. Wiesinger, MD^c

Martin Windpessl, MD^{d*}

^aDepartment of Internal Medicine I, Gastroenterology and Hepatology, Klinikum Wels-Grieskirchen, Wels, Austria

^bInstitute of Nuclear Medicine, Klinikum Wels-Grieskirchen, Wels, Austria

^cDepartment of Urology, Klinikum Wels-Grieskirchen, Wels, Austria

^dDepartment of Internal Medicine IV, Nephrology, Klinikum Wels-Grieskirchen, Wels, Austria

* Corresponding author

Dr. Martin Windpessl

Telephone: +43 (0)724241592194

Fax: +43 (0)72424153993

E-mail: martin.windpessl@klinikum-wegr.at

Funding source: None

The authors declare that they have no conflict of interests.

All authors had access to the data and a role in writing the manuscript.

Article type: Clinical Communication to the Editor

Key words: Cushings's syndrome, bariatric surgery, hypercortisolism

Running head: Cushing's syndrome after bariatric surgery

Unexpected Turnaround: Cushing's Syndrome New Recurrent Weight Gain After Bariatric Surgery

To the editor:

A 47-year-old woman with a history of bariatric surgery was admitted when an outpatient gastroscopy, performed because of epigastric pain, disclosed a deep marginal ulcer. There was no evidence of perforation on computed tomography imaging; incidentally, a right-sided adrenal adenoma was noted, measuring 33 mm in diameter with 10 Hounsfield units.

As the patient appeared cushingoid, a detailed history was obtained. Three years earlier, she had undergone Roux-en-Y gastric bypass surgery for class III obesity (weight at the time of surgery: 114 kg, corresponding to a body mass index (BMI) of $> 40 \text{ kg/m}^2$). She quickly lost weight thereafter, reaching a nadir of 77 kg 18 months later.

Subsequently, weight increased insidiously, despite no change in diet habits. During the following months, the patient developed hypertension, soon requiring triple therapy. In parallel, she noticed impure skin, xanthelasma, and frontal alopecia.

On present admission, the patient weighed 101 kg. She appeared plethoric, bilateral xanthelasma and teleangiectasias were evident (Figure). There were signs of easy skin bruising. Medication consisted of candesartan, a thiazide diuretic, and nebivolol; since surgery, she received regular vitamin B12 injections. She did not take inhalers, skin ointments, eye drops, or any herbal remedies. Laboratory results were in keeping with impaired fasting glucose (HbA1c 5.9%, compared to 5.3% at the time of surgery); potassium was 3.9 mmol/l, and cholesterol 210 mg/dl; thyroid function was normal.

The plasma corticotropin (ACTH) was fully suppressed (< 1.5 pg/ml), morning cortisol levels were between 18.1 and 25.6 mcg/dl on repeat testing (reference range 4.8-19.5). 24-hour urinary free cortisol was 311 mcg/24 (< 148). A dexamethason suppression test and late-night salivary cortisol further supported the diagnosis of ACTH-independent hypercortisolism.

The patient underwent unilateral robot-assisted adrenalectomy and received postoperative stress dose steroid cover to prevent withdrawal syndrome. Synacthen, administered on the second postoperative day, did not mount an adequate cortisol response (5.3 and 7.3 mcg/dl at 30 and 60 minutes, respectively). She was discharged on a tapering dose of hydrocortisone. On her last appointment, six months later, she remains well and has lost 7 kg, blood pressure is now well controlled with a single agent.

Discussion

Bariatric surgery has emerged as a useful treatment for persons with severe obesity, resulting in sustained weight loss and ultimately in longer life expectancy than conventional obesity care.¹ Obesity is also present in the vast majority of patients with endogenous hypercortisolism; however, given the well known association of weight and “physiologic hypercortisolism” (pseudo-Cushing’s), the low pre-test probability (a consequence of the rarity of Cushing’s syndrome and the continuous rise in the number of obese persons), major societies do not advocate routine evaluation for hypercortisolism prior to operations for weight loss, unless “clinically suspected”.² Cushing’s syndrome following bariatric surgery is seldom discovered. A multicenter study found 16 patients over a period of 15 years, who were diagnosed with Cushing’s syndrome following such operations (median time between bariatric surgery to treatment of hypercortisolism, 2.5 years).³

The clinical suspicion of Cushing's syndrome hinges on the recognition of the typical signs and symptoms of glucocorticoid excess which may emerge more or less simultaneously, as in the present case. Dermatologic changes such as fragile skin, ecchymoses, and teleangiectasias, are valuable in the differentiation between garden variety metabolic syndrome associated with obesity and genuine hypercortisolism.

References:

1. Carlsson LMS, Sjöholm K, Jacobson P, Andersson-Assarsson JC, Svensson PA, et al. Life Expectancy after Bariatric Surgery in the Swedish Obese Subjects Study. *N Engl J Med*. 2020;383:1535-1543.
2. Mechanick JJ, Youdim A, Jones DB, Garvey WT, Hurley DL, et al. The Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Obesity* (Silver Spring). 2013 Mar;21 Suppl 1(0 1):S1-27.
3. Javorsky BR, Carroll TB, Tritos NA, Salvatori R, Heaney AP, et al. Discovery of Cushing's Syndrome After Bariatric Surgery: Multicenter Series of 16 Patients. *Obes Surg*. 2015;25:2306-13.

Figure legend

Xanthelasma were evident at presentation. Six months post adrenalectomy, they had almost completely disappeared.

