

**Solutions to Exercise Set 6.**

12.2.(a) By symmetry,  $z_j = z_{N+1-j}$ , so  $\bar{z}_N = 0$ . Then using the Riemann approximation to an integral,  $\sigma_z^2 = (1/N) \sum z_j^2 = \sum \Phi^{-1}(j/(N+1))^2 (1/N) \rightarrow \int_0^1 \Phi^{-1}(x)^2 dx = \int_{-\infty}^{\infty} u^2 d\Phi(u) = 1$ .

(b) As in Example 1,  $\max(a_j - \bar{a})^2$  is bounded and  $\sum (a_j - \bar{a})^2 = m(N-m)/N$ . Therefore  $\delta_N$  goes to 0 if  $\max z_j^2/N \rightarrow 0$ . But  $\max z_j^2 = z_N^2 = \Phi^{-1}(N/(N+1))^2$ . Using the approximation  $1 - \Phi(x) \approx \phi(x)/x$  for large values of  $x$ , we find  $z_N^2 < 2 \log(N+1)$  for large  $N$ . Thus  $\delta_N \rightarrow 0$  and Theorem 12 gives  $S_N/\sqrt{\text{Var}(S_N)} \xrightarrow{\mathcal{L}} \mathcal{N}(0,1)$ , where  $\text{Var}(S_N) = N^2 \sigma_z^2 \sigma_a^2 / (N-1) \approx Nr(1-r)$ . Unlike Exercise 12.1(b), no further conditions are needed to conclude that  $S_N/\sqrt{N} \xrightarrow{\mathcal{L}} \mathcal{N}(0, r(1-r))$ .

12.6. (a)  $a(i) = b_1$  for  $i = 1, \dots, n$ ,  $a(i) = b_2$  for  $i = n+1, \dots, 2n$ , and  $a(i) = 0$  for  $i = 2n+1, \dots, 3n = N$ .

(b)  $\bar{a}_N = (b_1 + b_2)/3$ , and  $(1/N) \sum a(i)^2 = (b_1^2 + b_2^2)/3$ , so  $\sigma_a^2 = (b_1^2 + b_2^2)/3 - (b_1 + b_2)^2/9 = 2(b_1^2 - b_1 b_2 + b_2^2)/9$ . So  $\text{ES}_N = N \bar{z}_n \bar{a}_N = (N(N+1)/2)((b_1 + b_2)/3)$ , and  $\text{Var}(S_N) = (N^2/(N-1)) \sigma_z^2 \sigma_a^2 = (N^2(N+1)/12) \sigma_a^2$ . Since  $\max_j (a(j) - \bar{a}_N)^2 / \sigma_a^2$  is constant and  $\max_j (j - ((n+1)/2))^2 / \sigma_z^2 = [(N-1)^2/2] / [(N^2-1)/12] \rightarrow 6$ , we have

$$\frac{1}{N} \cdot \frac{\max_j (z_j - \bar{z}_N)^2}{\sigma_z^2} \cdot \frac{\max_j (a(j) - \bar{a}_N)^2}{\sigma_a^2} \rightarrow 0.$$

Hence,  $(S_N - \text{ES}_N)/\sqrt{\text{Var}(S_N)} \xrightarrow{\mathcal{L}} \mathcal{N}(0,1)$ . This reduces to  $\sqrt{N}((S_N/N^2) - (b_1 + b_2)/6) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma_a^2/12) = \mathcal{N}(0, (b_1^2 - b_1 b_2 + b_2^2)/54)$ .

(c) Thus for all  $(b_1, b_2)$ ,

$$\sqrt{N} \begin{pmatrix} b_1 & b_2 \end{pmatrix} \left( \begin{pmatrix} T_N/N^2 \\ V_N/N^2 \end{pmatrix} - \begin{pmatrix} 1/6 \\ 1/6 \end{pmatrix} \right) \xrightarrow{\mathcal{L}} \mathcal{N}\left(0, \frac{1}{54} \begin{pmatrix} b_1 & b_2 \end{pmatrix} \begin{pmatrix} 1 & -1/2 \\ -1/2 & 1 \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}\right)$$

This implies by the Cramér-Wold device,

$$\sqrt{N} \left( \begin{pmatrix} T_N/N^2 \\ V_N/N^2 \end{pmatrix} - \begin{pmatrix} 1/6 \\ 1/6 \end{pmatrix} \right) \xrightarrow{\mathcal{L}} \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \frac{1}{54} \begin{pmatrix} 1 & -1/2 \\ -1/2 & 1 \end{pmatrix}\right).$$

13.5.(a) Solving  $F(x_p|\theta) = p$ , we find  $x_p = p\theta/(1-p)$ . Therefore,

$$\sqrt{n} \left( X_{[np]} - \frac{p\theta}{(1-p)} \right) \xrightarrow{\mathcal{L}} \mathcal{N}\left(0, \frac{p(1-p)}{f(x_p|\theta)^2}\right) = \mathcal{N}\left(0, \frac{p\theta^2}{(1-p)^3}\right).$$

(b) For  $g(x) = (1-p)x/p$  and  $\hat{\theta}_n = (1-p)X_{[np]}/p$ , we have  $g'(x) = (1-p)/p$  and

$$\sqrt{n}(\hat{\theta}_n - \theta) \xrightarrow{\mathcal{L}} \mathcal{N}\left(0, \frac{p\theta^2(1-p)^2/p^2}{(1-p)^3}\right) = \mathcal{N}\left(0, \frac{\theta^2}{p(1-p)}\right).$$

The asymptotic variance is minimized by choosing  $p$  to maximize  $p(1-p)$ . This gives  $p = 1/2$ , so the median is the best quantile to use to estimate  $\theta$  (if you are only going to use one quantile).