Question 1 | Correct Mark 1.00 out of 1.00

Given an array `nums` of size `n`, return *the majority element*.

The majority element is the element that appears more than `⌊n / 2⌋` times. You may assume that the majority element always exists in the array.

Example 1:

```
Input: nums = [3,2,3]
Output: 3
```

Example 2:

```
Input: nums = [2,2,1,1,1,2,2]
Output: 2
```

Constraints:

- `n == nums.length`
- `1 <= n <= 5 * 10^4`
- `-2^31 <= nums[i] <= 2^31 - 1`

For example:

Input	Result
3 3 2 3	3
7 2 2 1 1 1 2 2	2

Answer: (penalty regime: 0 %)

```c
#include <stdio.h>

int main()
{
    int n;
    scanf("%d", &n);

    long long num, candidate = 0;
    int count = 0;

    for (int i = 0; i < n; i++)
    {
        scanf("%lld", &num);

        if (count == 0)
        {
            candidate = num;
            count = 1;
        }
        else if (num == candidate)
        {
            count++;
        }
        else
        {
            count--;
```

```
        }
    }

    printf("%lld", candidate);

    return 0;
}
```

	Input	Expected	Got	
✔	3 3 2 3	3	3	✔

Passed all tests! ✔

Correct

Marks for this submission: 1.00/1.00.

Back to Course