

STATISTICS WORKSHEET-3

Q1 to Q9 have only one correct answer. Choose the correct option to answer your question.

1. Which of the following is the correct formula for total variation?
 - a) Total Variation = Residual Variation – Regression Variation
 - b) Total Variation = Residual Variation + Regression Variation**
 - c) Total Variation = Residual Variation * Regression Variation
 - d) All of the mentioned
2. Collection of exchangeable binary outcomes for the same covariate data are called _____ outcomes.
 - a) random
 - b) direct
 - c) binomial**
 - d) none of the mentioned
3. How many outcomes are possible with Bernoulli trial?
 - a) 2**
 - b) 3
 - c) 4
 - d) None of the mentioned
4. If H_0 is true and we reject it is called **a) Type-I error**
 - b) Type-II error
 - c) Standard error
 - d) Sampling error
5. Level of significance is also called:
 - a) Power of the test
 - b) Size of the test**
 - c) Level of confidence
 - d) Confidence coefficient
6. The chance of rejecting a true hypothesis decreases when sample size is:
 - a) Decrease
 - b) Increase**
 - c) Both of them
 - d) None
7. Which of the following testing is concerned with making decisions using data?
 - a) Probability
 - b) Hypothesis**
 - c) Causal
 - d) None of the mentioned
8. What is the purpose of multiple testing in statistical inference?
 - a) Minimize errors
 - b) Minimize false positives
 - c) Minimize false negatives
 - d) All of the mentioned**

9. Normalized data are centred at ____ and have units equal to standard deviations of the original data

- a) 0
- b) 5
- c) 1
- d) 10

Q10 and Q15 are subjective answer type questions, Answer them in your own words briefly.

10. What Is Bayes' Theorem?

Answers:- **The probability of occurrence of an event related to any condition.** It is also considered for the case of conditional probability. Bayes theorem is also known as the formula for the probability of causes.

11. What is z-score?

Answers:- Number of standard deviation from the mean. The z-score can be calculated by subtracting mean by test value and dividing it by standard value. Where x is the test value, μ is the mean and σ is the standard value.

12. What is t-test?

Answers:- The t-test is any statistical hypothesis test in which the test statistic follows a Student's t-distribution under the null hypothesis.

13. What is percentile?

Answer:- A value on a scale of one hundred that indicates the percent of a distribution that is equal to or below it. a percentile score of 95 is a score equal to or better than 95 percent of the scores.

14. What is ANOVA?

Answer:- Analysis of variance (ANOVA) is a collection of statistical models and their associated estimation procedures (such as the "variation" among and between groups) used to analyze the differences among means.

15. How can ANOVA help?

Answer:- ANOVA is helpful for testing three or more variables. It is similar to multiple two-sample t-tests. However, it results in fewer type I errors and is appropriate for a range of issues. ANOVA groups differences by comparing the means of each group and includes spreading out the variance into diverse sources.

