

# Sun's analemma viewed from earth

Vincent Degrooff

July 24, 2025

Everything is vague to a degree you do not realize till you have tried to make it precise

---

Bertrand Russell,  
*The Philosophy of Logical Atomism*

What is noon? This trivial question hides a reality unsuspected at first glance. Any sensible person would tell you that noon is when their clock shows 12:00. Someone a bit more erudite could answer that noon is when the sun reaches its highest point in the sky. Are these two definitions equivalent?

Well, obviously, with the advent of time zones, the answer is no. But even at Greenwich, located right on the meridian, the two are close but do not match! Sometimes by 15 minutes! This is known at least since Ptolemy, the Greek astronomer, mathematician and geographer who lived in the 2nd century.


Let us look in figure 1 at the sunrise and sunset times on the top of *Monte Perdido* in the Pyrenees, located at 42°40'N, 00°02'E, very close to the meridian. Noon is computed as the average of sunrise and sunset times, and as expected, it is near 13:00 in winter (CET+1) and near 14:00 in summer (CEST+2). But if we zoom in on the data, we see that noon is not constant! It varies by more than 30 minutes over the year. And this is independent of the latitude!

Defining day length as the time between consecutive noons, we would find that it is not constant either and by extension, the hours, minutes and seconds would change from day to day... This is the definition of *apparent solar time*, which is the time measured by a sundial. Low-tech, but not very practical in our organized society.

Instead, we prefer to use the *mean solar day* which matches the sun's average rate of motion over the year. It is simply defined as 86 400 seconds, give or take a few cosmic crumbs that are not our business today.

Then, the natural question is: where is the sun at 12:00 if it is not at its highest point? We will answer this question by digging into the math and physics at play, and discover the beautiful figure drawn by the sun: the analemma.

Someone actually did this experiment and recorded the sun's position throughout the year [timelapse link]. On my side, I created interactive 3D plots of the analemma on Desmos, available [here] and [here].



**Figure 1.** Sunrise and sunset times at *Monte Perdido* in local time [data from `TimeAndDate`].


The analemma depends on three parameters:

- the Earth’s axial tilt angle  $\epsilon$ ;
- the eccentricity  $e$  of the Earth’s elliptic orbit;
- the argument of periapsis  $\omega$ , relating the delay from perihelion to spring equinox.

To understand why, let’s proceed step by step, starting without any of these parameters.

We should also note that latitude and longitude only translate the analemma to a different position in the sky, but do not change its shape.

For simplicity, let’s furthermore suppose that the year is exactly  $N = 365$  solar days of  $T_d = 86400 = 24 \times 60 \times 60$  seconds. This is obviously inaccurate, but sufficient to reveal the beauty of the phenomenon at play.



## 1 The simplest case

The Earth (E) is revolving counterclockwise around the sun (S) in a circular orbit of radius  $a$ , with angular velocity  $\dot{\phi} = \Omega$ . An observer (O) is rotating counterclockwise around the Earth's axis at a distance  $R$  with angular velocity  $\dot{\theta}$ .

The night sky completes a full rotation to an Earth-based observer in just under 24 hours. The extra  $\sim 4$  minutes account for the fact that, during that time, the Earth also moves slightly along its orbit around the sun - by about  $2\pi/N$  radians. As a result, Earth must rotate a bit more for the Sun to return to the same position in the sky.

This period is called the *sidereal day*, noted here  $T_s$ . We experience  $N$  solar days in a year, but the Earth actually completes  $N + 1$  sidereal days in the same time because of its prograde rotation:


$$\dot{\phi} = \Omega = \frac{2\pi}{NT_d}, \quad \dot{\theta} = \frac{2\pi}{T_s} = (N + 1)\Omega \quad (1)$$

We want to find the position of the sun at a given time  $t$  in the Earth's frame:

$$\begin{aligned} \vec{OS} &= \vec{OE} + \vec{ES}, \\ &= -R\mathbf{y}_1 - a\mathbf{x}_1, \\ &= -R\mathbf{y}_1 - a[\cos(\phi - \theta)\mathbf{y}_1 + \sin(\phi - \theta)\mathbf{y}_2]. \end{aligned} \quad (2)$$

Since  $R/a \approx 4 \times 10^{-5}$ , we will neglect the term  $R\mathbf{y}_1$ . From here on, we will use the direction vector  $s \approx \vec{OS}/\|\vec{OS}\|$  that neglects the  $R$  term. Rewriting  $\phi - \theta$ , and expressing 12:00 on day  $k$  as  $t_k = T_d/2 + kT_d$ , we get:

$$s(t) = -[\cos(2\pi t/T_d)\mathbf{y}_1 + \sin(2\pi t/T_d)\mathbf{y}_2] \implies s(t_k) = \mathbf{y}_1. \quad (3)$$



## 2 Effect of Earth's axial tilt

The Earth's axis is tilted by an angle  $\epsilon = 23.44^\circ$  with respect to the ecliptic plane. The basis  $[y]$  attached to the Earth's frame is now obtained after two rotations: first by  $\epsilon \in [0, \frac{\pi}{2}]$  around the  $I_2$  axis, then by  $\theta$  around the third axis:

$$\begin{aligned} [y] &= R_3(\theta)R_2(\epsilon)[I] \\ [x] &= R_3(\phi)R_2(-\epsilon)R_3(-\theta)[y]. \end{aligned} \quad (4)$$

The sun's position in the Earth's frame  $[y]$  is now given by:

$$s = -x_1 = \begin{bmatrix} -\cos(\phi)\cos(\epsilon)\cos(\theta) - \sin(\phi)\sin(\theta) \\ \cos(\phi)\cos(\epsilon)\sin(\theta) - \sin(\phi)\cos(\theta) \\ -\cos(\phi)\sin(\epsilon) \end{bmatrix}. \quad (5)$$

### Where is the sun at 12:00?

Once again, we insert  $t_k = kT_d + T_d/2$  with  $k = 0, 1, 2, \dots, N-1$  in  $s$  to derive the sun's position at noon. First let's express the angles  $\phi$  and  $\theta$  in terms of  $k$ , for both prograde and retrograde rotation:

$$\phi|_{12:00} = \Omega t_k = \frac{2\pi}{N}(k + \frac{1}{2}), \quad (6)$$

$$\theta|_{12:00} = \dot{\theta} t_k = \frac{2\pi}{N}(1 \pm N)(k + \frac{1}{2}) = \pm 2\pi k \pm \pi + \phi|_{12:00}, \quad (7)$$


$$\implies \cos(\theta)|_{12:00} = -\cos(\phi), \quad (8)$$

$$\implies \sin(\theta)|_{12:00} = -\sin(\phi). \quad (9)$$

With these, we can express the analemma as a parametric curve with parameter  $\mu \in [0, 2\pi)$ :

$$s|_{12:00}(\mu) = \frac{1}{2} \begin{bmatrix} 1 + \cos \epsilon \\ 0 \\ 0 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} -(1 - \cos \epsilon) \cos(2\mu) \\ (1 - \cos \epsilon) \sin(2\mu) \\ -2 \sin(\epsilon) \cos(\mu) \end{bmatrix}. \quad (10)$$

Based on equation (10), we can note that the analemma is the same for prograde and retrograde rotations. This curve is shown in figure 2 for various tilt angles  $\epsilon$ , in the observer's frame, that is, without the  $y_1$  component.



**Figure 2.** The analemma of 12:00, viewed from the ground at 00°N. The border of the figure is the horizon of the observer. Each month has been assigned a specific color.

### When is the sun at its highest point?


At solar noon and midnight, the vector  $\mathbf{x}_1$  has no  $y_2$  component: :

$$\begin{aligned} \cos(\epsilon) \cos(\Omega t) \sin((1 \pm N)\Omega t) &= \sin(\Omega t) \cos((1 \pm N)\Omega t) \\ \iff \pm \sin(N\Omega t)(1 - c \cos^2(\Omega t)) &= \frac{c}{2} \cos(N\Omega t) \sin(2\Omega t) \end{aligned} \quad (11)$$

where  $c = 1 - \cos(\epsilon)$ . This equation has no closed form solution for  $t^*$ , but we can approximate it around 12:00 with  $t^* = kT_d + T_d/2 + \delta_k = t_k + \delta_k$ ,  $k = 0, 1, 2 \dots N - 1$ :

$$\left\{ \begin{array}{l} \sin(N\Omega t^*) = -\sin(2\pi\delta_k/T_d) \\ \cos(N\Omega t^*) = -\cos(2\pi\delta_k/T_d) \\ \sin(2\Omega t^*) \approx \sin(2\Omega t_k) \\ \cos(\Omega t^*) \approx \cos(\Omega t_k) \end{array} \right\} \implies \frac{2\pi\delta_k}{T_d} \approx \tan\left(\frac{2\pi\delta_k}{T_d}\right) = \pm \frac{\frac{c}{2} \sin\left(2 \cdot 2\pi \frac{k+0.5}{N}\right)}{1 - c \cos^2\left(2\pi \frac{k+0.5}{N}\right)}. \quad (12)$$

Noon is off by  $\delta$  seconds from the mean solar time, and this shift oscillates twice a year as we can see with the factor at the numerator of (12) and in figure 3. The approximation can be simplified with  $\tan(x) \approx x$  ("approx. 2") and even further by removing the denominator ("approx. 3"), providing acceptable results until quite large tilt angles  $\epsilon$ .




**Figure 3.** Apparent noon time throughout the year for two different axial tilt angles  $\epsilon$ .

### An intuitive explanation

If we consider the Earth's rotating frame, the sun is revolving around the earth, but also moving up (winter and spring) and down (autumn and summer) its equatorial plane.

This vertical motion has a period of one year: the sun reaches its highest point at the summer solstice, and its lowest point at the winter solstice, for people in the northern hemisphere. Since the sun's total motion is constant in this circular orbit model, its horizontal motion must slow down when the vertical motion is at its highest *in absolute value*.

The sun's eastward velocity is thus at its highest at both solstices, and at its lowest at both equinoxes, producing a motion of period half a year. This eastward velocity can be observed in 3 as the slope of the apparent noon time. In fact, if the sun is moving faster to the East than the average pace, it will take more time to reach the zenith, shifting noon a bit later than the day before.



### 3 Effect of Earth's orbit eccentricity

The Earth's orbit is not a perfect circle, but an ellipse with semi-major axis  $a$  and semi-minor axis  $b$ . For the earth, the orbit eccentricity  $e = \sqrt{1 - b^2/a^2} \approx 0.017$  is relatively small, but enough to have a noticeable effect on the analemma. The eccentricity  $e \approx 0.66$  in the figure above is exaggerated for clarity.

In the early 17th century, Johannes Kepler stated his famous *laws of planetary motion*:

1. The orbit of a planet is an ellipse with the sun at one focus.
2. A line segment joining a planet and the sun sweeps out equal areas during equal intervals of time.
3. The square of the orbital period of a planet is proportional to the cube of the semi-major axis of its orbit.

#### Anomalies

To describe efficiently a planet's position along its elliptical orbit at a given time, astronomers use three angular parameters called *anomalies* to stress the difference with a circular orbit.

1. *True anomaly*  $\nu$ , the angle between the direction of perihelion and the Earth.
2. *Eccentric anomaly*  $E$ , an auxiliary angle used to simplify calculations.
3. *Mean anomaly*  $M = \Omega t$ , increasing linearly with time.

While  $M(t)$  increases linearly, Kepler's second law implies that  $\nu(t)$  does not: the Earth moves faster near perihelion and slower near aphelion. This second law states that the blue

and orange areas in the figure above are equal:

$$|AOP''| = |ASP'| = |AOP'| - |OSP'|, \quad (13)$$

$$\frac{a^2 M}{2} = \frac{a^2 E}{2} - \frac{(ae)(a \sin e)}{2}, \quad (14)$$

$$M = E - e \sin E. \quad (15)$$

The resulting equation (15) is known as *Kepler's equation* and has no closed form solution for  $E$  in terms of  $M$ . Once the eccentric anomaly  $E$  has been derived from the mean anomaly  $M$ , it remains to find the true anomaly  $\nu$ :

$$r = a(1 - e \cos E), \quad (16)$$

$$\cos \nu = \frac{\cos E - e}{1 - e \cos E}, \quad (17)$$

$$\sin \nu = \frac{\sqrt{1 - e^2} \sin E}{1 - e \cos E}, \quad (18)$$

$$r = a \frac{1 - e^2}{1 + e \cos \nu}. \quad (19)$$

The relationships (16), (17) and (18) were respectively derived by:

$$\begin{aligned} r^2 &= \|SD\|^2 + \|DP\|^2 = b^2 \sin^2 E + (ae - a \cos E)^2 \\ &= a^2 [(1 - e^2)(1 - \cos^2 E) + (e^2 - 2e \cos E + \cos^2 E)] \\ &= a^2 [1 - 2e \cos E + e^2 \cos^2 E] = a^2 (1 - e \cos E)^2, \end{aligned} \quad (20)$$

$$\begin{aligned} \|OD\| + \|DS\| &= \|OS\| \\ a \cos E - r \cos \nu &= ae \\ a \cos E - a(1 - e \cos E) \cos \nu &= ae, \end{aligned} \quad (21)$$

$$\begin{aligned} \|DP\| &= r \sin \nu = b \sin E \\ a(1 - e \cos E) \sin \nu &= a\sqrt{1 - e^2} \sin E. \end{aligned} \quad (22)$$

The true anomaly can be approximated from the mean anomaly via a Fourier expansion, but to keep the calculations tractable, we will limit ourselves to a first-order expansion:

$$\nu = M + (2e - \frac{1}{4}e^3) \sin(M) + \frac{5}{4}e^2 \sin(2M) + \frac{13}{12}e^3 \sin(3M) + \mathcal{O}(e^4), \quad (23)$$

$$= M + 2e \sin M + \mathcal{O}(e^2). \quad (24)$$

Perihelion ( $\nu = 0$ ) and winter solstice do not have to coincide, even though they are quite close currently for Earth which gets closest to the sun in early january. The delay between both events is fixed by introducing the angle  $\gamma$  between perihelion and the winter solstice.

Note also that the planet is assumed to be at perihelion at  $t = 0$ :

$$\nu(0) = M(0) = \Omega \cdot 0 = 0. \quad (25)$$



## Planetocentric longitude of perihelion

The planetocentric longitude of perihelion  $\lambda_p$  is defined as the angle between the direction of the planet's vernal equinox and the direction of its perihelion. Not directly available, this information can however be derived from first hand astronomical observations. The conventional reference frame of this data, is the equatorial coordinate system I, whose 1st-axis points in the direction Earth-Sun-Aries constellation and 3rd-axis points towards the Earth's north celestial pole. The parameters needed are listed below:

- The longitude of ascending node  $\Omega$ , measuring the angle in the ecliptic plane between reference  $x$ -axis and the planet's ascending node, i.e. where it crosses the ecliptic from south to north;
- The longitude of perihelion  $\varpi = \omega + \Omega$ , a composed angle where  $\omega$  is the argument of perihelion, measuring the angle in the planet's orbital plane between the ascending node and the perihelion;
- The planet's orbit inclination  $i$ , the angle between the ecliptic and the planet's orbital plane;
- The planet's axis orientation with respect to the reference frame, with its right ascension  $\alpha$  and declination  $\delta$ .
- The Earth's axial tilt, or obliquity  $\epsilon_0$ , the angle between the ecliptic and the Earth's equatorial plane.

The sun-planet position vector  $p$  at perihelion and the normal to the ecliptic plane  $n$  determine a basis  $(p, q, n)$ . This basis and the planet's rotation axis can be obtained through the following rotations:

$$a = [1 \ 0 \ 0] R_2(-\delta) R_3(\alpha) [I], \quad (26)$$

$$(p, q, n) = R_3(\lambda_p) R_3(\omega) R_1(i) R_3(\Omega) R_1(\epsilon_0) [I], \quad (27)$$

where only  $\lambda_p$  is unknown. The dot product  $n \cdot a$  provides the planet's obliquity  $\epsilon$ . The spring equinox is reached when the planet's rotation axis is orthogonal to its position vector, i.e.  $p \cdot a = 0$ , and oriented such that  $q \cdot a < 0$ . This allows us to find the planetocentric longitude of perihelion  $\lambda_p$ :

$$v \triangleq R_3(\omega) R_1(i) R_3(\Omega) R_1(\epsilon_0) R_3(-\alpha) R_2(\delta) \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad (28)$$

$$\begin{bmatrix} \cos \lambda_p & \sin \lambda_p \\ -\sin \lambda_p & \cos \lambda_p \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \stackrel{!}{=} \begin{bmatrix} 0 \\ -\sqrt{v_1^2 + v_2^2} \end{bmatrix} = \begin{bmatrix} 0 \\ -w \end{bmatrix}, \quad (29)$$

$$\implies \cos \lambda_p = -\frac{v_2}{w}, \quad \sin \lambda_p = \frac{v_1}{w}. \quad (30)$$

With  $\lambda_p$  now determined, the angle  $\gamma$  follows immediately:  $\gamma = \frac{\pi}{2} - \lambda$ . This closes the construction by linking the solstitial direction to the orbital geometry.

### Where is the sun at 12:00?

With the elliptical orbit, the basis  $[y]$  must now account for the position of winter solstice before the Earth's axis obliquity is applied, while the  $[x]$  basis remains unchanged.

$$[y] = R_3(\theta)R_2(\epsilon)R_3(-\gamma)[I], \quad (31)$$

$$[x] = R_3(\nu)[I] \quad (32)$$

To account for the rotation  $\gamma$ , we set an initial angle  $\theta_0$ . Even though it does not align  $y_1$  with  $x_1$  at  $t = 0$ ,  $\theta_0 = \pm\gamma$  is our preferred choice as it centers the analemma in the East-West direction over the year:

$$\theta(t) = \theta_0 + (1 \pm N)\Omega t = \gamma + (1 \pm N)\Omega t. \quad (33)$$

Taking into account the varying distance  $r$  of the Earth to the sun from equation (16), and the expressing  $x_1$  in the basis  $[y]$  with equations (31) and (32), we write the position of the sun in the Earth's frame as:

$$s = -\frac{r}{a}x_1 = \frac{r}{a} \begin{bmatrix} -\cos(\epsilon)\cos(\nu + \gamma)\cos(\theta) - \sin(\nu + \gamma)\sin(\theta) \\ \cos(\epsilon)\cos(\nu + \gamma)\sin(\theta) - \sin(\nu + \gamma)\cos(\theta) \\ -\sin(\epsilon)\cos(\nu + \gamma) \end{bmatrix}. \quad (34)$$

As both  $r$  and  $\nu$  depend on the eccentricity, we find the required first order approximations:

$$r \cos(\nu + \gamma) = \cos(M + \gamma) - e \left[ 2 \sin(M + \gamma) \sin M + \cos(M + \gamma) \cos M \right] + \mathcal{O}(e^2) \quad (35)$$

$$r \sin(\nu + \gamma) = \sin(M + \gamma) + e \left[ 2 \cos(M + \gamma) \sin M - \sin(M + \gamma) \cos M \right] + \mathcal{O}(e^2) \quad (36)$$

Let's first recall the expressions for noon, the Earth's angular position  $\theta$ , and let's derive a useful relation for integer multiples of the angular position  $\mu$ :

$$t_k = kT_d + \frac{T_d}{2}, \quad (37)$$

$$M_k = \Omega t_k = 2\pi \frac{k}{N} + \frac{\pi}{N}, \quad (38)$$

$$\theta_k = \gamma + (1 \pm N)M_k = \pm 2\pi k \pm \pi + M_k + \gamma. \quad (39)$$

Playing around with the trigonometric identities, we find:

$$r_k \cos(\nu_k) \cos(\theta_k) \approx -\cos^2(M + \gamma) + \frac{e}{4} \left[ 2 \cos M + 3 \cos(M + 2\gamma) - \cos(3M + 2\gamma) \right], \quad (40)$$

$$r_k \sin(\nu_k) \sin(\theta_k) \approx -\sin^2(M + \gamma) + \frac{e}{4} \left[ 2 \cos M - 3 \cos(M + 2\gamma) + \cos(3M + 2\gamma) \right], \quad (41)$$

$$r_k \cos(\nu_k) \sin(\theta_k) \approx -\frac{1}{2} \sin(2M + 2\gamma) + \frac{e}{4} \left[ 4 \sin M + 3 \sin(M + 2\gamma) - \sin(3M + 2\gamma) \right], \quad (42)$$

$$r_k \sin(\nu_k) \cos(\theta_k) \approx -\frac{1}{2} \sin(2M + 2\gamma) - \frac{e}{4} \left[ 4 \sin M - 3 \sin(M + 2\gamma) + \sin(3M + 2\gamma) \right]. \quad (43)$$

Inserting the approximations (40)–(43) into equation (34) allows us to approximate the analemma up to first order in  $e$  with a curve parametrized by  $\mu = M + \gamma, \in [0, 2\pi]$ :

$$s|_{12:00}(\mu) = \frac{1}{2} \begin{bmatrix} 1 + \cos \epsilon \\ 0 \\ 0 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} -(1 - \cos \epsilon) \cos(2\mu) \\ (1 - \cos \epsilon) \sin(2\mu) \\ -2 \sin(\epsilon) \cos(\mu) \end{bmatrix} + \frac{e}{4} \begin{bmatrix} \delta_1 \\ \delta_2 \\ \delta_3 \end{bmatrix} + \mathcal{O}(e^2), \quad (44)$$

$$\begin{bmatrix} \delta_1 \\ \delta_2 \\ \delta_3 \end{bmatrix} = \begin{bmatrix} -2(1 + \cos \epsilon) \cos(\mu - \gamma) + (1 - \cos \epsilon) [3 \cos(\mu + \gamma) - \cos(3\mu - \gamma)] \\ 4(1 + \cos \epsilon) \sin(\mu - \gamma) - (1 - \cos \epsilon) [3 \sin(\mu + \gamma) - \sin(3\mu - \gamma)] \\ 2 \sin \epsilon [3 \cos \gamma - \cos(2\mu - \gamma)] \end{bmatrix} \quad (45)$$


Quite an expression! The terms with a factor  $(1 - \cos \epsilon)$  or  $\sin \epsilon$  only slightly modify the analemma's shape for moderate tilt angles  $\epsilon$ . The impact of  $\delta_1$  is also quite small for an observer on Earth as he only sees the analemma's projection on the celestial sphere, i.e. the second and third components of the normalized position vector  $s$ .

Let's first consider the case  $\gamma = 0$  to measure the effect of the Earth's orbit eccentricity without the complication of the perihelion shift. In that case, the vertical symmetry of the analemma is broken. The term  $e \sin(\mu - 0)$  of  $\delta_2$ , highlighted in red, tends to push the analemma towards the East during the first half of the year and towards the West during the second half. As a consequence, the analemma's southern loop of winter solstice gets widened, while the northern loop of summer solstice gets narrowed. This effect is illustrated in figures 4 and 5. The effect would be exactly the opposite if we had  $\gamma = \pi$ , that is, if perihelion happened on summer solstice.


Let's then consider the case  $\gamma = \pi/2$ . Now, the term  $e \sin(\mu - \gamma)$  becomes  $-e \cos(\mu)$ : it pushes the analemma towards the East during spring and summer, and towards the West during autumn and winter. Therefore, the analemma's northern loop of summer solstice gets shifted towards the East, while the southern loop of winter solstice gets shifted towards the West. Both loops then keep their original width.

The situation with arbitrary angles  $\gamma$  allows a smooth transition between both cases and should just be enjoyed visually in figure 6. There, we observe that the first-order approximation remains highly accurate for eccentricity values up to 0.3.


Until now, we have only considered the analemma at 12:00 on the equator, both in the mathematical model and in figures 2-6. From other latitudes, the analemma would appear the same but simply lower in the sky. Earlier in the day, the analemma would be rotated lower and eastward, while later in the day, it would be rotated lower and westward.



**Figure 4.** 12:00 Earth analemma, viewed from the ground at 00°N. Introducing the eccentricity breaks vertical symmetry, while perihelion shift  $\gamma$  breaks horizontal symmetry.



**Figure 5.** 12:00 Mars analemma, viewed from the ground at 00°N. The martian calendar is made of 24 months, grouped here in 12 pairs.



**Figure 6.** Noon analemma, viewed from the ground at  $00^\circ\text{N}$ , for a tilt angle  $\epsilon = 35^\circ$ , eccentricities  $e = 0.01, 0.05, 0.15, 0.3, 0.6, 0.9$  and perihelion shifts  $\gamma = \frac{0\pi}{6}, \frac{1\pi}{6}, \frac{2\pi}{6}, \frac{3\pi}{6}, \frac{4\pi}{6}, \frac{5\pi}{6}, \frac{6\pi}{6}$ .

### When is the sun at its highest point?

Again, at solar noon and midnight, the position vector  $\mathbf{x}_1$  has no component  $y_2$  in the East-West direction. Equation (34) gives us the condition

$$\cos \epsilon \cos(\nu + \gamma) \sin \theta - \sin(\nu + \gamma) \cos \theta = 0, \quad (46)$$

where  $\nu$  and  $\theta$  are functions of time  $t$ . As said previously, the true anomaly  $\nu$  has no closed form solution, while the rotation angle  $\theta$  is given by equation (39). This situation, complicated by the eccentricity of the orbit, has obviously no analytical solution either. Let's use our approximation of  $\nu$  in equation (24)

$$\begin{aligned} & \cos \epsilon \left( \cos(M + \gamma) - 2e \sin(M + \gamma) \sin M \right) \sin(\gamma + (1 \pm N)M) \\ & - \left( \sin(M + \gamma) + 2e \cos(M + \gamma) \sin M \right) \cos(\gamma + (1 \pm N)M) = \mathcal{O}(e^2), \end{aligned} \quad (47)$$

and expand the high frequency sine/cosine terms to reach, with  $\mu = M + \gamma$ ,

$$\begin{aligned} & \sin(N\Omega t_k) \left[ \underbrace{\pm(\cos \epsilon \cos^2 \mu + \sin^2 \mu)}_C \underbrace{\pm(1 - \cos \epsilon) \sin(\mu - \gamma) \sin 2\mu e}_D \right] \\ & = \cos(N\Omega t_k) \left[ \underbrace{\frac{1}{2}(1 - \cos \epsilon) \sin(2\mu)}_A + \underbrace{2 \sin(\mu - \gamma) (\cos^2 \mu + \cos \epsilon \sin^2 \mu)}_B e \right] + \mathcal{O}(e^2). \end{aligned} \quad (48)$$

We will use the same assumption as in equation (12) that the time  $t^*$  satisfying equation (48) is close to noon every day:  $t_k = kT_d + T_d/2 + \delta_k = M_k + \delta_k$ , with  $k = 0, 1, 2 \dots N - 1$  and  $\delta_k \ll NT_d$ . This lets us decouple the periodic terms of low and high frequencies:

$$\begin{aligned} \sin(N\Omega t_k) &= -\sin(2\pi\delta_k/T_d) & \sin(\ell\Omega t_k) &\approx \sin(\ell M_k) \\ \cos(N\Omega t_k) &= -\cos(2\pi\delta_k/T_d) & \cos(\ell\Omega t_k) &\approx \cos(\ell M_k) \end{aligned} \quad \text{for } \ell \text{ small.} \quad (49)$$

Introducing the approximations (49) in equation (48), we can now isolate the time shift  $\delta_k$ :

$$\tan\left(\frac{2\pi\delta_k}{T_d}\right) = \frac{A + eB}{C + eD} + \mathcal{O}(e^2) = \frac{A}{C} + e \frac{BC - AD}{C^2} + \mathcal{O}(e^2). \quad (50)$$

After going through some algebra, we finally find an approximation for the noon shift  $\delta_k$  for a planet with axial tilt  $\epsilon$ , orbit eccentricity  $e$  and solstice-perihelion shift  $\gamma$ , on day  $k$  from winter solstice through  $\mu = \gamma + 2\pi/N \cdot (k + \frac{1}{2})$ :

$$\tan\left(\frac{2\pi\delta_k}{T_d}\right) = \pm \frac{\frac{1}{2}(1 - \cos \epsilon) \sin(2\mu)}{1 - (1 - \cos \epsilon) \cos^2 \mu} \pm e \frac{2 \cos \epsilon \sin(\mu - \gamma)}{[1 - (1 - \cos \epsilon) \cos^2 \mu]^2} + \mathcal{O}(e^2). \quad (51)$$

Assuming that the tilt angle is moderate so that the denominator remains close to 1 and that  $\delta_k \ll T_d$ , we can derive an over-simplified approximation for the noon shift:

$$\delta_k \approx \pm \frac{T_d}{2\pi} \left[ \frac{1 - \cos \epsilon}{2} \sin(2\mu) + 2e \cos \epsilon \sin(\mu - \gamma) \right], \quad (52)$$

which is made of two harmonic terms representing the influence of:

1. the planet's axial tilt  $\epsilon$ , with a 6 month-period starting at winter solstice and an amplitude  $24 \cdot 60(1 - \cos \epsilon)/4\pi \approx 09\text{m}30\text{s}$  for Earth;
2. the planet's orbit eccentricity  $e$ , with a 12 month-period starting at perihelion and an amplitude  $24 \cdot 60 \cdot e \cos \epsilon/\pi \approx 08\text{m}10\text{s}$  for Earth.

In order to compute a numerical solution of equation (46) with Newton-Raphson, we first need to relate the mean anomaly  $M$  to the eccentric anomaly  $\nu$  by solving the Kepler equation (15), and then to the true anomaly  $\nu$  with the relations (17) and (18). We also need  $d \cos \nu / dM$  and  $d \sin \nu / dM$ :

$$\frac{dE}{dM} = \frac{1}{1 - e \cos E} = \frac{1}{r} \quad (53)$$

$$\frac{d \cos \nu}{dM} = \frac{d \cos \nu}{dE} \frac{dE}{dM} = \frac{-(1 - e^2) \sin E}{(1 - e \cos E)^3} = -\frac{(1 + e \cos \nu)^2}{(1 - e^2)^{3/2}} \sin \nu = -\zeta \sin \nu, \quad (54)$$


$$\frac{d \sin \nu}{dM} = \frac{d \sin \nu}{dE} \frac{dE}{dM} = \frac{\sqrt{1 - e^2}(\cos E - e)}{(1 - e \cos E)^3} = +\frac{(1 + e \cos \nu)^2}{(1 - e^2)^{3/2}} \cos \nu = +\zeta \cos \nu. \quad (55)$$

From here we can derive the function  $f(M)$  whose root  $M$  gives the solar noon time, along with its derivative  $f'(M)$ :


$$f(M) = \cos \epsilon [\cos \nu \cos \gamma - \sin \nu \sin \gamma] \sin (\gamma + (1 \pm N)M) - [\sin \nu \cos \gamma + \cos \nu \sin \gamma] \cos (\gamma + (1 \pm N)M), \quad (56)$$

$$f'(M) = [(1 \pm N) - \zeta \cos \epsilon] [\sin \nu \cos \gamma + \cos \nu \sin \gamma] \sin (\gamma + (1 \pm N)M) + [(1 \pm N) \cos \epsilon - \zeta] [\cos \nu \cos \gamma - \sin \nu \sin \gamma] \cos (\gamma + (1 \pm N)M). \quad (57)$$


In figure 7, we compare the numerical solution, the full approximation ("approx. 1"), the simplified approximation ("approx. 2") and the data presented earlier in figure 1. We use the Earth's parameters  $\epsilon = 23.44^\circ$ ,  $e = 0.0167$  and  $\gamma = 12.93^\circ$ .



**Figure 7.** Due to the low eccentricity, the full approximation is almost indistinguishable from the numerical solution, both matching very well the data.




**Figure 8.** On mars,  $\epsilon = 25.19^\circ$ ,  $e = 0.0934$ ,  $\gamma = 341^\circ$ , due to the higher eccentricity, solar noon is shifted significantly more than on Earth.



**Figure 9.** For a planet with a reasonable axial tilt  $\epsilon = 10^\circ$ , but a high eccentricity  $e = 0.5$  and with  $\gamma = 60^\circ$ , the noon shift gets quite large at its peak. The approximations are no longer very accurate.





**Figure 10.** Equations of time for various angles  $\gamma$ , with  $e = 0.05$  and  $\epsilon = 35^\circ$ .

## References

- [1] B. A. Archinal et al. “Report of the IAU Working Group on Cartographic Coordinates and Rotational Elements: 2009”. In: *Celestial Mechanics and Dynamical Astronomy* 109.2 (Feb. 2011), pp. 101–135. ISSN: 1572-9478. DOI: 10.1007/s10569-010-9320-4. URL: <https://doi.org/10.1007/s10569-010-9320-4>.
- [2] Jean Eudes Arlot. *Les échelles de temps*. 2017. URL: <https://perso.imcce.fr/jean-eudes-arlot/ecole-ete-aout-2017/cours/cours-le%20temps%20en%20physique%20et%20en%20astronomie.pdf> (visited on 07/23/2025).
- [3] Pierre Barbier. *The Equation of Time*. 2012. URL: <https://pbarbier.com/eqtime/eqtime.html> (visited on 07/23/2025).
- [4] Jet Propulsion Laboratory. *Approximate Positions of the Planets*. URL: [https://ssd.jpl.nasa.gov/planets/approx\\_pos.html](https://ssd.jpl.nasa.gov/planets/approx_pos.html) (visited on 07/23/2025).
- [5] Action Physics. *Equation of time*. URL: <https://www.actionphysics.uk/index.php/equation-of-time/> (visited on 07/23/2025).