

Every value is $f^*(2\sigma)$ (smallest injected fraction detected at 2σ median significance; lower is better), median over draws. Residual children indented; best test per row **bold**. '>.1' = never reached 2σ .

method	Eucl.	Maha.	MMD	SparKer
SupCon	.042	.068	.024	.023
SupCon+SIGReg	.087	.055	.022	.023
NPLM-dist	.040	.051	.025	.034
NPLM+cwSIGReg	.034	.039	.021	.025
SupCon $\rightarrow$ res	.052	.041	.025	.028
SupCon $\rightarrow$ res-nplm	.014	.031	.025	.027
SupCon $\rightarrow$ res	.047	.040	.028	.027
SupCon $\rightarrow$ res-nplm	.035	.040	.022	.024
NPLM-dist $\rightarrow$ res	.027	.036	.026	.026
NPLM-dist $\rightarrow$ res-nplm	.082	.038	.032	.026
NPLM $\rightarrow$ res	.042	.031	.025	.027
NPLM $\rightarrow$ res-nplm	>.1	.090	.027	.027

Table 1: CIFAR-100: pre-discovery (frozen) f^* , median over draws.

method	Eucl.	Eucl.(a)	Maha.	MMD	SparKer	SpK(a)
SupCon	>.1	.023	>.1	.018	.028	.024
SupCon+SIGReg	.024	.011	.044	.017	.016	.016
NPLM-dist	.033	.023	.029	.020	.022	.016
NPLM+cwSIGReg	.030	.034	.025	.016	.016	.010
SupCon $\rightarrow$ res	>.1	>.1	>.1	>.1	>.1	>.1
SupCon $\rightarrow$ res-nplm	.015	.022	.035	.013	.008	.016
SupCon $\rightarrow$ res	>.1	>.1	>.1	>.1	>.1	>.1
SupCon $\rightarrow$ res-nplm	.025	.014	.038	.016	.014	.015
NPLM-dist $\rightarrow$ res	>.1	>.1	>.1	>.1	>.1	>.1
NPLM-dist $\rightarrow$ res-nplm	>.1	>.1	>.1	>.1	>.1	>.1
NPLM $\rightarrow$ res	>.1	>.1	>.1	>.1	>.1	>.1
NPLM $\rightarrow$ res-nplm	>.1	>.1	>.1	>.1	>.1	>.1

Table 2: CIFAR-100: post-discovery f^* , derived cut (np pool). Eucl.(a)/SpK(a) = anchor-aware.

CIFAR-100 (100 classes, $b=0.01$)

method	Eucl.	Eucl.(a)	Maha.	MMD	SparKer	SpK(a)
SupCon	.027	.019	.030	.025	.024	.021
SupCon+SIGReg	.033	.025	.042	.025	.021	.021
NPLM-dist	.056	.022	.027	.022	.031	.033
NPLM+cwSIGReg	.030	.029	.035	.029	.024	.025
SupCon $\rightarrow$ res	.066	.021	.039	.028	.026	.024
SupCon $\rightarrow$ res-nplm	.025	.032	.029	.024	.021	.025
SupCon $\rightarrow$ res	.053	.021	.047	.023	.027	.025
SupCon $\rightarrow$ res-nplm	.030	.020	.029	.022	.022	.021
NPLM-dist $\rightarrow$ res	.058	.021	.051	.023	.025	.025
NPLM-dist $\rightarrow$ res-nplm	.037	.038	.022	.022	.026	.021
NPLM $\rightarrow$ res	.040	.027	.036	.028	.027	.025
NPLM $\rightarrow$ res-nplm	.031	.032	.040	.032	.026	.023

Table 3: CIFAR-100: post-discovery f^* , $s/\sqrt{b}$ cut.