Supplementary Notes for "Intermediate Precision of Single-Molecule Break-Junction Conductance: Measurement, Uncertainty Budget, and Design Rules"

Viraaj Minhas

These notes hold material moved from the paper for length. Section, table and figure numbers without the prefix S refer to the paper; every number is generated by the same analysis code and macro files as the paper's.

S1. Two excluded cells

Two excluded cells, one within-round observation. Two further single-duplex cells hold two sessions from the same round that differ only in ramp rate: 3 against 10 (differing by 0.005 log units) and 3 against 20 (by 0.255). They are one occasion and are excluded; including them gives a factor of 1.43 rather than 1.46. The 0.255 difference has two readings: ramp rate acting as an influence quantity, or variation within a round. No other cell of the 64 sessions varies ramp rate at fixed molecule, bias and gain, so the two cannot be separated here; ramp rate is therefore held fixed in every group and named in the measurand. The mapping from ramp-rate setting to retraction speed is not in the deposit.

S2. Tunnelling filter, kept fraction and yield

Between 75% and 90% of traces are kept per session; the two sessions of a headline pair differ in kept fraction by at most 0.08, and across the seven pairs the absolute kept-fraction difference is negatively associated with the absolute median difference (Spearman -0.86, exact permutation $p = 0.024$; signed differences -0.29, $p = 0.56$). A negative association of absolute differences runs opposite to a yield-driven shift, and with seven pairs and differences of at most 0.08 the corpus gives no support for attributing the component to yield, consistent with Section VII. Stricter selection raises the component modestly, as the plateau filter does: with the stricter $R^2 < 0.90$ criterion the headline component is 0.185, and with no filter 0.150, against 0.163.

S3. Homogeneity, window sensitivity and shape

Homogeneity. Per-group standard deviations span 0.010 to 0.417 across the 15 groups. A parametric bootstrap of the spread of log per-group SD under a common τ gives $p = 0.27$, Bartlett's test on sessions-within-group residuals $p = 0.30$, and the power of the bootstrap test against a two-point mixture of 0.5 and 1.5 times τ is 10%. On the seven headline groups, which the budget assumes, per-group SDs span 0.069 to 0.266 and both tests give p near 0.95, but with one degree of freedom per group they have almost no power. Heterogeneity is neither detected nor excluded.

Window sensitivity. Restricting the samples entering the median to $-6 < \log_{10}(G/G_0) < -1.5$ gives 0.162 for the headline against 0.163 unrestricted; -5 to -1.5 gives 0.157, -6 to -2 gives 0.138 and -4 to -1.5 gives 0.140. The component is stable to the noise-floor cut and falls modestly when the window removes the upper or lower part of the molecular distribution.

Shape. Five shape statistics of the per-group spreads were compared with 20 000 corpora simulated under a Gaussian session effect of the same τ and group sizes; all lie inside the null (smallest two-sided $p = 0.54$). With 15 groups this check has little power and supports no claim about the distribution.

S4. The mixture extension

The mixture extension. Predictions were fixed before any mixture trace was loaded. The component transfers to the new specimen class (0.225 log units, five of eight groups clearing their null). The three-session groups give $\tau = 0.250$ (chi-square on 6 df 0.16 to 0.55) against 0.189 (5 df, 0.12 to 0.46) from the two-session mixture groups, so the exchangeable-draw assumption behind the allocation rule is not contradicted, a check with power only against large departures. Across the 19 pairs with a recorded round or date, the difference does not grow with the gap between them (round-number and calendar gaps pooled, and pairs within three-session groups dependent, so the check is coarse) (Spearman -0.15, $p = 0.54$). One prediction failed: adding the mixture groups widened the bootstrap interval (0.137 against 0.090 in log units), because they are more heterogeneous.

S5. Candidate remedies: summary table, the two-campaign test and the AFM deposit

A force-spectroscopy deposit [3] with four constructs, each in three independent sessions covering six retract speeds, separates a shared offset from an occasion-by-condition interaction: against a null that permutes session identity within speed (2000 permutations), a common offset is detectable in one construct of four (22.4%, $p < 0.001$; the others 2.4, 5.2, 1.2%, $p > 0.3$), while the session-by-speed interaction exceeds per-cell bootstrap noise in 3 of 4 (excess 5.4 to 8.7%, parametric bootstrap $p \leq 0.011$; $p = 0.27$ in the other).

TABLE S1
CANDIDATE REMEDIES AND WHAT THE MEASUREMENT SAYS

remedy	result
a more reproducible summary statistic	a leave-one-group-out search selects a tail: the 0.95 quantile in 9 of 9 folds on the registered nine groups (+68%), the 0.05 quantile in 14 of 15 folds on the current set (out-of-sample between-session SD 66% below the median's). Rejected: the molecular signal shrinks with the session noise (signal-to-session-noise 1.42 at the 0.05 quantile, 1.11 at the 0.95, 1.47 for the median; best of 14 statistics, post hoc, 1.80)
a co-measured reference standard	three molecules in the same two campaigns shift −0.38, +0.13, +0.22 log units (7 to 9 per-session bootstrap SE each), shared fraction −0.50; crossed AFM deposit: shared offset in 1 of 4 constructs, interaction in 3 of 4; another laboratory's bare-gold reference stage does not predict the sample offset (Spearman −0.11, 95% CI −0.80 to 0.70, $n = 8$). Same-session co-measurement of a reference molecule is untested
a within-session self-diagnostic	none of 5 predictors on the 18 registered sessions; on the 14 headline sessions the smallest p is 0.05; drift predicts error on 33 sessions (exploratory) and fails to replicate on 70 AFM and 8 Delft sessions
trace selection, two surrogates	k-means class ($k = 3$, 50-bin histograms): component ratio 1.07 [0.95, 1.16] as registered (17 groups), 1.02 on the headline; plateau-length filter (54% of traces kept): 0.207 on the headline against 0.163; the published pipelines of [1], [2] were not run
standardising to a common decay time	within-session regression of trace median on the log time to fall one decade after entering the window: SD to 0.50 of its value on the headline, 0.42 on the held-out mixture groups; a noise placebo (trace median plus noise) gives 0.47 and 0.41, so the reduction is not evidence. Pre-registered on the public deposits: Delft NBD 0.97, Delft PAH 0.68, Liverpool 7.5 (a same-day pair, 1 df), geometric mean 1.70; stop condition met (noise placebo 0.89)




Fig. S1. The two-campaign reference-standard test. Three molecules measured in the same two campaigns shift in different directions, so a campaign-level co-measured standard would not correct the effect.

S6. D-STUDY FIGURE

Fig. S2 is the D-study of Section VIII: the expanded uncertainty of a reported conductance against the number of independent sessions, for four trace counts per session.

S7. OTHER LABORATORIES AND TECHNIQUES: EACH ANALYSIS

Fig. S3 collects every value of this note with the paper's own.




Fig. S2. D-study for the headline component: 95% expanded uncertainty of a reported conductance against the number of independent sessions, for four trace counts per session, with k from the Welch-Satterthwaite degrees of freedom. The curve assumes the corpus component transfers to the user's molecule.




Fig. S3. Standard deviation of a single reported conductance at each nesting level: this corpus with both estimators, two published MCBJ datasets recomputed from their tables, three public deposits from two other laboratories re-analysed with this paper's estimator at their nesting levels (the same-sample rows are repeatability-type), and a thesis between-day upper bound. Chi-square intervals on the stated degrees of freedom; arrows give upper limits beyond the axis.

Published tables. From Table S1 and Fig. S3 of [1] (OPE3, 100 mV, eleven datasets on six samples, log-normal-fit peak) the between-dataset SD is 0.169 log units, and the sample nesting splits it into 0.099 within sample and 0.200 between samples. From Table 1 of [2] (OPV3-2BT-H, eight datasets on three samples) it is 0.063 raw and 0.032 after clustering.

Delft, small deposit [4]: its files hold two runs of norbornadiene NBD-1 three months apart at identical settings and two of NBD-3 six days apart. Pre-registered, with the same median estimator, tunnelling filter and re-partition null, both pairs clear their null; the medians differ by a factor of 1.62 for the three-month pair and 1.05 for the six-day pair, a pooled SD of 0.106 log units (2 df), 0.06 to 0.17 across noise-floor settings with the same verdict at each.

Delft, large deposit [5]: two instruments, four molecules, a folder per sample with a bare-gold reference stage and a

molecule stage. Pre-registered with 0.10 V traces only (two samples mix biases and were screened per trace, keeping 1447 and 2254 of 3000 drawn; a first run without that screen is preserved with the released files and gave 0.39) and up to 3000 traces per session. Three nesting levels are reported separately. Consecutive days of one sample: 0.05 (2 df). Three samples recorded within two weeks, all on the LC instrument: 0.24 (chi-square on 2 df 0.12 to 1.5), that laboratory's intermediate-precision figure under the definition of Section II. Samples recorded between 2020 and 2023: 0.34 (2 df, 0.18 to 2.1), a campaign-level quantity that plausibly includes solution batches. The pre-registered bound of 0.35 applied to the pooled between-sample value over the three groups, which pools the two between-sample levels. As registered it was exceeded (0.41, 5 df). After submission of that run the inventory was checked file by file: the folders LC/2-OS/23-2-2023 and Piezo/2-OS/23-2-2023 hold the same 18480 files (equal checksums and sizes), and the file headers of that folder (input range, breaking speed, sampling rate, settling time, amplifier setting) match every LC session and none of the Piezo sessions. The folder is therefore counted once, as an LC recording; the Piezo/2-OS group keeps its two remaining samples, and the pooled value becomes 0.29 (4 df), inside the band. Both runs are released (pah_mcbj.py as registered, pah_mcbj_corrected.py). Pooling all three levels including the day level would give 0.24 on 6 df. The sample-level SD is 5.3 times the day-level SD, but on 2 and 2 df the ratio is not resolved ($F = 28$, two-sided $p = 0.07$, 95% interval on the variance ratio 0.7 to 1091), and the day level comes from the Piezo instrument, the sample level from the LC instrument; sample and campaign are not separated either. The one LC recording of 2-OS prepared ex situ lies 0.87 log units below the mean of the two Piezo samples of the same molecule, a single between-instrument comparison on different days that cannot be separated from a preparation difference.

Liverpool [6]: two compounds carry second measurement folders; one is a byte-identical copy of the first (identical hashes and timestamps) and is excluded, and the MnPARA 300 mV pair is a repeat of the same sample 1.8 hours later on the same day, a repeatability-condition repeat, not an occasion. At the pre-registered noise floor ($10^{-5}\,G_0$) it clears its null with an SD of 0.040 (medians differ by a factor of 1.14); at floors of 10^{-6} and $10^{-4}\,G_0$ the SD is 0.15 and 0.21, also clearing. For this pair the noise-floor choice is not a small matter, and it is not used as a repeatability figure.

A between-day bound. Imidazole and benzimidazole STM-BJ peaks averaged over at least four experiments on different days with at least 5000 traces each show "day-to-day variation in G1-3 peak positions … less than 3%" [7], an upper bound of 0.013 log units on a between-day component, under preparation conditions the thesis does not specify.

One example each from two other techniques. The same procedure gives, for one AFM force-spectroscopy deposit [3], a median ratio of between-session spread to within-session bootstrap standard error of 2.2 (24 groups of three sessions), and for one localisation-microscopy deposit on a nuclear-pore reference standard [8] a between-date SD of the ring radius of 0.38 nm against a mean per-localisation precision of 3.5 nm in the deposited files, a ratio of 0.11.




Fig. S4. Standard deviation of the reported value against the number of sessions a fixed budget of 1500 traces is split across, headline component. All points are the variance model with the headline τ and σ_{trace}; the shaded band marks the one to three sessions per group the corpus observes, and the square the third session, available only in the mixture groups. Short sessions add start-up variation (Section VIII), so the curve overstates the gain.




Fig. S5. Independent sessions per molecule required for each of 26 comparisons at matched settings, with the headline component, logarithmic scale. The observed differences include session noise, so the counts are biased low.

S8. Pre-registration ledger

Table S2 lists every pre-registered analysis with its archived file, registered criterion and outcome; the registration files are released unchanged, with their hashes.

S9. The session in recording order: methods, tables and the allocation figure

Data and series. Recording order is the cell order of the deposit's trace files (Data_LPF_R100.mat), identical to the analysis cache for all 64 sessions; there is no per-trace clock, so the order is an assumption. The released registration carries a dated post-run addendum (its Section 10); the pre-run file is released beside it as PREREG_noise_v1.md, and its hash is the first line of PREREG_noise.sha256. The deposit's processing (SMBJClassifier, DPP.py) starts a trace at the last sample above the 10 V saturation output and ends it at the third sample below 1 mV of amplifier output (one before the last when fewer than four fall below), so every trace ends at a fixed recorded current. Block statistic b_T: median of the

per-trace medians of 50 consecutive kept traces (primary); b_P: median of the pooled samples, the paper's indication. Floor c_0: variance of the block statistic over 200 random permutations of trace order within the session. Semivariogram $\gamma(h)$ at lags 1 to 20 blocks within sessions, averaged with equal weight per session; V = mean $\gamma(4..10)/c_0$; null: 2000 within-session shuffles. Gate: G is the weighted residual sum of squares of a linear (random-walk) fit divided by that of an exponential-plus-floor fit; g_{95} from 2000 random-walk-plus-floor surrogates matched at lag 1 with the actual block counts; power from Ornstein-Uhlenbeck surrogates at the fitted parameters; specificity from fractional ($d = 0.45$) and linear-trend surrogates. Step: γ_X is the session-pair-equal cross-session semivariance of b_T within a group, S the model-free plateau (mean γ over lags 10 to 20), $R = \sum \gamma_X / \sum S$ with equal group weights.

Table S3 gives the per-laboratory values. The Liverpool MnPARA folder is excluded: its 1778 files carry five copy timestamps within eight seconds and the per-trace medians trend across the file numbering, so the index is not a recording clock. The PAH dependence sits in the two LC sessions (2 of 2 clear their envelope) and in one Piezo session (1 of 5); the NBD pooled-sample statistic shows more dependence than the per-trace one (1.81 against 1.38), so trace-length weighting carries extra dependence there. The NBD runs contain thirteen scripted pauses of about 18 minutes; the semivariance across a pause lies above the within-run value at the equivalent lag in three runs and below it in three, with one to six pauses per run, so the pre-declared clock question is uninformative, as registered.

Placebos and controls (DNA corpus). Trace length (A1): 4.18 times its floor; block-level regression removes 21% of the excess and the trace-level version with the residual's own floor removes none, so trace length explains 0 to 21%. End level (A2): 2.64 times its floor; the registered block-level regression removed 96%, a figure inflated by shared sampling noise; at trace level with the residual's own floor it removes 81% of the excess and leaves $V = 1.76$ against a shuffle 95th percentile near 1.1. Per-block pass rate of the tunnelling filter (A3): 34%. Ceiling-censored statistic (A4): the logged value divided a pooled-sample statistic by a per-trace floor and is void; computed consistently, $V = 2.33$, inside the registered interval. Start-up (PF): 2.41 after dropping the first quarter. Shuffled sessions (PB): 1.00. The registered positive control (PC) compares two expressions of the same quantity and is void. Composition (PE): regressing the block median on the per-block pass rate and on the fraction of traces with at least 20 samples changes the cross-session excess by -2%.

What moves (post hoc). Per-trace quantile series through the same pipeline, block-level drift SD relative to the median's: trace maximum 0.03, 0.9 quantile 0.99, 0.1 quantile 0.65, end level 0.56, minimum 0.25. The end level correlates with the trace median at $r = 0.60$ (median over sessions). The deposit's exponential-decay class ($R^2 \geq 0.95$, 57 sessions) follows the molecular block median with slope 0.17 (quartiles 0.03 to 0.46).

Peak and weight (post hoc, after review). For each headline pair, the 0.05-decade histogram mode on the paper's window (smoothed), the fraction of samples within 0.5 decade of it, and the pooled-sample median: reweighting the second session so that its near-mode fraction equals the first session's removes 11% of the squared median shifts (position term 0.174 against weight term 0.055 log units, as between-session SDs); the mode itself has a between-session SD of 0.460 because it jumps between plateau families; the fitted peak of PREREG_peak.md, where it converges, moves less than half as much as the median in 2 of 5 single-duplex groups and more in the others. The median therefore follows the distribution's bulk, which is what a session moves; a peak-based indication shows less of the movement in some groups and more in others.

Block-wise null (post hoc, after review). The paper's re-partition null deals single traces into pseudo-sessions and assumes exchangeable traces. Dealing contiguous blocks of 100 traces instead, so that the serial dependence stays inside each pseudo-session, raises the null's 95th percentile for the headline groups from 0.040 (median 0.047 under blocks) to 0.070 log units; the observed s_I is 2.3 times that value.

Table S4 gives, per headline session, the three indications a reader may prefer; the per-trace-median block statistic of Section VI agrees with the pooled median on every quantity reported there.

Allan view. Per-trace median series: Allan-variance slope -0.84 up to 50 traces and -0.52 beyond (white gives -1); Allan deviation at 200 traces 2.1 times the white extrapolation; Hadamard over Allan at long averaging 0.96. The block-length Allan minimum lies between 50 and 200 traces in 30 of 58 sessions and is undefined below 400 traces in 24, so the pooled range rather than a per-session minimum is the design quantity.

Budget quantities. Effective number of independent 50-trace blocks in 1000 traces 6.1 (model-free) or 7.9 (fitted model); within-session standard error of a 1000-trace median 0.087 log units against the trace-resampling 0.022 (median over the headline sessions), inflation 3.2 (1.2 to 6.2); s_I over the drift-inclusive within-session SE 1.9; drift share of s_I^2 under the stationary reading 29%. The paper's $(s_I/u)^2$ rule is unchanged: about 12 sessions for $u = 0.05$ log units.

Step statistics (descriptive, branch B). $R = 1.59$ (group bootstrap 1.31 to 1.85); 12 of 15 groups and 6 of 7 headline groups above 1. Against the fitted short-range process the one-sided p is 0.0005; against a fractional ($d = 0.45$) process matched to the within-session curve the 95th percentile of R is 1.49; a random walk gives a median of 2.3. The direct mean square of the headline session differences is 0.0250 against 0.0304 from $\tau^2 + S/n_{\mathrm{eff}}$; the difference is the block-against-pooled difference of the two statistics. Fig. S5 gives the sessions each of the 26 molecule comparisons at matched settings needs, from the headline component (Section VIII of the paper). Fig. S4 is the paper's model-based allocation curve (sixty sessions of a hundred traces against one of 6000 are 6.4 times more precise under the model); the within-session term it uses is the resampling term, so its gains are upper bounds twice over.

REFERENCES

[1] D. Cabosart *et al.*, "A reference-free clustering method for the analysis of molecular break-junction measurements," *Appl. Phys. Lett.*, vol. 114, no. 14, art. 143102, 2019.

[2] N. D. Bamberger, J. A. Ivie, K. N. Parida, D. V. McGrath, and O. L. A. Monti, "Unsupervised segmentation-based machine learning as an advanced analysis tool for single molecule break junction data," *J. Phys. Chem. C*, vol. 124, no. 33, pp. 18302–18315, 2020.

[3] A.-M. Tsirigoni, M. Goktas, and K. G. Blank, "Chain sliding versus β-sheet formation upon shearing single α-helical coiled coils," Zenodo, 2023, doi: 10.5281/zenodo.7713350.

[4] L. Ornago and S. van der Poel, "Break junction data underlying the publication: Exploring the impact of select anchor groups for norbornadiene/quadricyclane single-molecule switches," 4TU.ResearchData, 2023, doi: 10.4121/022fea93-0e9a-4477-8363-41f709e96ca4.

[5] T. Baum, S. van der Poel, S. Fernández, M. Vilas-Varela, D. Peña, and H. S. J. van der Zant, "Fast-breaking MCBJ data underlying the thesis: 'Electronic transport signatures of two-electron interactions in all-organic single-molecule junctions'," 4TU.ResearchData, ver. 1, 2024, doi: 10.4121/18ef279f-b771-466e-bc7c-04b365c7e7c8.

[6] A. Vezzoli, N. Ferri, and R. J. Nichols, "Single-molecule conductance data for a series of ligands and their metal carbonyl complexes," Univ. Liverpool Data Catalogue, 2023, doi: 10.17638/datacat.liverpool.ac.uk/2244.

[7] X. Pan, "Single molecule conductance spectroscopy: probing the gold-bio interface at the atomic scale," Ph.D. dissertation, Dept. Chem., Boston Univ., Boston, MA, USA, 2024, ch. 4, p. 50 and Table 4-1. hdl: 2144/49785

[8] J. V. Thevathasan *et al.*, "Nuclear pores as versatile reference standards for quantitative superresolution microscopy," *Nat. Methods*, vol. 16, no. 10, pp. 1045–1053, 2019, doi: 10.1038/s41592-019-0574-9.

TABLE S2
PRE-REGISTERED ANALYSES: ARCHIVED FILE, REGISTERED CRITERION, OUTCOME

date	file	registered criterion	outcome	adverse
2026-08-17	PREREGISTRATION.md	peak-based understatement factor $F > 2$; K1: if median $F \leq 2$, "there is no paper. Report and stop"; K2: the pseudo-session control must collapse; K3: at least 8 usable groups, else a pilot; K4: report any qualitative dependence of F on the estimator	stop condition met ($F = 1.57$, archived 2026-08-17; F is the between-session SD over the pooled-histogram bootstrap SE; its equality with the current fitted-peak ratio is coincidental); peak mode-jumps; the analysis continued with the median, adopted post hoc, contrary to K1. K2 met (pseudo-session τ 0.069 against 0.214). K3 met at registration (13 groups); the current headline rests on seven groups after later corrections. K4 met: F is 1.43 and 1.50 for the two peak estimators and 7.94 for the median of per-trace medians (control 3, printed in 2026-08 and archived from a 2026-10 rerun of the unchanged script), so the understatement factor depends on the estimator, as Section III reports	yes
2026-08-18	PREREG_estimator.md	most stable quantile on nine groups; kill if the winner tracks the instrument	numerical criteria met (9 of 9 folds, +68%); physical criterion fired; q0.95 rejected	yes
2026-08-18	PREREG_standard.md	shared campaign offset: decision rule, no direction	"not shared" (signs disagree)	no
2026-08-18	PREREG_mixture.md	mixture-decomposition ratio $R > 0.70$; kill if the null matches	stop condition met (null 0.946 against 0.993)	yes
2026-09-27	PREREG_set2.md	mixture sessions: τ in [0.08, 0.30]; null cleared; combined CI narrows	first two met; CI widened	yes
2026-09-27	PREREG_classfilter.md	k-means class ratio ≤ 0.5; kill if its CI includes 1	stop condition met (1.07)	yes
2026-09-30	PREREG_delft.md	small Delft deposit: a pair clears; pooled SD in [0.05, 0.35]	both met	no
2026-09-30	PREREG_liverpool.md	Liverpool: a pair clears; pooled SD in [0.05, 0.35]	first met; second failed (0.029, pulled down by a duplicate folder found and excluded afterwards; 0.040 after exclusion, still outside the band)	yes
2026-10-01	PREREG_pah.md	large Delft deposit: τ in [0.05, 0.35]; null; day $<$ sample; gold reference null	bound exceeded as registered (0.41); 0.29 after a duplicated folder was found and counted once (inventory correction, both runs released); others met; bias screen omitted in first run (0.39; deviation)	yes
2026-10-01	PREREG_drift_replication.md	drift predicts error on AFM: $\rho > 0$, $p < 0.05$	failed (+0.04, $p = 0.77$)	yes
2026-10-01	PREREG_peak.md	fitted-peak component in [0.104, 0.234], the 1.5× band around the then-current 0.156 (nine groups); kill if more than 2 of 18 fits fail	band met (0.165 on the 5 of the registered nine groups that converge); the archived script counted 7 failures over all 37 sessions (deviation); recounted on the registered 18, 4; the kill fires either way: estimator not well defined on these distributions. Current headline: 0.183 on 4 groups	yes
2026-10-07	PREREG_lite_ mixture_holdout.md	decay-time standardisation, written before the mixture groups were loaded: SD ratio ≤ 0.60 there	met (0.42), but a noise placebo gives 0.41, so it is not evidence	no
2026-10-07	PREREG_casemix.md	the same on the public deposits: geometric-mean ratio ≤ 0.80 success, ≥ 1 kill; amended before any result (Delft data sections, duplicate folder)	stop condition met (1.70; Delft NBD 0.97, PAH 0.68, Liverpool 7.5)	yes
2026-10-10	PREREG_noise.md	three primaries: serial dependence of the block median (shuffle $p < 0.01$; apparatus placebos below half of the excess); identifiability gate (calibrated G, range bound, surrogate specificity, power); occasion step ($R \geq 1.5$, interval above 1, parametric null)	serial dependence: statistics met ($V = 2.60$, $p = 0.0005$) but the end-level placebo met its criterion, so the registered consequence applies (attribution withdrawn; the placebo is a trace quantile in this deposit, Note S9); gate failed on specificity against long memory (the code tested the random-walk surrogate only; the outcome is the same under the registered text); step not evaluated as a claim; the registered positive control is an identity and is void; one stray line removed from the script between stages, recorded	yes

TABLE S3
RECORDING-ORDER ANALYSIS BY LABORATORY (BLOCK LENGTH 50 TRACES)

	DNA corpus	Delft NBD	Delft PAH
records (traces each)	58 (1000 to 3217)	7 (1333 to 9957)	7 (9985 to 10000)
V, b_T (shuffle p)	2.60 (0.0005)	1.38 (0.001)	1.19 (0.001)
V, b_P	2.79	1.81	1.23
records above own shuffle 95th pct	51 of 58	4 of 7	3 of 7
floor SD, plateau SD (log units)	0.138, 0.227	0.083, 0.115	0.118, 0.145
fitted range (traces)	97 (55 to 374; headline)	874 (127 to 2889)	1558 (642 to 6001)
gate (power)	fail (0.89)	underpowered (0.37)	underpowered (0.07)
R session-to-session over block-to-block	1.59 (1.31 to 1.85)	2.15	4.49

TABLE S4
FOUR INDICATIONS PER HEADLINE SESSION (FIRST, SECOND SESSION): POOLED-SAMPLE MEDIAN, SMOOTHED 0.05-DECADE HISTOGRAM MODE, FITTED PEAK (PREREG_PEAK.MD) WHERE IT CONVERGES; $\log_{10}(G/G_0)$

group	median	mode	fitted peak
Delta/MM1/100	−3.74, −3.51	−4.03, −3.13	no fit
Delta/MM1/30	−3.46, −3.22	−3.83, −3.53	no fit
Delta/MM2/100	−4.01, −3.82	−4.53, −4.43	−4.42, −4.42
Delta/PM/30	−2.90, −3.00	−2.83, −2.73	−2.79, −2.81
Beta/MM1/100	−3.56, −3.18	−3.53, −2.33	no fit
Beta/MM2/300	−4.06, −4.19	−4.08, −4.63	−4.18, −4.57
Beta/MM3/100	−3.59, −3.81	−3.83, −4.38	−3.93, −4.26