UNIVERSITY OF SRI JAYEWARDENEPURA

Faculty of Applied Sciences
Department of Computer Science

FIRST PROGRESS REVIEW

Doctor of Philosophy (PhD) in Computer Science

Chapter 1

Introduction

Calibrated Selective Autonomy in Multi-Agent Clinical Artificial Intelligence: A Clinician-Validated, Privacy-Preserving Framework for Brain-Tumour Diagnosis and Surgical-Planning Support in Low-Resource Settings

Candidate: KH Viraj Sameera
Mode: Full-time, 36 months (commenced 1 October 2026)
Principal Supervisor: Dr. WMKS Ilmini
Co-Supervisor: Dr. WGCW Kumara
Clinical Supervisor: Dr. AM Deepal K Attanayake

Version 1.1 · 11 November 2026 · Draft for supervisor review

Document Information

Item	Detail
Document	First Progress Review, Chapter 1: Introduction
Research title	Calibrated Selective Autonomy in Multi-Agent Clinical Artificial Intelligence: A Clinician-Validated, Privacy-Preserving Framework for Brain-Tumour Diagnosis and Surgical-Planning Support in Low-Resource Settings
Degree and department	PhD in Computer Science, Department of Computer Science, Faculty of Applied Sciences, University of Sri Jayewardenepura
Version and date	1.1, 11 November 2026
Status	Draft for supervisor review; not for circulation outside the supervisory team
Basis	Research proposal accepted by the Board on 4 November 2026; Research Paper Bank of 559 open-access papers with structured methodology extraction for every paper
Ethics status	KDU Faculty of Medicine ERC application submitted 10 November 2026 (decision awaited); USJ application to follow

Version history

Version	Date	Changes
1.0	10 November 2026	First complete draft for supervisor review.
1.1	11 November 2026	Quality revision: title page, document information, contents, list of tables and figures, abbreviations, running headers; progress section updated (methodology extracted for all 559 papers; KDU ethics application submitted); references standardised to APA 7th edition.

Contents

1. Introduction.....1
1.1 Background of the Research.....1
1.1.1 The brain-tumour pathway and the state of computational support.....2
1.1.2 The Sri Lankan context and the case for a national registry.....3
1.2 Problem Statement and Research Gap.....4
1.2.1 Why accuracy alone is not enough.....4
1.2.2 Identified research gaps.....4
1.2.3 Problem statement.....5
1.3 Aim, Objectives and Research Questions.....5
1.3.1 Aim.....5
1.3.2 Research objectives.....6
1.3.3 Research questions.....6
1.4 Scope, Limits of Autonomy and Limitations.....7
1.4.1 Scope.....7
1.4.2 Limits of AI autonomy.....7
1.4.3 Limitations and assumptions.....8
1.5 Justification and Significance.....8
1.6 Progress Since Registration.....9
1.7 Structure of the Report.....10
References.....10

List of Figures

Figure 1.1: The brain-tumour care pathway, the specialised AI agent that supports each stage, and the clinician who retains final authority throughout (from the research proposal).....2

List of Tables

Table 1.1: Research gaps, the evidence that confirms them, and the objective that addresses each.....5
Table 1.2: Progress since registration (1 October to 11 November 2026).....9

Abbreviations

Abbreviation	Meaning
AI	Artificial intelligence
BraTS	Brain Tumor Segmentation challenge
ERC	Ethics Review Committee
EU	European Union
GLOBOCAN	Global Cancer Observatory estimates (IARC)
HITL	Human-in-the-loop
KDU	General Sir John Kotelawala Defence University
LLM	Large language model
MRI	Magnetic resonance imaging

Abbreviation	Meaning
NCCP	National Cancer Control Programme
WHO	World Health Organization

1. Introduction

This chapter introduces the research for the First Progress Review. It restates the background and the problem that motivate the study, updates them with evidence gathered during the first months of candidature, and sets out the aim, objectives, research questions, scope and significance of the work. The chapter is based on the research proposal accepted by the Board on 4 November 2026 and on the literature collected in the Research Paper Bank (559 open-access papers grouped by objective; see Chapter 2). Where the evidence reviewed since registration sharpens or qualifies a claim made in the proposal, this is stated explicitly.

1.1 Background of the Research

Artificial intelligence (AI) now supports diagnosis, treatment planning and clinical documentation at a scale no human reader could match (Topol, 2019; Khalighi et al., 2024). In medical imaging alone, a meta-analysis of 503 studies found high diagnostic accuracy for deep learning across ophthalmology, breast and respiratory imaging, while noting that very few of those studies were prospective or externally validated (Aggarwal et al., 2021). The same capability makes unsupervised deployment hazardous. Performance degrades under dataset shift: a pneumonia detector generalised poorly outside its training hospitals because it had learned site-specific artefacts rather than disease (Zech et al., 2018; Finlayson et al., 2021). And a model that reports a confident answer gives the clinician no signal that it has left the region in which it is competent (Kompa et al., 2021; Guo et al., 2017).

These findings have made human-in-the-loop (HITL) AI the dominant design philosophy for safety-critical medicine. HITL inserts control points at consequential moments so that a clinician can validate, refine or override a recommendation using findings, comorbidity and patient preferences that no model can see (Olawade et al., 2026). The evidence reviewed for this progress report shows, however, that simply placing a human beside an AI is not enough. A meta-analysis of 106 human-AI studies found that, on average, human-AI combinations performed worse than the better of the human or the AI alone in decision tasks (Malone et al., 2024); radiologists' benefit from AI assistance varied widely and incorrect AI output harmed their performance (Yu et al., 2024); and explanations increased acceptance of wrong recommendations (Bansal et al., 2021). By contrast, systems that decide *which* cases the AI should handle and which should go to the clinician

have improved both accuracy and workload: complementarity-driven deferral in mammography and tuberculosis screening reduced false positives and clinician reads at the same time (Dvijotham et al., 2023). The open engineering question, and the one this research addresses, is therefore not whether a human should be in the loop, but when, on which cases, and on what calibrated evidence.

1.1.1 The brain-tumour pathway and the state of computational support

The brain-tumour pathway is long, information-dense and unforgiving. It runs from non-specific neurological symptoms and a scan, through multi-sequence magnetic resonance imaging (MRI), biopsy and microscopic examination of tissue, laboratory tests on that tissue, a multidisciplinary decision, surgery and follow-on therapy, to years of surveillance scanning (Figure 1.1). Since 2021 the World Health Organization (WHO) classification has been integrated, so a diagnosis combines tissue appearance with molecular results (Louis et al., 2021), making the task multi-source by definition. Brain tumours carry disproportionate mortality and affect working-age adults and children (Price et al., 2024). Surgery must balance removing as much tumour as possible against preserving the regions controlling speech, movement and vision: extent of resection is repeatedly associated with survival (Sanai & Berger, 2008; Molinaro et al., 2020), while removal of healthy tissue causes permanent loss of function. The pathway is also dependent: an error in outlining the tumour propagates into grading, grading into treatment selection, and treatment selection into outcome.




Figure 1.1: The brain-tumour care pathway, the specialised AI agent that supports each stage, and the clinician who retains final authority throughout (from the research proposal).

Computational support has advanced rapidly for individual sub-tasks. Deep learning dominates tumour detection, outlining and classification on MRI, benchmarked on the

BraTS challenges (Menze et al., 2015; Baid et al., 2021), with nnU-Net establishing strong, self-configuring baselines (Isensee et al., 2021) that remain hard to beat when validated rigorously (Isensee et al., 2024). Computational pathology reaches clinical-grade analysis of whole digitised slides with weak, slide-level labels (Lu et al., 2021), and deep learning has achieved neuropathologist-level integrated classification of adult-type diffuse gliomas (Wang et al., 2023). Fusion of pathology and radiology foundation models now supports WHO 2021 glioma subtyping (Saueressig et al., 2026), and molecular classification has been brought into the operating theatre (Hollon et al., 2023; Vermeulen et al., 2023). Yet individual models solve individual tasks: no accepted architecture assembles them into a supervised workflow with defined accountability.

Two further developments frame this work. The first is agentic and multi-agent AI, in which a coordinating component routes work between specialised models or language-model agents. Multi-agent systems have outperformed single models on medical benchmarks (Kim et al., 2024; Chen et al., 2025), and an autonomous agent working inside a simulated electronic health record recently exceeded physicians on emergency cases (Ferber et al., 2026). However, a scoping review found only seven eligible studies of agentic AI in healthcare and a single randomised trial (Collaco et al., 2026), and audits of multi-agent systems show high failure rates and false consensus (Cemri et al., 2025; Zhu et al., 2025). The second is federated learning, which lets institutions train a shared model without disclosing raw data and has been proven for glioblastoma boundary detection across 71 sites on six continents (Pati et al., 2022).

1.1.2 The Sri Lankan context and the case for a national registry

GLOBOCAN 2024 estimates approximately 700 new brain and nervous-system cancer cases and around 587 deaths annually in Sri Lanka, with a five-year prevalence of roughly 2,500 (International Agency for Research on Cancer, 2024). Deaths are high relative to new cases, and care is concentrated in a few tertiary units served by a small pool of neurosurgeons, neuroradiologists and neuropathologists. A National Cancer Registry exists for incidence and mortality reporting (National Cancer Control Programme, 2024), but there is no imaging- and tissue-linked brain-tumour registry; patients are tracked in fragmented, institution-specific records. Clinically this impedes continuity of care, multidisciplinary review and outcome auditing. Scientifically it makes locally valid AI almost impossible to build, because models trained on foreign cohorts do not transfer reliably to local scanners, protocols, laboratory practice or case mix.

Evidence from comparable settings supports this concern. The BraTS-Africa challenge was created because routine-care MRI in Sub-Saharan Africa, often from older or low-field scanners with poorer contrast and from patients presenting later, challenges models developed on Western benchmark data; the winning teams needed heavy augmentation and ensembling to cope with a small local dataset (Adewole et al., 2026; Ankomah et al., 2025), and reviews of AI in low- and middle-income countries report few real-world evaluations, limited data and poor fit to local workflows (Ciecierski-Holmes et al., 2022; Yevudza et al., 2024). The project therefore separates two layers. The **platform layer** is Sri Lanka's first National Brain Tumour Patient Registry, which gives the research authentic data and users and is a real-world deliverable in its own right. The **research layer**, the focus of this PhD, is the design, formalisation and validation of the AI framework that makes such a platform safe, data-efficient and privacy-preserving.

1.2 Problem Statement and Research Gap

1.2.1 Why accuracy alone is not enough

A model that is 95% accurate is not automatically safe: the danger lies in the 5% of confident mistakes, and in the absence of a mechanism by which the system can recognise the limits of its own competence. Modern deep networks are systematically over-confident unless explicitly calibrated (Guo et al., 2017), and calibration worsens precisely when a model meets data from a new hospital (Ovadia et al., 2019). Benchmarks of failure detection in medical imaging found that no published confidence-scoring method reliably prevented silent failures (Bungert et al., 2023; Bernhardt et al., 2022). The issue is therefore not a shortage of accurate models but the absence of a governing mechanism that decides, case by case, whether an output is fit to be acted upon. Where specialist review time is the scarcest resource, that mechanism also determines whether AI reduces clinician workload or merely relocates it (Chen et al., 2024).

1.2.2 Identified research gaps

The proposal identified six gaps. The literature reviewed since registration (Chapter 2) confirms each of them and adds specific evidence, summarised in Table 1.1.

Table 1.1: Research gaps, the evidence that confirms them, and the objective that addresses each.

Gap	What the literature shows	Objective
No principled policy for when to defer	Selective classification, learning to defer and conformal prediction exist but address a single model and a single decision with a uniform error cost (Geifman & El-Yaniv, 2017; Mozannar & Sontag, 2020; Angelopoulos & Bates, 2021); cost- and capacity-aware deferral has been studied only outside medicine or on simulated experts (Alves et al., 2024; Verma & Nalisnick, 2022).	O2
Unqualified hand-offs between agents	Medical multi-agent systems route by question difficulty or LLM self-rating, not calibrated uncertainty (Kim et al., 2024; Lou et al., 2026); unsupported observations propagate downstream in 16.6% of cases (Zhu et al., 2025); optimising components does not optimise the system (Bedi et al., 2025).	O1
Fusion that does not survive real data	Fusion beats single sources on complete cohorts (Chen et al., 2022; Saueressig et al., 2026) but most methods assume every source is present; missing-modality work is simulated on BraTS (Zhang et al., 2022; Liu et al., 2023).	O3
Methods that assume abundant, standardised data	Public benchmarks are mostly Western and pre-processed (Baid et al., 2021); routine-care MRI from low-resource settings needed a dedicated benchmark and adaptation (Adewole et al., 2026; Ankomah et al., 2025); transfer and self-supervised learning help with limited data (Boyd et al., 2024; Tang et al., 2022).	O3
Data sharing in a few small, heterogeneous hospitals	Federation works at international scale (Pati et al., 2022; Zenk et al., 2025) but the effect of aggregation on calibration, and the small-federation regime, are little studied (Manthe et al., 2024; Koutsoubis et al., 2024).	O3
No way to measure the quality of the escalation decision	Reporting guidelines require human factors and safety in situ (Vasey et al., 2022; Sounderajah et al., 2025) but do not define escalation-quality metrics; selective prediction can change the pattern of clinician errors (Jabbour et al., 2025).	O4

1.2.3 Problem statement

These gaps identify the central research problem as the absence of a clinically responsible, formally governed multi-agent AI framework that can support the brain-tumour pathway across imaging, tissue analysis, combined diagnosis and pre-operative surgical planning while explicitly managing uncertainty and preserving clinician authority. Such a framework must decide, for each case and each sub-decision, when an agent can provide reliable decision support and when the case must be escalated to a qualified clinician. It must remain robust to the small, heterogeneous, non-standardised datasets of Sri Lankan institutions, and enable privacy-preserving collaboration without centralising raw patient data.

1.3 Aim, Objectives and Research Questions

1.3.1 Aim

The aim is to design, develop and evaluate an uncertainty-aware, human-in-the-loop multi-agent AI framework for brain-tumour care, integrating imaging, tissue analysis, combined

clinical decision support and surgical planning, and to evaluate its feasibility within a national brain-tumour registry in Sri Lanka. Human oversight is implemented as an explicit, measurable mechanism rather than treated as a guiding principle.

1.3.2 Research objectives

The objectives are unchanged from the accepted proposal and are deliberately limited to four:

- **O1.** To design and develop a multi-agent AI architecture for the brain-tumour care pathway (patient intake, MRI analysis, tissue analysis, diagnosis and staging, surgical planning and follow-up), in which an orchestrator routes tasks, propagates uncertainty across agent hand-offs, and keeps an auditable record of every AI output and every clinician validation.
- **O2.** To develop and evaluate a calibrated, uncertainty-aware escalation mechanism that decides when an agent may provide a decision-support output and when a case or sub-decision must be referred to a clinician, based on selective prediction, learning to defer and the clinical cost of error.
- **O3.** To develop and evaluate the data-efficient models the agents depend on (MRI detection, outlining, classification and anatomical reconstruction; analysis of digitised biopsy slides; and their combination with available laboratory markers) on small, heterogeneous Sri Lankan data, including privacy-preserving federated training across institutions, and to benchmark their performance and calibration against established approaches.
- **O4.** To clinically evaluate the human-in-the-loop framework, covering diagnostic and decision-support performance, the appropriateness of escalation decisions, clinician workload, usability, trust and safety, with surgical planning evaluated as a proof-of-concept component.

1.3.3 Research questions

The research questions correspond one-to-one with the objectives:

- **RQ1.** How can specialised AI agents be orchestrated across the brain-tumour care pathway so that uncertainty is propagated between them, information is coordinated, and an auditable record of AI outputs and clinician validations is maintained?

- **RQ2.** How can uncertainty estimation and calibration be incorporated into the framework to determine when the system should provide a decision-support output and when it should defer to a clinician?
- **RQ3.** How effectively can data-efficient and multi-source methods detect, outline, classify and grade brain tumours from small and heterogeneous Sri Lankan MRI and biopsy-slide data, including under federated training across hospitals, and how do their performance and calibration compare with established approaches?
- **RQ4.** To what extent can the human-in-the-loop framework improve clinically relevant decision support while maintaining appropriate clinician involvement, acceptable workload, usability, trust and safety, including the feasibility of MRI-based surgical-planning support?

1.4 Scope, Limits of Autonomy and Limitations

1.4.1 Scope

Depth of investigation is differentiated across three tiers. The first and primary tier is the uncertainty-aware orchestration and escalation mechanism, supported by data-efficient MRI models; this is the central methodological contribution. The second is computational pathology and combined decision support, which remain decision support rather than autonomous diagnosis. The third is privacy-preserving federated learning. The surgical-planning component is a proof of concept covering patient-specific anatomical reconstruction, outlining of the tumour and structures at risk, and visualisation of surgical risk; it makes no autonomous surgical decisions. The registry is enabling infrastructure. The study is limited to participating Sri Lankan institutions and to tumours included in the registry under predefined eligibility criteria.

1.4.2 Limits of AI autonomy

Autonomy is conditional, uncertainty-aware and task-specific. The system does not independently diagnose, decide treatment or perform surgery. Each agent produces a decision-support output with a calibrated uncertainty; cases exceeding predefined thresholds are escalated, and the final decision remains the responsibility of the treating clinician. The system never performs or controls an operation: it plans, predicts, flags and recommends, and the neurosurgeon executes and validates every consequential step. This follows international guidance that places human autonomy, transparency and

accountability among the core principles of AI for health (World Health Organization, 2021), the requirement in the EU AI Act that high-risk systems be designed so that people can oversee and interrupt them (European Union, 2024), and the FUTURE-AI consensus on traceability and usability (Lekadir et al., 2025). Three boundaries follow: bounded decision authority (no diagnosis, grade or treatment decision of record is issued by the AI); a conservative default (under uncertainty or novelty the system defers); and transparency of basis (every autonomous action carries its evidence, confidence and a model-facts statement of the conditions under which it was validated (Sendak et al., 2020; Mitchell et al., 2019)).

1.4.3 Limitations and assumptions

Component performance may vary across tumour types, imaging protocols, institutions and patient populations, and the federated component depends on the number of participating institutions, computing infrastructure and connectivity. Intra-operative guidance, robotic actuation, new laboratory assays, radiotherapy and chemotherapy planning, paediatric-specific modelling and medical-device certification are outside scope. Three further limitations are stated plainly. **Cohort size:** national incidence means the registry will accrue hundreds rather than thousands of fully linked cases, which bounds subgroup analyses and makes data-efficient methods necessary. **Reference standard:** clinician-confirmed diagnoses serve as ground truth, so the framework is evaluated against expert consensus. **Study type:** this is not a randomised trial of patient outcomes; validation measures diagnostic performance, workload, escalation appropriateness, safety and trust in an assistive workflow (Vasey et al., 2022), and no survival benefit is claimed. The work assumes that ethics approval and data-sharing agreements are secured before data access (the KDU application was submitted on 10 November 2026), that units can supply usable MRI and digitisable biopsy material, and that clinician time is available, though scarce. Each assumption has a planned mitigation.

1.5 Justification and Significance

Scientific. How an AI agent should decide when it is competent to act and when it must defer is a computer-science problem before it is a clinical one. It sits at the intersection of uncertainty quantification (Gal & Ghahramani, 2016; Lakshminarayanan et al., 2017), calibration (Guo et al., 2017; Ovadia et al., 2019), selective prediction and learning to defer (Geifman & El-Yaniv, 2017; Mozannar & Sontag, 2020; Dvijotham et al., 2023) and

multi-agent orchestration (Liu et al., 2025). The literature treats these as separate concerns. Formalising selective autonomy over a chain of dependent sub-decisions, and evaluating it on an explicit workload-versus-safety frontier, is methodologically novel and transferable beyond this clinical domain. Recent preprints that add conformal risk control to a single routing step (Li et al., 2026) or a safety governor to a tumour-board agent (Liu et al., 2026) show that the field is moving in this direction, which makes a rigorous, clinically validated treatment timely.

Clinical and national. Brain-tumour care is an unusually strong test case: it requires integration of several evidence sources (Louis et al., 2021), uncertainty must propagate along a sequential chain, the cost of error is severe and asymmetric (Sanai & Berger, 2008; Molinaro et al., 2020), and specialist review time is the binding constraint. Establishing Sri Lanka's first imaging- and tissue-linked brain-tumour registry delivers durable value to the participating units independently of the research findings, and an associated de-identified benchmark addresses the near-absence of South Asian cases in public brain-tumour datasets (Baid et al., 2021; Adewole et al., 2026).

Ethical, regulatory and methodological. A framework that makes oversight measurable and auditable rather than declared addresses a requirement that is becoming mandatory (European Union, 2024; Sounderajah et al., 2025). The constraints of the setting also make the results generalisable: small, heterogeneous data and barriers to centralisation are the normal condition of clinical AI in most of the world, so methods demonstrated under scarcity, heterogeneity and federation reach further than those requiring large curated corpora (Pati et al., 2022; Ciecierski-Holmes et al., 2022).

1.6 Progress Since Registration

The candidature started on 1 October 2026. The main activities completed in the first six weeks are summarised in Table 1.2.

Table 1.2: Progress since registration (1 October to 11 November 2026).

Activity	Status and date	Outcome
Research proposal	Accepted by the Board; handed over to the evaluators on 4 November 2026	Objectives, research questions and scope fixed for the candidature
Literature collection	Completed on 10 November 2026	Research Paper Bank of 559 open-access papers (O1 143, O2 109, O3 146, O4 161), each with its full text
Methodology extraction	Completed on 11 November 2026	Structured record for every paper: study design, data and sample size, preprocessing, model, training, evaluation, baselines, results, statistics and code/data availability

Activity	Status and date	Outcome
Ethics	KDU Faculty of Medicine ERC application submitted on 10 November 2026; decision awaited	University of Sri Jayewardenepura application to follow the KDU decision
Progress review chapters	Chapters 1 to 3 drafted (version 1.1, 11 November 2026)	Introduction, literature review and research methodology ready for supervisor review
Supervision and tracking	Progress website and log book maintained	Meetings, action items, documents and the Paper Bank available to the supervisory team

1.7 Structure of the Report

Chapter 2 reviews the literature by objective and identifies the open problems. Chapter 3 sets out the research methodology: design-science research with a mixed-methods evaluation, the agent architecture and escalation rule, the work packages, data, analysis and ethics. Chapter 4 (in preparation) will report preliminary work and results on public data, followed by the work plan and timeline.

References

Adewole, M., Rudie, J. D., Gbadamosi, A., et al. (2026). Brain tumor segmentation in Sub-Saharan Africa patient population: The BraTS-Africa challenge. *Neuro-Oncology Advances*. https://doi.org/10.1093/noajnl/vdag082

Aggarwal, R., Sounderajah, V., Martin, G., et al. (2021). Diagnostic accuracy of deep learning in medical imaging: a systematic review and meta-analysis. *npj Digital Medicine*. https://doi.org/10.1038/s41746-021-00438-z

Alves, J. V., Leitao, D., Jesus, S., Sampaio, M. O. P., Liebana, J., Saleiro, P., et al. (2024). Cost-Sensitive Learning to Defer to Multiple Experts with Workload Constraints. *Transactions on Machine Learning Research*. https://doi.org/10.48550/arxiv.2403.06906

Angelopoulos, A. N., & Bates, S. (2021). A Gentle Introduction to Conformal Prediction and Distribution-Free Uncertainty Quantification. *arXiv preprint*. https://arxiv.org/abs/2107.07511

Ankomah, C. T., Ayivor, L. E., Nyame, I., et al. (2025). How We Won BraTS-SSA 2025: Brain Tumor Segmentation in the Sub-Saharan African Population Using Segmentation-Aware Data Augmentation and Model Ensembling. *arXiv preprint*. https://arxiv.org/abs/2510.03568

Baid, U., Ghodasara, S., Mohan, S., Bilello, M., Calabrese, E., Bakas, S., et al. (2021). The RSNA-ASNR-MICCAI BraTS 2021 Benchmark on Brain Tumor Segmentation and Radiogenomic Classification. *arXiv preprint*. https://arxiv.org/abs/2107.02314

Bansal, G., Wu, T., Zhou, J., Fok, R., Nushi, B., Kamar, E., et al. (2021). Does the Whole Exceed its Parts? The Effect of AI Explanations on Complementary Team Performance. *ACM CHI 2021*. https://arxiv.org/abs/2006.14779

Bedi, S., Mlauzi, I., Shin, D., Koyejo, S., & Shah, N. H. (2025). The Optimization Paradox in Clinical AI Multi-Agent Systems. *arXiv preprint*. https://arxiv.org/abs/2506.06574

Bernhardt, M., Ribeiro, F. D. S., & Glocker, B. (2022). Failure Detection in Medical Image Classification: A Reality Check and Benchmarking Testbed. *Transactions on Machine Learning Research*. https://arxiv.org/abs/2205.14094

Boyd, A., Ye, Z., Prabhu, S. P., Tjong, M. C., Zha, Y., et al. (2024). Stepwise Transfer Learning for Expert-level Pediatric Brain Tumor MRI Segmentation in a Limited Data Scenario. *Radiology: Artificial Intelligence*. https://doi.org/10.1148/ryai.230254

Bungert, T. J., Kobelke, L., & Jaeger, P. F. (2023). Understanding Silent Failures in Medical Image Classification. *MICCAI 2023*. https://doi.org/10.48550/arXiv.2307.14729

Cemri, M., Pan, M. Z., Yang, S., Agrawal, L. A., Chopra, B., Tiwari, R., et al. (2025). Why Do Multi-Agent LLM Systems Fail?. *NeurIPS 2025 Datasets and Benchmarks Track*. https://arxiv.org/abs/2503.13657

Chen, M., Wang, Y., Wang, Q., et al. (2024). Impact of human and artificial intelligence collaboration on workload reduction in medical image interpretation. *npj Digital Medicine*. https://doi.org/10.1038/s41746-024-01328-w

Chen, R. J., Lu, M. Y., Wang, J., Williamson, D. F. K., Rodig, S. J., Lindeman, N. I., et al. (2022). Pathomic Fusion: An Integrated Framework for Fusing Histopathology and Genomic Features for Cancer Diagnosis and Prognosis. *IEEE Transactions on Medical Imaging*. https://arxiv.org/abs/1912.08937

Chen, X., Yi, H., You, M., Liu, W., Wang, L., Li, H., et al. (2025). Enhancing diagnostic capability with multi-agents conversational large language models. *npj Digital Medicine (2025) 8:159*. https://doi.org/10.1038/s41746-025-01550-0

Ciecierski-Holmes, T., Singh, R., Axt, M., et al. (2022). Artificial intelligence for strengthening healthcare systems in low- and middle-income countries: a systematic scoping review. *npj Digital Medicine*. https://doi.org/10.1038/s41746-022-00700-y

Collaco, B. G., Haider, S. A., Prabha, S., Gomez-Cabello, C. A., Genovese, A., Wood, N. G., et al. (2026). The role of agentic artificial intelligence in healthcare: a scoping review. *npj Digital Medicine (vol. 9, article 345)*. https://doi.org/10.1038/s41746-026-02517-5

Dvijotham, K., Winkens, J., Barsbey, M., Ghaisas, S., Stanforth, R., Pawlowski, N., et al. (2023). Enhancing the reliability and accuracy of AI-enabled diagnosis via complementarity-driven deferral to clinicians. *Nature Medicine 29(7):1814-1820*. https://doi.org/10.1038/s41591-023-02437-x

European Union. (2024). Regulation (EU) 2024/1689 laying down harmonised rules on artificial intelligence (Artificial Intelligence Act), Article 14: Human oversight. *Official Journal of the European Union*. https://eur-lex.europa.eu/eli/reg/2024/1689/oj

Ferber, D., Hilgers, L., Höper, C., Kinny-Köster, B., Eckardt, J. N., Egger-Heidrich, K., et al. (2026). Towards autonomous medical artificial intelligence agents. *Nature, 655:1282*. https://doi.org/10.1038/s41586-026-10675-5

Finlayson, S. G., Subbaswamy, A., Singh, K., et al. (2021). The clinician and dataset shift in artificial intelligence. *New England Journal of Medicine*, 385(3), 283-286. https://doi.org/10.1056/NEJMc2104626

Gal, Y., & Ghahramani, Z. (2016). Dropout as a Bayesian Approximation: Representing Model Uncertainty in Deep Learning. *ICML 2016*. https://arxiv.org/abs/1506.02142

Geifman, Y., & El-Yaniv, R. (2017). Selective Classification for Deep Neural Networks. *NeurIPS 2017*. https://arxiv.org/abs/1705.08500

Guo, C., Pleiss, G., Sun, Y., & Weinberger, K. Q. (2017). On Calibration of Modern Neural Networks. *ICML 2017*. https://arxiv.org/abs/1706.04599

Hollon, T., Jiang, C., Chowdury, A., Nasir-Moin, M., et al. (2023). Artificial-intelligence-based molecular classification of diffuse gliomas using rapid, label-free optical imaging. *Nature Medicine*. https://doi.org/10.1038/s41591-023-02252-4

International Agency for Research on Cancer. (2024). GLOBOCAN 2024: Sri Lanka fact sheet. *Global Cancer Observatory*, IARC/WHO. https://gco.iarc.who.int/media/globocan/factsheets/populations/144-sri-lanka-fact-sheet.pdf

Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J., & Maier-Hein, K. H. (2021). nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation. *Nature Methods*. https://arxiv.org/abs/1809.10486

Isensee, F., Wald, T., Ulrich, C., et al. (2024). nnU-Net Revisited: A Call for Rigorous Validation in 3D Medical Image Segmentation. *MICCAI 2024 (LNCS)*. https://doi.org/10.1007/978-3-031-72114-4_47

Jabbour, S., Fouhey, D., Banovic, N., Shepard, S. D., Kazerooni, E., Sjoding, M. W., et al. (2025). Selective Prediction Reduces the Negative Effects of Automation Bias Overall but Increases False Negatives. *arXiv preprint*. https://arxiv.org/abs/2508.07617

Khalighi, S., Reddy, K. M., Midya, A., et al. (2024). Artificial intelligence in neuro-oncology: advances and challenges in brain tumor diagnosis, prognosis, and precision treatment. *npj Precision Oncology*. https://doi.org/10.1038/s41698-024-00575-0

Kim, Y., Park, C., Jeong, H., Chan, Y. S., Xu, X., McDuff, D., et al. (2024). MDAgents: An Adaptive Collaboration of LLMs for Medical Decision-Making. *NeurIPS 2024*. https://arxiv.org/abs/2404.15155

Kompa, B., Snoek, J., & Beam, A. L. (2021). Second opinion needed: communicating uncertainty in medical machine learning. *npj Digital Medicine*. https://doi.org/10.1038/s41746-020-00367-3

Koutsoubis, N., Waqas, A., Yilmaz, Y., Ramachandran, R. P., Schabath, M., & Rasool, G. (2024). Future-Proofing Medical Imaging with Privacy-Preserving Federated Learning and Uncertainty Quantification: A Review. *arXiv preprint*. https://arxiv.org/abs/2409.16340

Lakshminarayanan, B., Pritzel, A., & Blundell, C. (2017). Simple and Scalable Predictive Uncertainty Estimation using Deep Ensembles. *NeurIPS 2017*. https://arxiv.org/abs/1612.01474

Lekadir, K., Frangi, A. F., Porras, A. R., et al. (2025). FUTURE-AI: international consensus guideline for trustworthy and deployable artificial intelligence in healthcare. *BMJ*. https://doi.org/10.1136/bmj-2024-081554

Li, X., Jiang, M., Xu, G., Xia, J., Chiu, C., Jia, M., et al. (2026). CRC-Router: Risk-Constrained Routing for Medical Agentic AI Systems. *arXiv preprint*. https://arxiv.org/abs/2609.30714

Liu, F., Niu, Y., Zhang, Q., Wang, K., Dong, Z., Wong, I. N., et al. (2025). A foundational architecture for AI agents in healthcare. *Cell Reports Medicine*. https://doi.org/10.1016/j.xcrm.2025.102374

Liu, H., Wei, D., Lu, D., et al. (2023). M3AE: Multimodal Representation Learning for Brain Tumor Segmentation with Missing Modalities. *AAAI Conference on Artificial Intelligence*. https://doi.org/10.1609/aaai.v37i2.25253

Liu, Y., Zhang, Z., Liu, R., Xitang, M., Shin, S., & Lee, H. (2026). TumorBoard: Evidence-Grounded Multi-Agent Decision Support for Longitudinal Neuro-Oncology. *arXiv preprint*. https://arxiv.org/abs/2608.03190

Lou, J., Yang, Y., Yu, Z., Fu, Z., Han, W., Huang, Q., et al. (2026). CXRAgent: Director-Orchestrated Multi-Stage Reasoning for Chest X-Ray Interpretation. *IEEE Transactions on Medical Imaging*. https://doi.org/10.1109/tmi.2026.3713585

Louis, D. N., Perry, A., Wesseling, P., et al. (2021). The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro-Oncology*, 23(8), 1231-1251. https://doi.org/10.1093/neuonc/noab106

Lu, M. Y., Williamson, D. F. K., Chen, T. Y., Chen, R. J., Barbieri, M., & Mahmood, F. (2021). Data-efficient and weakly supervised computational pathology on whole-slide images (CLAM). *Nature Biomedical Engineering*. https://arxiv.org/abs/2004.09666

Malone, T. W., Vaccaro, M. A., Campero, A., et al. (2024). When combinations of humans and AI are useful: A systematic review and meta-analysis. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-024-02024-1

Manthe, M., Duffner, S., & Lartizien, C. (2024). Federated brain tumor segmentation: An extensive benchmark. *Medical Image Analysis*. https://doi.org/10.1016/j.media.2024.103270

Menze, B. H., Jakab, A., Bauer, S., et al. (2015). The multimodal brain tumor image segmentation benchmark (BRATS). *IEEE Transactions on Medical Imaging*, 34(10), 1993-2024. https://doi.org/10.1109/TMI.2014.2377694

Mitchell, M., Wu, S., Zaldivar, A., Barnes, P., Vasserman, L., Hutchinson, B., et al. (2019). Model Cards for Model Reporting. *ACM FAT* 2019*. https://arxiv.org/abs/1810.03993

Molinaro, A. M., Hervey-Jumper, S., Morshed, R. A., et al. (2020). Association of maximal extent of resection of contrast-enhanced and non-contrast-enhanced tumor with survival within molecular subgroups of patients with newly diagnosed glioblastoma. *JAMA Oncology*, 6(4), 495-503. https://doi.org/10.1001/jamaoncol.2019.6143

Mozannar, H., & Sontag, D. (2020). Consistent Estimators for Learning to Defer to an Expert. *ICML 2020*. https://arxiv.org/abs/2006.01862

National Cancer Control Programme. (2024). National cancer incidence and mortality data, Sri Lanka. *Ministry of Health*, Sri Lanka. https://www.nccp.health.gov.lk/

Olawade, D. B., Plabon, S. B., Ojo, A., Ogunbona, M. A., Makanjuola, B. D., & Olasilola, O. R. (2026). Human in the loop artificial intelligence in healthcare: applications, outcomes, and implementation challenges. *International Journal of Medical Informatics*, 213, 106362. https://doi.org/10.1016/j.ijmedinf.2026.106362

Ovadia, Y., Fertig, E., Ren, J., Nado, Z., Sculley, D., Nowozin, S., et al. (2019). Can You Trust Your Model's Uncertainty? Evaluating Predictive Uncertainty Under Dataset Shift. *NeurIPS 2019*. https://arxiv.org/abs/1906.02530

Pati, S., Baid, U., Edwards, B., Sheller, M., Wang, S. H., Bakas, S., et al. (2022). Federated learning enables big data for rare cancer boundary detection. *Nature Communications*. https://arxiv.org/abs/2204.10836

Price, M., Ostrom, Q. T., et al. (2024). CBTRUS statistical report: primary brain and other central nervous system tumors diagnosed in the United States in 2017-2021. *Neuro-Oncology*, 26(Suppl. 6), vi1-vi85. https://doi.org/10.1093/neuonc/noae145

Sanai, N., & Berger, M. S. (2008). Glioma extent of resection and its impact on patient outcome. *Neurosurgery*, 62(4), 753-766. https://doi.org/10.1227/01.neu.0000318159.21731.cf

Saueressig, C., Scholz, D., Raffler, P., Delbridge, C., et al. (2026). Multimodal fusion of pathology and radiology foundation models for WHO 2021 glioma subtyping. *npj Precision Oncology*. https://doi.org/10.1038/s41698-026-01366-5

Sendak, M. P., Gao, M., Brajer, N., & Balu, S. (2020). Presenting machine learning model information to clinical end users with model facts labels. *npj Digital Medicine*, 3, 41. https://doi.org/10.1038/s41746-020-0253-3

Sounderajah, V., Guni, A., Liu, X., et al. (2025). The STARD-AI reporting guideline for diagnostic accuracy studies using artificial intelligence. *Nature Medicine*. https://doi.org/10.1038/s41591-025-03953-8

Tang, Y., Yang, D., Li, W., Roth, H. R., Landman, B., Xu, D., et al. (2022). Self-Supervised Pre-Training of Swin Transformers for 3D Medical Image Analysis. *CVPR 2022*. https://arxiv.org/abs/2111.14791

Topol, E. J. (2019). High-performance medicine: the convergence of human and artificial intelligence. *Nature Medicine*, 25, 44-56. https://doi.org/10.1038/s41591-018-0300-7

Vasey, B., Nagendran, M., Campbell, B., Clifton, D. A., Collins, G. S., McCulloch, P., et al. (2022). Reporting guideline for the early-stage clinical evaluation of decision support systems driven by artificial intelligence: DECIDE-AI. *Nature Medicine*. https://doi.org/10.1038/s41591-022-01772-9

Verma, R., & Nalisnick, E. (2022). Calibrated Learning to Defer with One-vs-All Classifiers. *ICML 2022*. https://arxiv.org/abs/2202.03673

Vermeulen, C., Pages-Gallego, M., Kester, L., Kranendonk, M. E. G., et al. (2023). Ultra-fast deep-learned CNS tumour classification during surgery. *Nature*. https://doi.org/10.1038/s41586-023-06615-2

Wang, W., Zhao, Y., Teng, L., et al. (2023). Neuropathologist-level integrated classification of adult-type diffuse gliomas using deep learning from whole-slide pathological images. *Nature Communications*. https://doi.org/10.1038/s41467-023-41195-9

World Health Organization. (2021). Ethics and governance of artificial intelligence for health: WHO guidance. *WHO*. https://www.who.int/publications/i/item/9789240029200

Yevudza, W. E., Buckman, V., Darko, K., et al. (2024). Neuro-oncology access in Sub-Saharan Africa: A literature review of challenges and opportunities. *Neuro-Oncology Advances*. https://doi.org/10.1093/noajnl/vdae057

Yu, F., Moehring, A., Banerjee, O., et al. (2024). Heterogeneity and predictors of the effects of AI assistance on radiologists. *Nature Medicine*. https://doi.org/10.1038/s41591-024-02850-w

Zech, J. R., Badgeley, M. A., Liu, M., Costa, A. B., Titano, J. J., & Oermann, E. K. (2018). Variable generalization performance of a deep learning model to detect pneumonia in chest radiographs: a cross-sectional study. *PLOS Medicine*, 15(11), e1002683. https://doi.org/10.1371/journal.pmed.1002683

Zenk, M., Baid, U., Pati, S., et al. (2025). Towards fair decentralized benchmarking of healthcare AI algorithms with the Federated Tumor Segmentation (FeTS) challenge. *Nature Communications*. https://doi.org/10.1038/s41467-025-60466-1

Zhang, Y., He, N., Yang, J., et al. (2022). mmFormer: Multimodal Medical Transformer for Incomplete Multimodal Learning of Brain Tumor Segmentation. *MICCAI 2022 (LNCS)*. https://doi.org/10.1007/978-3-031-16443-9_11

Zhu, Y., Gu, L., Wang, Z., Sang, H., Sui, D., Tang, W., et al. (2025). Auditing medical multi-agent AI reveals risks of false consensus. *arXiv preprint*. https://arxiv.org/abs/2510.10185