Assignment 8 CS374

Harsh Patel(201701021) Viraj Patel(201701439)

Assigned by:

Prof. Arnab Kumar

October 3, 2019

Contents

| | bic Spline Interpolation |
|-----|---|
| 1.1 | 1 |
| | 1.1.1 Graph |
| Cu | bic Spline Interpolation and Quadratic Lagrange In- |
| | polation |
| 2.1 | • |
| 2.2 | |
| | 2.2.1 Graph |
| Nıı | merical Integration and Differentiation |
| 3.1 | Q1 |
| | 3.1.1 Equation |
| | 3.1.2 Table |
| 3.2 | |
| | 3.2.1 Equation |
| | 3.2.2 Table |
| 3.3 | Q3 |
| | 3.3.1 Equation |
| | 3.3.2 Table |
| 3.4 | |
| | 3.4.1 Table |
| 3.5 | Q5 |
| | 3.5.1 Equation |
| | 3.5.2 Table |
| For | rward and Central Difference at x=1 |
| 4.1 | |
| | 4.1.1 Equation |
| | 4.1.2 Table |
| | 4.1.2 Table |

| 4.2 | Q2 . | | | | | | | | | | | | 13 |
|-----|-------|----------|----|--|--|--|--|--|--|--|--|--|----|
| | 4.2.1 | Equation | ١. | | | | | | | | | | 13 |
| | 4.2.2 | Table | | | | | | | | | | | 13 |

1 Cubic Spline Interpolation

| X | 0 | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|--------|--------|--------|--------|--------|--------|
| У | 2 | 2.1592 | 3.1697 | 5.4332 | 9.1411 | 14.406 | 21.303 |

1.1 Equation

$$S''(x) = M_j \tag{1}$$


$$S(x) = \frac{M_{j-1}}{x_j - x_{j-1}} \frac{(x_j - x)^3}{6} + \frac{M_j}{x_j - x_{j-1}} \frac{(x - x_{j-1})^3}{6} + Ax + B$$
 (2)

$$A = D - C \tag{3}$$

$$B = Cx_j - Dx_{j-1} \tag{4}$$

$$\frac{M_{j-1}}{6}(x_j - x_{j-1}) + \frac{M_j}{3}(x_{j+1} - x_{j-1}) + \frac{M_{j+1}}{6}(x_{j+1} - x_j) = \frac{y_{j+1} - y_j}{x_{j+1} - x_j} + \frac{y_j - y_{j-1}}{x_j - x_{j-1}}$$
(5)

1.1.1 Graph



Plot of Cubic Spline Interpolation

2 Cubic Spline Interpolation and Quadratic Lagrange Interpolation

2.1 Equation

$$S''(x) = M_j \tag{6}$$

$$S(x) = \frac{M_{j-1}}{x_j - x_{j-1}} \frac{(x_j - x)^3}{6} + \frac{M_j}{x_j - x_{j-1}} \frac{(x - x_{j-1})^3}{6} + Ax + B$$
 (7)

$$A = D - C \tag{8}$$


$$B = Cx_i - Dx_{i-1} \tag{9}$$

$$\frac{M_{j-1}}{6}(x_j - x_{j-1}) + \frac{M_j}{3}(x_{j+1} - x_{j-1}) + \frac{M_{j+1}}{6}(x_{j+1} - x_j) = \frac{y_{j+1} - y_j}{x_{j+1} - x_j} + \frac{y_j - y_{j-1}}{x_j - x_{j-1}}$$
(10)

$$y = \frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)}y_0 + \frac{(x-x_0)(x-x_2)}{(x_1-x_0)(x_1-x_2)}y_1 + \frac{(x-x_0)(x-x_1)}{(x_2-x_0)(x_2-x_1)}y_2$$
(11)

2.2 Graph

2.2.1 Graph



Plot of Cubic Spline Interpolation and Quadratic Lagrange Interpolation

3 Numerical Integration and Differentiation

3.1 Q1

3.1.1 Equation

$$\int_0^{\pi} e^x \cos 4x = \frac{e^{\pi} - 1}{17} \tag{12}$$

3.1.2 Table

| n | h | Tn | Sn | TnError | SnError |
|-----|----------|-----------|-----------|-----------|--------------|
| 2 | 1.570796 | 26.516336 | 22.715077 | 25.213942 | 2.141268e+01 |
| 4 | 0.785398 | 3.249050 | -4.506711 | 1.946657 | 5.809105e+00 |
| 8 | 0.392699 | 1.624525 | 1.083017 | 0.322132 | 2.193769e-01 |
| 16 | 0.196350 | 1.375723 | 1.292788 | 0.073329 | 9.605410e-03 |
| 32 | 0.098175 | 1.320312 | 1.301842 | 0.017918 | 5.520189e-04 |
| 64 | 0.049087 | 1.306848 | 1.302360 | 0.004454 | 3.379643e-05 |
| 128 | 0.024544 | 1.303506 | 1.302392 | 0.001112 | 2.101451e-06 |
| 256 | 0.012272 | 1.302672 | 1.302394 | 0.000278 | 1.311722e-07 |
| 512 | 0.006136 | 1.302463 | 1.302394 | 0.000069 | 8.195633e-09 |

3.2 Q2

3.2.1 Equation

$$\int_0^1 x^{5/2} = 2/7 \tag{13}$$

3.2.2 Table

| n | h | Tn | Sn | TnError | SnError |
|-----|----------|----------|----------|--------------|--------------|
| 2 | 0.500000 | 0.338388 | 0.284518 | 5.267406e-02 | 1.196489e-03 |
| 4 | 0.250000 | 0.298791 | 0.285593 | 1.307721e-02 | 1.217400e-04 |
| 8 | 0.125000 | 0.288975 | 0.285702 | 3.260454e-03 | 1.179823e-05 |
| 16 | 0.062500 | 0.286529 | 0.285713 | 8.142822e-04 | 1.108410e-06 |
| 32 | 0.031250 | 0.285918 | 0.285714 | 2.034940e-04 | 1.020813e-07 |
| 64 | 0.015625 | 0.285765 | 0.285714 | 5.086654e-05 | 9.279914e-09 |
| 128 | 0.007812 | 0.285727 | 0.285714 | 1.271601e-05 | 8.363088e-10 |
| 256 | 0.003906 | 0.285717 | 0.285714 | 3.178946e-06 | 7.492457e-11 |
| 512 | 0.001953 | 0.285715 | 0.285714 | 7.947314e-07 | 6.685430e-12 |

3.3 Q3

3.3.1 Equation

$$\int_0^5 \frac{1}{1 + (x - \pi)^2} = \arctan(5 - \pi) + \arctan(\pi)$$
 (14)

3.3.2 Table

| n | h | Tn | Sn | TnError | SnError |
|-----|----------|----------|----------|----------|--------------|
| 2 | 2.500000 | 2.166655 | 2.625095 | 0.173111 | 2.853292e-01 |
| 4 | 1.250000 | 2.268668 | 2.302672 | 0.071099 | 3.709437e-02 |
| 8 | 0.625000 | 2.332270 | 2.353471 | 0.007496 | 1.370512e-02 |
| 16 | 0.312500 | 2.337813 | 2.339660 | 0.001953 | 1.059309e-04 |
| 32 | 0.156250 | 2.339277 | 2.339765 | 0.000489 | 1.079988e-06 |
| 64 | 0.078125 | 2.339644 | 2.339766 | 0.000122 | 6.743238e-08 |
| 128 | 0.039062 | 2.339736 | 2.339766 | 0.000031 | 4.216919e-09 |
| 256 | 0.019531 | 2.339759 | 2.339766 | 0.000008 | 2.635929e-10 |
| 512 | 0.009766 | 2.339764 | 2.339766 | 0.000002 | 1.647504e-11 |

3.4 Q4

$$\exp\left(-x^2\right) \qquad x\epsilon[0,10] \tag{15}$$

3.4.1 Table

| n | h | Tn | Sn | TnError | SnError |
|-----|----------|----------|----------|--------------|--------------|
| 4 | 2.500000 | 1.254826 | 0.839768 | 3.685992e-01 | 4.645874e-02 |
| 8 | 1.250000 | 0.889428 | 0.767629 | 3.201353e-03 | 1.185979e-01 |
| 16 | 0.625000 | 0.886227 | 0.885160 | 1.886302e-11 | 1.067118e-03 |
| 32 | 0.312500 | 0.886227 | 0.886227 | 1.110223e-16 | 6.287748e-12 |
| 64 | 0.156250 | 0.886227 | 0.886227 | 2.220446e-16 | 2.220446e-16 |
| 128 | 0.078125 | 0.886227 | 0.886227 | 2.220446e-16 | 0.000000e+00 |
| 256 | 0.039062 | 0.886227 | 0.886227 | 5.551115e-16 | 4.440892e-16 |
| 512 | 0.019531 | 0.886227 | 0.886227 | 7.771561e-16 | 2.220446e-16 |

3.5 Q5

3.5.1 Equation

$$\arctan\left(x^2+1\right) \qquad x\epsilon[0,2] \tag{16}$$

3.5.2 Table

| | n | h | Tn | Sn |
|----|---|----------|----------|----------|
| - | 4 | 0.500000 | 2.177450 | 2.174418 |
| | 8 | 0.250000 | 2.175061 | 2.174265 |
| 1 | 6 | 0.125000 | 2.174461 | 2.174261 |
| 3 | 2 | 0.062500 | 2.174311 | 2.174261 |
| 6 | 4 | 0.031250 | 2.174274 | 2.174261 |
| 12 | 8 | 0.015625 | 2.174264 | 2.174261 |
| 25 | 6 | 0.007812 | 2.174262 | 2.174261 |
| 51 | 2 | 0.003906 | 2.174261 | 2.174261 |

4 Forward and Central Difference at x=1

4.1 Q1

4.1.1 Equation

$$\arctan\left(x^2 - x + 1\right) \tag{17}$$

4.1.2 Table

| | h | ForwardDiff | CentralDiff | ErrorForward | ErrorCentral |
|---|---------|-------------|-------------|--------------|--------------|
| • | 0.10000 | 0.520855 | 0.495856 | 0.012500 | -0.001042 |
| | 0.05000 | 0.511460 | 0.498960 | 0.006250 | -0.000260 |
| | 0.02500 | 0.505990 | 0.499740 | 0.003125 | -0.000065 |
| | 0.01250 | 0.503060 | 0.499935 | 0.001563 | -0.000016 |
| | 0.00625 | 0.501546 | 0.499984 | 0.000781 | -0.000004 |

4.2 Q2

4.2.1 Equation

$$\arctan\left(100x^2 - 199x + 100\right)$$
 (18)

4.2.2 Table

| h | ForwardDiff | CentralDiff | ErrorForward | ErrorCentral |
|---------|-------------|-------------|--------------|--------------|
| 0.10000 | 3.409790 | 0.200294 | 2.487500 | -0.124792 |
| 0.05000 | 2.594051 | 0.390426 | 1.243750 | -0.031198 |
| 0.02500 | 1.675666 | 0.469776 | 0.621875 | -0.007799 |
| 0.01250 | 1.109328 | 0.492262 | 0.310938 | -0.001950 |
| 0.00625 | 0.808388 | 0.498054 | 0.155469 | -0.000487 |