

AN EXPLICIT FORMULA FOR THE LODAY ASSEMBLY

VIRGIL CHAN

ABSTRACT. We give an explicit description of the Loday assembly map on homotopy groups when restricted to a certain subgroup coming from the Atiyah-Hirzebruch spectral sequence. This proves and generalises a formula about the Loday assembly map on the first homotopy group written down in the survey article by Lück and Reich. Furthermore, we use this new result to prove the Loday assembly for the integral group ring of cyclic group of order 2 is a bijection on the second homotopy group, and is an injection when one considers n -fold direct sum of cyclic group of order 2.

CONTENTS

1. Introduction	1
1.1. Notations	3
2. The Setups	3
2.1. Loday's Pairing	3
2.2. Extending the Pairing	5
2.3. Relationship with the Classical Pairings in Algebraic K-theory	7
2.4. Definition of the Loday Assembly	10
3. The Main Theorem	10
3.1. A Subgroup of the Source of Loday Assembly	11
3.2. Statement of the Main Theorem and the Proof	12
3.3. Applications to the Second K-groups of Integral Group Rings	18
Index	24
References	25

1. INTRODUCTION

The general consensus is that computing homotopy groups is difficult. Hence, investigating algebraic K-theory of rings is not an easy task. This is why assembly map comes into play—to approximate homotopy theory with a generalised homology theory. In his dissertation [Lod76], Loday wrote down the first example of K-theoretic assembly map

$$\alpha_{\text{Loday}} : BG_+ \wedge \mathbb{K}_R^{\text{GW}} \rightarrow \mathbb{K}_{R[G]}^{\text{GW}}$$

(see [Lod76, Proposition 4.1.1 on page 356] or Definition 2.9 below). Here, the target $\mathbb{K}_R^{\text{GW}}$ is the Gersten-Wagoner non-connective algebraic K-theory spectrum of the ring R , and the domain represents the generalised homology theory of the classifying space BG of the group G with coefficients in $\mathbb{K}_R^{\text{GW}}$. Furthermore, he related the cokernel of the induced map on homotopy groups with the classical Whitehead groups defined by J. H. C. Whitehead in [Whi50] and Hatcher-Wagoner in [HW73] when $R = \mathbb{Z}$. Motivated by the connection between

the classical Novikov Conjecture and the rational injectivity of the L-theoretic assembly map, Hsiang cast the following conjecture in his ICM address:

Conjecture 1.1 (K-theoretic Novikov Conjecture, [Hsi84])

When R is regular, and G is torsion-free, the map

$$\pi_i(\alpha_{\text{Loday}}) \otimes \text{id}_{\mathbb{Q}} : \pi_i(BG_+ \wedge \mathbb{K}_R^{\text{GW}}) \otimes \mathbb{Q} \rightarrow K_i(R[G]) \otimes \mathbb{Q} \quad (1.0.1)$$

is injective for all i .

Attempts on answering Conjecture 1.1 often leads to the creation of new mathematics, most notably the Topological Cyclic Homology, which allowed Bökstedt-Hsiang-Madsen [BHM93, Theorem 9.13 on page 535] to prove Conjecture 1.1 when G is homologically finite. A stronger statement is also possible:

Conjecture 1.2 (Classical Farrell-Jones Conjecture)

When R is regular and G is torsion-free, the map

$$\pi_i(\alpha_{\text{Loday}}) : \pi_i(BG_+ \wedge \mathbb{K}_R^{\text{GW}}) \rightarrow K_i(R[G]) \quad (1.0.2)$$

is an isomorphism for all i .

This conjecture can be understood as a conceptual approach to compute algebraic K-theory of a group ring via a homology theory, but it has important implications for a range of topics, notably the Borel Conjecture concerning the topological rigidity of closed spherical manifolds, and the Kaplansky Conjecture about the idempotents in a group ring.

One might ask why torsion-free is a necessary condition in Conjecture 1.2. As stated before, the cokernel of

$$\pi_i(\alpha_{\text{Loday}}) : \pi_i(BG_+ \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}}) \rightarrow K_i(\mathbb{Z}[G])$$

can be identified with the Whitehead groups $\text{Wh}_i(G)$. So, non-vanishing $\text{Wh}_i(G)$ implies the non-surjectivity of $\pi_i(\alpha_{\text{Loday}})$. This point is known, for example, when $G = C_p$ is the cyclic group of prime order p [Coh73, 11.5 on page 45] for $i = 1$. What happens with non-injectivity?

Conjecture 1.3

For a regular ring R , is there a group G with torsion that makes $\pi_i(\alpha_{\text{Loday}})$ not injective for some i ?

Ullmann-Wu gives an example (the Klein four-group) where non-injectivity can occur when R is a finite field [UW17]. In their case, non-injectivity occurs at the second homotopy group. However, it is currently not known whether such an example exists $R = \mathbb{Z}$.

When $R = \mathbb{Z}$, Conjecture 1.3 is known to be false for $i = 0, 1$. See [LR05, Lemma 2 on page 709] for example. This paper contributes to Conjecture 1.2 by giving an explicit description of $\pi_i(\alpha_{\text{Loday}})$ when restricted onto a subgroup of $\pi_i(BG_+ \wedge \mathbb{K}_R^{\text{GW}})$ coming from the Atiyah-Hirzebruch spectral sequence (see Theorem 3.5), and to Conjecture 1.3 by investigating injectivity phenomenon of $\pi_i(\alpha_{\text{Loday}})$ for cyclic group of order 2 (see Example 3.8 and Example 3.11).

We organise this paper as follow. We begin by review the pairing map defined by Loday in Section 2, which plays a key role in defining the assembly map. The main theorem is stated in Section 3, with examples about cyclic group of order 2.

1.1. Notations. Throughout this paper, we assume all rings are unital, associative, and satisfy the Invariant Basis Property. I.e., $R^n \cong R^m$ if and only if $n = m$. We always work with a discrete group G , and denote by BG its classifying space obtained from the bar construction.

By a spectrum, we mean a sequence $\mathbb{E} = \{\mathbb{E}_n\}_{n=1}^\infty$ of topological spaces, together with continuous maps

$$f_n : \mathbb{S}^1 \wedge \mathbb{E}_n \rightarrow \mathbb{E}_{n+1}.$$

For a ring R , we will use $\mathbb{K}_R^{\text{GW}}$ to denote the Gersten-Wagoner non-connective algebraic K-theory spectrum [Ger72; Wag72], whose n -th space is given by

$$(\mathbb{K}_R^{\text{GW}})_n := K_0(\Sigma^n R) \times BGL(\Sigma^n R)^+,$$

where $\Sigma^n R$ is the n -fold suspension of R .

2. THE SETUPS

2.1. Loday's Pairing. Let R, S be rings, and fix an isomorphism

$$\xi : R^m \otimes S^n \rightarrow (R \otimes S)^{mn}. \quad (2.1.1)$$

Under this setting, the tensor product of matrices gives a group homomorphism

$$GL(m, R) \times GL(n, S) \rightarrow GL(mn, R \otimes S),$$

sending elementary matrices to elementary matrices. Hence we have an induced map

$$f_{m,n}^{R,S} : BGL(m, R)^+ \times BGL(n, S)^+ \rightarrow BGL(mn, R \otimes S)^+. \quad (2.1.2)$$

Write

$$i_{mn} : BGL(mn, R \otimes S)^+ \rightarrow BGL(R \otimes S)^+$$

to be map induced by the canonical inclusion

$$GL(mn, R \otimes S) \rightarrow GL(R \otimes S).$$

We then define

Definition 2.1 (The map $\gamma_{m,n}$, [Lod76, page 332])

The map

$$\gamma_{m,n} : BGL(m, R)^+ \times BGL(n, S)^+ \rightarrow BGL(R \otimes S)^+ \quad (2.1.3)$$

is defined by the formula:

$$\gamma_{m,n}(x, y) := i_{mn} \circ f_{m,n}^{R,S}(x, y) - i_{mn} \circ f_{m,n}^{R,S}(x, y_0) - i_{mn} \circ f_{m,n}^{R,S}(x_0, y). \quad (2.1.4)$$

Here, the minus sign on the RHS comes from the H-group structure of $BGL(R \otimes S)^+$; and x_0 (resp. y_0) is the base-point in $BGL(m, R)^+$ (resp. $BGL(n, S)^+$). (I.e., represented by the $m \times m$ identity matrix.)

The choice of the isomorphism ξ says the map γ is well-defined up to weak-homotopy.

One gets a map

$$\gamma : BGL(R)^+ \times BGL(S)^+ \rightarrow BGL(R \otimes S)^+ \quad (2.1.5)$$

by letting $m, n \rightarrow \infty$. This map, again, is well-defined up to weak-homotopy. Moreover, it is homotopy trivial on the wedge product $BGL(R)^+ \vee BGL(S)^+$. So we have

Definition 2.2 (The Loday pairing map γ_{Loday} , [Lod76, Section 2.1.7 on page 333])

The map

$$\gamma_{\text{Loday}} : BGL(R)^+ \wedge BGL(S)^+ \rightarrow BGL(R \otimes S)^+ \quad (2.1.6)$$

is defined to be the filler of the following homotopy commutative diagram:

$$\begin{array}{ccc} BGL(R)^+ \times BGL(S)^+ & \xrightarrow{\gamma} & BGL(R \otimes S)^+ \\ \text{proj} \downarrow & \nearrow \gamma_{\text{Loday}} & \\ BGL(R)^+ \wedge BGL(S)^+ & & \end{array} \quad (2.1.7)$$

This allows us to define a multiplicative structure on algebraic K-theory.

Definition 2.3 (Loday product $\star$, [Lod76, Section 2.1.10 on page 335])

For each integers $i, j \geq 1$, we define the product

$$\star : K_i(R) \otimes K_j(S) \rightarrow K_{i+j}(R \otimes S) \quad (2.1.8)$$

by

$$[f] \star [g] := [\gamma_{\text{Loday}} \circ (f \circ g)], \quad (2.1.9)$$

for which f and g are spheroids

$$\begin{aligned} f : \mathbb{S}^i &\rightarrow BGL(R)^+, \\ g : \mathbb{S}^j &\rightarrow BGL(S)^+. \end{aligned}$$

Proposition 2.4 (Properties of the Loday Product $\star$)

The product map $\star$ is

- (i) ([Lod76, Theorem 2.1.11 on page 335]) natural in R and S , associative, bilinear,
- (ii) ([Lod76, Theorem 2.1.12 on page 335]) graded commutative:

$$[f] \star [g] = (-1)^{ij} ([g] \star [f]) \quad (2.1.10)$$

whenever $[f] \in K_i(R)$ and $[g] \in K_j(S)$.

- (iii) ([Lod76, Proposition 2.2.3 on page 337]) When $i = j = 1$, the product $\{u\} \star \{v\}$ is the inverse of the Steinberg symbol:

$$\{u\} \star \{v\} = -\{u, v\}_{\text{St}}, \quad (2.1.11)$$

where we write the Abelian group $K_2(R \otimes S)$ additively.

2.2. Extending the Pairing. We want to modify the Loday pairing map

$$\gamma_{\text{Loday}} : BGL(R)^+ \wedge BGL(S)^+ \rightarrow BGL(R \otimes S)^+$$

to a map

$$\gamma'_{\text{Loday}} : K_R^+ \wedge K_S^+ \rightarrow K_{R \otimes S}^+.$$

for which

$$K_R^+ := K_0(R) \times BGL(R)^+, \quad (2.2.1)$$

so that one has an extended external product

$$K_p(R) \otimes K_q(S) \rightarrow K_{p+q}(R \otimes S)$$

for all $p, q \geq 0$. Moreover, this external product should be coherent with the isomorphism

$$K_p(R) \xrightarrow{\cong} K_{p+1}(\Sigma R) \quad (2.2.2)$$

promised by the **Gersten-Wagoner delooping** [Ger72; Wag72]:

$$K_R^+ \simeq \Omega BGL(\Sigma R)^+. \quad (2.2.3)$$

This is the place where we need the close form of the isomorphism in Equation (2.2.2):

$$\begin{cases} [f] \mapsto [f] \star \{\tau\} & \text{if } p \geq 1 \text{ ([Lod76, Corollary 2.3.6])}, \\ [P] \mapsto [P] \# \{\tau\} := \{p \otimes \tau + (1-p) \otimes 1\} & \text{if } p = 0 \text{ ([Lod76, page 328])}, \end{cases} \quad (2.2.4)$$

where

(i) the element $\tau \in GL(\Sigma \mathbb{Z})$ is given by

$$\tau := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & \cdots \\ 1 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 1 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}; \quad (2.2.5)$$

(ii) and p is the projection operator associated to the finitely generated projective R -modules P .

Equally important, the extended external product should be related to the classical product

$$\# : K_p(R) \otimes K_q(S) \rightarrow K_{p+q}(R \otimes S) \quad (2.2.6)$$

defined by Milnor [Mil72] for $p, q \geq 0$, and $p + q \leq 2$.

There is an obvious way to do so—tensor product of modules gives a homomorphism

$$K_0(R) \otimes K_0(S) \rightarrow K_0(R \otimes S).$$

So one might use this to extend the pairing map

$$\gamma_{\text{Loday}} : BGL(R)^+ \wedge BGL(S)^+ \rightarrow BGL(R \otimes S)^+$$

by

$$\begin{aligned} (K_0(R) \times BGL(R)^+) \wedge (K_0(S) \times BGL(S)^+) &\rightarrow K_0(R \otimes S) \times BGL(R \otimes S)^+ \\ ([P], x) \wedge ([Q], y) &\mapsto ([P \otimes Q], \gamma_{\text{Loday}}(x, y)). \end{aligned}$$

This does not work well, because the induced product map

$$K_0(R) \otimes K_i(S) \rightarrow K_i(R \otimes S)$$

is the zero map for all $i > 0$.

Here, we propose a method to extend the original pairing and show this method might be a correct one.

Definition 2.5 (The modified Loday pairing γ'_{Loday})

The map

$$\gamma'_{\text{Loday}} : K_R^+ \wedge K_S^+ \rightarrow K_{R \otimes S}^+$$

is defined to be the filler of the following homotopy commutative diagram:

$$\begin{array}{ccc} K_R^+ \wedge K_S^+ & \xrightarrow{\gamma'_{\text{Loday}}} & K_{R \otimes S}^+ \\ \simeq \downarrow & & \downarrow \simeq \\ \Omega BGL(\Sigma R)^+ \wedge \Omega BGL(\Sigma S)^+ & \xrightarrow{\Omega^2 \gamma_{\text{Loday}}} & \Omega^2 BGL(\Sigma^2(R \otimes S))^+. \end{array} \quad (2.2.7)$$

Furthermore, we define a new product map

$$\star' : K_i(R) \otimes K_j(S) \rightarrow K_{i+j}(R \otimes S), \quad (2.2.8)$$

for all $i, j \geq 0$ by

$$[f] \star' [g] := [\gamma'_{\text{Loday}} \circ (f \circ g)], \quad (2.2.9)$$

The two products are related in the following way:

Proposition 2.6

For each $i, j \geq 1$, if $[f] \in K_i(R)$ and $[g] \in K_j(S)$, then

$$[f] \star' [g] = [f] \star ((-1)^j [g]). \quad (2.2.10)$$

In particular, for $\{u\} \in K_1(R)$ and $\{v\} \in K_1(S)$, the $\star'$ -product is given by the Steinberg symbol

$$\{u\} \star' \{v\} = \{u, v\}_{\text{St}}. \quad (2.2.11)$$

Proof. We consider the following commutative diagram

$$\begin{array}{ccc} K_i(R) \otimes K_j(S) & \xrightarrow{\star'} & K_{i+j}(R \otimes S) \\ \downarrow \cong & & \downarrow \cong \\ (-\star \{\tau\}) \otimes (-\star \{\tau\}) & & (-\star \{\tau\}) \star \{\tau\} \\ \downarrow & & \downarrow \\ K_{i+1}(\Sigma R) \otimes K_{j+1}(\Sigma S) & \xrightarrow{\star} & K_{i+j+2}(\Sigma^2(R \otimes S)). \end{array} \quad (2.2.12)$$

If $[f] \in K_i(R)$ and $[g] \in K_j(S)$, then the lower part of the diagram gives

$$\begin{aligned} ([f] \star \{\tau\}) \star ([g] \star \{\tau\}) &= [f] \star (\{\tau\} \star [g]) \star \{\tau\} && \text{(associativity)} \\ &= [f] \star ((-1)^j ([g] \star \{\tau\})) \star \{\tau\} && \text{(graded-commutativity)} \\ &= ([f] \star ((-1)^j [g]) \star \{\tau\}) \star \{\tau\}. && \text{(associativity)} \end{aligned}$$

So we must have

$$[f] \star' [g] = [f] \star ((-1)^j [g]).$$

□

2.3. Relationship with the Classical Pairings in Algebraic K-theory. Milnor defined multiplication structure on lower K-groups

$$\# : K_i(R) \otimes K_j(R) \rightarrow K_{i+j}(R \otimes R') \quad (2.3.1)$$

for $i, j \geq 0$, and $i + j \leq 2$ in [Mil72]. Actually, Milnor only defined the multiplication internally (i.e., when $R = R'$). But one can mimic his definition to obtain an external product as in Equation (2.3.1), so that when $R = R'$ it becomes Milnor's version.

The multiplication in Equation (2.3.1) is given by the formula:

$$\left\{ \begin{array}{ll} [P] \# [Q] &= [P \otimes Q] && \text{if } i = j = 0, \\ [P] \# \{u\} &= \{p \otimes u + (1 - p) \otimes 1\} && \text{if } i = 0, \text{ and } j = 1, \\ \{u\} \# \{v\} &= \{u, v\}_{\text{St}} && \text{if } i = j = 1. \end{array} \right. \quad (2.3.2)$$

We omit the case $i = 0, j = 2$ here. The main tool we need is

Proposition 2.7 ([Mil72, Lemma 8.9 on page 70])

The multiplication $\#$ in (2.3.1) is associative and bilinear for $i, j > 0$, and $i + j \leq 2$.

It is also straight forward to verify that $\#$ is graded-commutative. We now relate $\#$ and $\star'$.

Theorem 2.8

For every $[P] \in K_0(R)$, $[Q] \in K_0(R')$, $\{u\} \in K_1(R)$, and $\{v\} \in K_1(R')$, we have

- (i) $[P] \star' [Q] = [P] \# [Q]$,
- (ii) $[P] \star' \{v\} = [P] \# (-\{v\})$,
- (iii) $\{u\} \star' \{v\} = \{u\} \# \{v\}$.

The author, however, does not know if the product

$$\star' : K_0(R) \otimes K_2(S) \rightarrow K_2(R \otimes S) \quad (2.3.3)$$

is compatible with Milnor's product

$$\# : K_0(R) \otimes K_2(S) \rightarrow K_2(R \otimes S). \quad (2.3.4)$$

defined in [Mil72, page 51].

Proof. We refer readers to Equation (2.2.4), and the commutative diagram in Equation (2.2.12).

(i) When $i = j = 0$, the diagram in (2.2.12) becomes

$$\begin{array}{ccc} K_0(R) \otimes K_0(R') & \xrightarrow{\star'} & K_0(R \otimes R') \\ (-\# \{\tau\}) \otimes (-\# \{\tau\}) \cong & & \cong ((-\# \{\tau\}) \star \{\tau\}) \\ \downarrow & & \downarrow \\ K_1(\Sigma R) \otimes K_1(\Sigma R') & \xrightarrow{\star} & K_2(\Sigma^2(R \otimes R')). \end{array} \quad (2.3.5)$$

We then compute

$$\begin{aligned} ([P] \# \{\tau\}) \star ([Q] \# \{\tau\}) &= -\{[P] \star' \{\tau\}, [Q] \# \{\tau\}\}_{\text{St}} \\ &= \{[P] \# \{\tau\}, (-[Q]) \# \{\tau\}\}_{\text{St}} && (\text{Bilinearity of } \{-, -\}_{\text{St}}) \\ &= ([P] \# \{\tau\}) \# ((-[Q]) \# \{\tau\}) \\ &= [P] \# (\{\tau\} \# (-[Q])) \# \{\tau\} && (\text{Associativity of } \#) \\ &= [P] \# (-[Q] \# \{\tau\}) \# \{\tau\} && (\text{Graded-commutativity of } \#) \\ &= -([P] \# ([Q] \# \{\tau\}) \# \{\tau\}) && (\text{Bilinearity of } \#) \\ &= -\{[P] \# ([Q] \# \{\tau\}), \{\tau\}\}_{\text{St}} \\ &= ([P] \# ([Q] \# \{\tau\})) \star \{\tau\} \end{aligned}$$

$$= ([P] \# [Q]) \# \{\tau\} \star \{\tau\}, \quad (\text{Associativity of } \#)$$

proving

$$[P] \star' [Q] = [P] \# [Q].$$

(ii) When $i = 0$, $j = 1$, Diagram (2.2.12) becomes

$$\begin{array}{ccc} K_0(R) \otimes K_1(R') & \xrightarrow{\star'} & K_1(R \otimes R') \\ (-\# \{\tau\}) \otimes (-\star \{\tau\}) \downarrow \cong & & \cong \downarrow ((-\star \{\tau\}) \star \{\tau\}) \\ K_1(\Sigma R) \otimes K_2(\Sigma R') & \xrightarrow{\star} & K_3(\Sigma^2(R \otimes R')). \end{array} \quad (2.3.6)$$

We then compute

$$\begin{aligned} ([P] \# \{\tau\}) \star (\{v\} \star \{\tau\}) &= (([P] \# \{\tau\}) \star \{v\}) \star \{\tau\} && (\text{Associativity of } \star) \\ &= (-\{[P] \# \{\tau\}, \{v\}\}_{\text{st}}) \star \{\tau\} && (\text{Proposition 2.4}) \\ &= (-([P] \# \{\tau\}) \# \{v\}) \star \{\tau\} && (\text{Equation (2.3.2)}) \\ &= (-([P] \# (\{\tau\} \# \{v\}))) \star \{\tau\} && (\text{Associativity of } \#) \\ &= ([P] \# (\{u\} \# \{\tau\})) \star \tau && (\text{Graded-commutativity of } \#) \\ &= (([P] \# \{u\}) \# \{\tau\}) \star \{\tau\} && (\text{Associativity of } \star) \\ &= \{[P] \# \{v\}, \{\tau\}\}_{\text{st}} \star \{\tau\} && (\text{Equation (2.3.2)}) \\ &= (-([P] \# \{v\}) \star \{\tau\}) \star \{\tau\} && (\text{Proposition 2.4}) \\ &= ((-[P] \# \{v\}) \star \{\tau\}) \star \{\tau\} && (\text{Bilinearity of } \star) \\ &= (([P] \# (-\{v\})) \star \{\tau\}) \star \{\tau\} && (\text{Bilinearity of } \#), \end{aligned}$$

proving

$$[P] \star' \{v\} = [P] \# (-\{v\}).$$

(iii)

$$\{u\} \star' \{v\} = \{u\} \star (-\{v\}) \quad (\text{Proposition 2.6})$$

$$= -(\{u\} \star \{v\}) \quad (\text{Bilinearity of } \star)$$

$$= \{\{u\}, \{v\}\}_{\text{St}} \quad (\text{Proposition 2.4})$$

$$= \{u\} \# \{v\}.$$

□

2.4. Definition of the Loday Assembly. We use the modified Loday pairing γ'_{Loday} to define the Loday assembly map:

Definition 2.9 (The Loday Assembly Map α_{Loday})

Let R be a ring, and G be a group. The **Loday assembly map** is a map of spectra

$$\alpha_{\text{Loday}} : BG_+ \wedge \mathbb{K}_R^{\text{GW}} \rightarrow \mathbb{K}_{R[G]}^{\text{GW}}, \quad (2.4.1)$$

whose components are given by the composition of maps of spaces:

$$BG_+ \wedge K_{\Sigma^m R}^+ \xrightarrow{j^+ \wedge \text{id}} K_{\mathbb{Z}[G]_+}^+ \wedge K_{\Sigma^m R}^+ \xrightarrow{\gamma'_{\text{Loday}}} K_{\Sigma^m R[G]}^+. \quad (2.4.2)$$

Here, the map

$$j^+ : BG_+ \rightarrow K_{\mathbb{Z}[G]}^+ \quad (2.4.3)$$

is induced by the group homomorphism

$$j : G \rightarrow GL(\mathbb{Z}[G])$$

$$g \mapsto \begin{bmatrix} g & 0 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ 0 & 0 & 1 & 0 & \cdots \\ 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}. \quad (2.4.4)$$

3. THE MAIN THEOREM

The source of the Loday assembly

$$\alpha_{\text{Loday}} : \pi_{i+1}(BG_+ \wedge \mathbb{K}_R^{\text{GW}}) \rightarrow K_{i+1}(R[G]) \quad (3.0.1)$$

on homotopy groups can be computed by the Atiyah-Hirzebruch spectral sequence (AHSS):

$$\begin{aligned} \text{AHSS}(BG)_{p,q}^2 &\cong H_p(BG; K_q(R)) \\ &\Rightarrow \pi_{p+q}(BG_+ \wedge \mathbb{K}_R^{\text{GW}}), \end{aligned} \quad (3.0.2)$$

and in particular,

$$\begin{aligned} E_{1,i}^2 &\cong H_1(BG; K_i(R)) \\ &\cong G_{ab} \otimes K_i(R). \end{aligned} \quad (3.0.3)$$

Because the spectral sequence (3.0.3) is concentrated on the right-half plane, it follows that the differential

$$d_{1,i}^2 : E_{1,i}^2 \rightarrow E_{-1,i+1}^2 \quad (3.0.4)$$

has to be zero. So the group $E_{1,i}^\infty$ is a quotient of $E_{1,i}^2$, and we shall see it is also a subgroup of the source $\pi_{i+1}(BG_+ \wedge \mathbb{K}_R^{\text{GW}})$.

3.1. A Subgroup of the Source of Loday Assembly. Fix a spectrum $\mathbb{E}$. For any CW complex X , we have a homeomorphism

$$X_+ \cong X \vee \mathbb{S}^0, \quad (3.1.1)$$

and hence an isomorphism of homotopy groups

$$\pi_{i+1}(X_+ \wedge \mathbb{E}) \cong \pi_{i+1}(X \wedge \mathbb{E}) \oplus \pi_{i+1}(\mathbb{S}^0 \wedge \mathbb{E}). \quad (3.1.2)$$

Fix a skeletal filtration

$$\text{point} = X^{(0)} \subseteq X^{(1)} \subseteq \cdots \subseteq X \quad (3.1.3)$$

on X , the identification in Equation (3.1.1) gives a skeletal filtration

$$X_+^{(p)} \cong X^{(p)} \vee \mathbb{S}^0 \quad (3.1.4)$$

on X_+ . Thus we have a splitting for the AHSS

$$\text{AHSS}(X_+)^r_{p,q} \cong \text{AHSS}(X)^r_{p,q} \oplus \text{AHSS}(\mathbb{S}^0)^r_{p,q} \quad (3.1.5)$$

for $r \geq 1$. Tracing through the construction of the AHSS (see, for example [Wic15]), we see that the splitting in Equation 3.1.1 respects the filtration on homotopy groups:

$$\mathcal{F}_1[\pi_{i+1}(X_+ \wedge \mathbb{E})] \cong \mathcal{F}_1[\pi_{i+1}(X \wedge \mathbb{E})] \oplus \mathcal{F}_1[\pi_{i+1}(\mathbb{S}^0 \wedge \mathbb{E})], \quad (3.1.6)$$

for which

$$\mathcal{F}_1[\pi_{i+1}(X_+ \wedge \mathbb{E})] := \text{im} \left(\pi_{i+1} \left(X_+^{(1)} \wedge \mathbb{E} \right) \rightarrow \pi_{i+1}(X_+ \wedge \mathbb{E}) \right). \quad (3.1.7)$$

The subgroup

$$\mathcal{F}_1[\pi_{i+1}(X_+ \wedge \mathbb{E})] \subseteq \pi_{i+1}(X_+ \wedge \mathbb{E}) \quad (3.1.8)$$

is what we are interested in, and it can be identified as:

$$\begin{aligned} \mathcal{F}_1[\pi_{i+1}(X_+ \wedge \mathbb{E})] &\cong \mathcal{F}_1[\pi_{i+1}(X \wedge \mathbb{E})] \oplus \mathcal{F}_1[\pi_{i+1}(\mathbb{S}^0 \wedge \mathbb{E})] \\ &= \text{AHSS}(X)_{1,i}^\infty \oplus \pi_{i+1}(\mathbb{E}). \end{aligned} \quad (3.1.9)$$

3.2. Statement of the Main Theorem and the Proof. Now, let us work with $\mathbb{E} = \mathbb{K}_R^{\text{GW}}$, and $X = BG$. Then Equation (3.1.9) says

$$\begin{aligned} \mathcal{F}_1 [\pi_{i+1}(BG_+ \wedge \mathbb{K}_R^{\text{GW}})] &\cong \mathcal{F}_1 [\pi_{i+1}(BG \wedge \mathbb{K}_R^{\text{GW}})] \oplus \mathcal{F}_1 [\pi_{i+1}(\mathbb{S}^0 \wedge \mathbb{K}_R^{\text{GW}})] \\ &= \text{AHSS}(BG)_{1,i}^\infty \oplus \pi_{i+1}(\mathbb{K}_R^{\text{GW}}) \\ &= \text{AHSS}(BG)_{1,i}^\infty \oplus K_{i+1}(R). \end{aligned} \quad (3.2.1)$$

So the source of the Loday assembly

$$\alpha_{\text{Loday}} : \pi_{i+1}(BG_+ \wedge \mathbb{K}_R^{\text{GW}}) \rightarrow K_{i+1}(R[G])$$

contains

$$\text{AHSS}(BG)_{1,i}^\infty \oplus K_{i+1}(R)$$

as a subgroup. In fact, we already know what the Loday assembly is doing when restricted onto the summand $K_{i+1}(R)$ —it is induced by the inclusion

$$R \rightarrow R[G]$$

of rings. See [Lod76, before Lemma 4.1.2 on page 356]. What we want now is to study the Loday assembly when restricted onto the summand $\text{AHSS}(BG)_{1,i}^\infty$.

The splitting of the AHSS in Equation (3.1.5) says the differentials of $\text{AHSS}(BG_+)^2_{p,q}$ splits into two parts, which we illustrate here:

$$\begin{array}{ccc} \begin{array}{c} \uparrow q \\ \begin{array}{cccc} K_5(R) & H_1(BG; K_5(R)) & H_2(BG; K_5(R)) & \cdots \\ K_4(R) & H_1(BG; K_4(R)) & H_2(BG; K_4(R)) & \cdots \\ K_3(R) & H_1(BG; K_3(R)) & H_2(BG; K_3(R)) & \cdots \\ K_2(R) & H_1(BG; K_2(R)) & H_2(BG; K_2(R)) & \cdots \\ K_1(R) & H_1(BG; K_1(R)) & H_2(BG; K_1(R)) & \cdots \\ K_0(R) & H_1(BG; K_0(R)) & H_2(BG; K_0(R)) & \cdots \end{array} \\ \rightarrow p \end{array} & \cong & \begin{array}{c} \uparrow q \\ \begin{array}{cccc} 0 & H_1(BG; K_5(R)) & H_2(BG; K_5(R)) & \cdots \\ 0 & H_1(BG; K_4(R)) & H_2(BG; K_4(R)) & \cdots \\ 0 & H_1(BG; K_3(R)) & H_2(BG; K_3(R)) & \cdots \\ 0 & H_1(BG; K_2(R)) & H_2(BG; K_2(R)) & \cdots \\ 0 & H_1(BG; K_1(R)) & H_2(BG; K_1(R)) & \cdots \\ 0 & H_1(BG; K_0(R)) & H_2(BG; K_0(R)) & \cdots \end{array} \\ \rightarrow p \end{array} \oplus \begin{array}{c} \uparrow q \\ \begin{array}{cccc} K_5(R) & 0 & 0 & \cdots \\ K_4(R) & 0 & 0 & \cdots \\ K_3(R) & 0 & 0 & \cdots \\ K_2(R) & 0 & 0 & \cdots \\ K_1(R) & 0 & 0 & \cdots \\ K_0(R) & 0 & 0 & \cdots \end{array} \\ \rightarrow p \end{array} \\ \text{AHSS}(BG_+)^2_{p,q} & & \text{AHSS}(BG)^2_{p,q} \quad \text{AHSS}(\mathbb{S}^0)^2_{p,q} \end{array} \quad (3.2.2)$$

(We omit the negative K-groups because they are annoying to include.) We remark that the groups $\text{AHSS}(BG)_{0,q}^2$ are trivial because the reduced homology $\tilde{H}_0(BG)$ is trivial, for BG being path-connected. Consequently,

Lemma 3.1

Let G be a group, and R be a ring. The differentials

$$d_{1,q}^2 : \text{AHSS}(BG_+)_{1,q}^2 \rightarrow \text{AHSS}(BG_+)_{-1,q+1}^2 \quad (3.2.3)$$

in the E^2 -page of the AHSS of BG_+ associated to the K-theory spectrum $\mathbb{K}_R^{\text{GW}}$ are trivial for all q .

Proof. The splitting of the AHSS in Equation (3.1.5) says the differential

$$d_{1,q}^2 : \text{AHSS}(BG_+)_{1,q}^2 \rightarrow \text{AHSS}(BG_+)_{-1,q+1}^2$$

splits into direct sum of the two differentials

$$\text{AHSS}(BG)_{1,q}^2 \rightarrow \text{AHSS}(BG)_{-1,q+1}^2, \quad (3.2.4)$$

$$\text{AHSS}(\mathbb{S}^0)_{1,q}^2 \rightarrow \text{AHSS}(\mathbb{S}^0)_{-1,q+1}^2. \quad (3.2.5)$$

From Diagram (3.2.2), we see that the first (resp. second) map is trivial because the codomain (resp. domain) is trivial, hence proving the claim. $\square$

Lemma 3.1 that says the group $\text{AHSS}(BG_+)_{1,i}^\infty$ is a quotient of $\text{AHSS}(BG_+)_{1,i}^2$, and therefore we have a diagram

$$\begin{array}{ccc} \text{AHSS}(BG)_{1,i}^2 \cong G_{ab} \otimes K_i(R) & \dashrightarrow & K_{i+1}(R[G]) \\ \downarrow & & \uparrow \alpha_{\text{Loday}} \\ \text{AHSS}(BG)_{1,i}^\infty & \hookrightarrow & \pi_{i+1}(BG \wedge \mathbb{K}_R^{\text{GW}}). \end{array} \quad (3.2.6)$$

We would like to study the filler. (Perhaps we should explain the absent of the disjoint base-point “+” in Diagram (3.2.6) is not a typo. As pointed out in Equation (3.2.1), $\pi_{i+1}(BG_+ \wedge \mathbb{K}_R^{\text{GW}})$ contains $K_{i+1}(R) \oplus \text{AHSS}(BG)_{1,i}^\infty$ as a subgroup, and we want to know what the Loday assembly is doing on the term $\text{AHSS}(BG)_{1,i}^\infty$.)

Let us go back to the E^1 -page of the AHSS. Unlike the later pages, the E^1 -page is not homotopy invariant—it depends on the choice of cell structure on BG . **For our purpose, we are using the skeletal filtration on the bar construction for BG .** Then we have:

$$\text{AHSS}(BG)_{1,i}^1 \cong \pi_{i+1} \left(\left(\bigvee_{g \in G} \mathbb{S}^1 \right) \wedge \mathbb{K}_R^{\text{GW}} \right). \quad (3.2.7)$$

The filler we want to study is then induced by the composition

$$\pi_{i+1} \left(\left(\bigvee_{g \in G} \mathbb{S}^1 \right) \wedge \mathbb{K}_R^{\text{GW}} \right) \xrightarrow{(i_G)_*} \pi_{i+1}(BG \wedge \mathbb{K}_R^{\text{GW}}) \xrightarrow{\alpha_{\text{Loday}}} K_{i+1}(R[G]), \quad (3.2.8)$$

for which the first arrow is induced by the inclusion

$$\mathbf{i}_G : \bigvee_{g \in G} \mathbb{S}^1 \hookrightarrow BG \quad (3.2.9)$$

of the 1-skeleton of BG into the BG . This follows from the construction of the AHSS. (See [Wic15] for example.) When $G = \langle t \rangle$ is the infinite cyclic group, we know the circle $\mathbb{S}^1$ is a model for $B\langle t \rangle$. Moreover,

Lemma 3.2

The inclusion

$$\mathbf{i}_t : \mathbb{S}^1 \rightarrow B\langle t \rangle \quad (3.2.10)$$

that sends $\mathbb{S}^1$ to the 1-cell of $B\langle t \rangle$ labelled by the generator t is a weak equivalence.

Proof. It is clear that the induced group homomorphism

$$(\mathbf{i}_t)_* : \pi_i(\mathbb{S}^1) \rightarrow \pi_i(B\langle t \rangle) \quad (3.2.11)$$

is an isomorphism for $i \neq 1$, because the source and target are both trivial. So we need to check the map

$$(\mathbf{i}_t)_* : \pi_1(\mathbb{S}^1) \rightarrow \pi_1(B\langle t \rangle) \quad (3.2.12)$$

is an isomorphism.

The map in Equation (3.2.12) is just an endomorphism of infinite cyclic group. In particular, it is surjective by the very definition of $\mathbf{i}_t$. (The generator is in the image.) So the First Isomorphism Theorem says the map in Equation (3.2.12) must have trivial kernel, and hence an isomorphism. $\square$

The Whitehead Theorem then asserts $\mathbf{i}_t$ is a homotopy equivalence. Thus when $G = \langle t \rangle$ is the infinite cyclic group, we have the following commutative diagram

$$\begin{array}{ccc} \bigvee_{g \in \langle t \rangle} \mathbb{S}^1 & \xrightarrow{\mathbf{i}_{\langle t \rangle}} & B\langle t \rangle \\ & \nwarrow \mathbf{i}_t & \uparrow \simeq \mathbf{i}_t \\ & & \mathbb{S}^1 \end{array} \quad (3.2.13)$$

The upshot is that we are able to describe the composition in Equation (3.2.8) when $G = \langle t \rangle$.

Proposition 3.3

When $G = \langle t \rangle$ is the infinite cyclic group, the composition in Equation (3.2.8) can be identified as

$$\begin{array}{ccccc}
\pi_{i+1} \left(\left(\bigvee_{g \in \langle t \rangle} \mathbb{S}^1 \right) \wedge \mathbb{K}_R^{\text{GW}} \right) & \xrightarrow{(\mathbf{i}_{\langle t \rangle})} & \pi_{i+1} (B \langle t \rangle \wedge \mathbb{K}_R^{\text{GW}}) & \xrightarrow{\alpha_{\text{Loday}}} & K_{i+1} (R[t^{\pm 1}]) \\
\uparrow (\mathbf{i}_t)_* & & \uparrow (\mathbf{i}_t)_* \cong & & \parallel \\
\pi_{i+1} (\mathbb{S}^1 \wedge \mathbb{K}_R^{\text{GW}}) & \xrightarrow{\text{id}} & \pi_{i+1} (\mathbb{S}^1 \wedge \mathbb{K}_R^{\text{GW}}) & \longrightarrow & K_{i+1} (R[t^{\pm 1}]) \\
\parallel & & \parallel & & \parallel \\
K_i(R) & \xrightarrow{\text{id}} & K_i(R) & \longrightarrow & K_{i+1} (R[t^{\pm 1}]),
\end{array} \tag{3.2.14}$$

where the bottom composition sends $[f] \in K_i(R)$ to

$$[f] \star \{\mu_{t^{-1}}\} \in K_{i+1} (R[t^{\pm 1}]),$$

for which $\mu_{t^{-1}} \in \text{Aut} (R[t^{\pm 1}])$ is multiplication by t^{-1} .

We remind readers the commutativity of the top-left square comes from Diagram (3.2.13).

Proof. The class $[f] \in K_i(R)$ is represented by the spheroid

$$f : \mathbb{S}^i \rightarrow K_R^+,$$

where

$$K_R^+ := K_0(R) \times BGL(R)^+.$$

Taking suspension yields

$$\mathbb{S}^1 \wedge \mathbb{S}^i \xrightarrow{\text{id} \wedge f} \mathbb{S}^1 \wedge K_R^+ \xrightarrow{j^+ \wedge \text{id}} K_{\mathbb{Z}[t^{\pm 1}]}^+ \wedge K_R^+ \xrightarrow{\gamma'_{\text{Loday}}} K_{R[t^{\pm 1}]}^+.$$

This composition represents the element $\alpha_{\text{Loday}}([f]) \in K_{i+1} (R[t^{\pm 1}])$, as well as the product

$$[f] \star' \{j^+\} \in K_{i+1} (R[t^{\pm 1}]).$$

Now, j^+ is induced by the group homomorphism

$$\begin{aligned}
j : \langle t \rangle &\rightarrow GL(\mathbb{Z}[t^{\pm 1}]) \\
t &\mapsto \begin{bmatrix} t & 0 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ 0 & 0 & 1 & 0 & \cdots \\ 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.
\end{aligned}$$

So $\{j^+\} = \{\mu_t\}$, and thus

$$\begin{aligned}
[f] \star' \{j^+\} &= [f] \star' \{\mu_t\} \\
&= [f] \star (-\{\mu_t\}) \\
&= [f] \star \{\mu_{t^{-1}}\}
\end{aligned}
\tag{Proposition 2.6}$$

as desired. □

For arbitrary group G , we recall the isomorphism

$$\begin{aligned}
G &\cong \text{Hom}(\langle t \rangle, G) \\
g &\mapsto (\varphi_g : t \mapsto g),
\end{aligned}
\tag{3.2.15}$$

allowing us to extend Proposition 3.3 to arbitrary G .

Proposition 3.4

For an arbitrary group G , under the canonical isomorphism

$$\pi_{i+1} \left(\left(\bigvee_{g \in G} \mathbb{S}^1 \right) \wedge \mathbb{K}_R^{\text{GW}} \right) \cong \bigoplus_{g \in G} K_i(R),
\tag{3.2.16}$$

the composition in (3.2.8) can be identified as

$$\begin{array}{ccccc}
\pi_{i+1} \left(\left(\bigvee_{g \in G} \mathbb{S}^1 \right) \wedge \mathbb{K}_R^{\text{GW}} \right) & \xrightarrow{(\mathbf{i}_G)_*} & \pi_{i+1} (BG \wedge \mathbb{K}_R^{\text{GW}}) & \xrightarrow{\alpha_{\text{Loday}}} & K_{i+1} (R[G]) \\
\parallel & & \parallel & & \parallel \\
\bigoplus_{g \in G} K_i(R) & \longrightarrow & \pi_{i+1} (BG \wedge \mathbb{K}_R^{\text{GW}}) & \longrightarrow & K_{i+1} (R[G]),
\end{array}
\tag{3.2.17}$$

where the bottom composition sends the element $[f] \in K_i(R)$ in the summand in $\bigoplus_{g \in G} K_i(R)$ labelled by $g \in G$ to the element

$$[f] \star \{\mu_{g^{-1}}\} \in K_{i+1} (R[G]),$$

for which $\mu_{g^{-1}} \in \text{Aut} (R[G])$ is multiplication by g^{-1} .

Proof. Fix $g \in G$, and consider the commutative diagram

$$\begin{array}{ccc}
\pi_{i+1}(\mathbb{S}^1 \wedge \mathbb{K}_R^{\text{GW}}) & \xrightarrow{(\varphi_g)_*} & \pi_{i+1}\left(\left(\bigvee_{g \in G} \mathbb{S}^1\right) \wedge \mathbb{K}_R^{\text{GW}}\right) \\
\downarrow (\mathbf{i}_t)_* \cong & & \downarrow (\mathbf{i}_G)_* \\
\pi_{i+1}(B\langle t \rangle \wedge \mathbb{K}_R^{\text{GW}}) & \xrightarrow{(\varphi_g)_*} & \pi_{i+1}(BG \wedge \mathbb{K}_R^{\text{GW}}) \\
\downarrow \alpha_{\text{Loday}} & & \downarrow \alpha_{\text{Loday}} \\
K_{i+1}(R[t^{\pm 1}]) & \xrightarrow{(\varphi_g)_*} & K_{i+1}(R[G]).
\end{array} \tag{3.2.18}$$

The top square commutes by chasing the definitions of the arrows. The bottom square commutes by the naturality of α_{Loday} via the group homomorphism

$$\varphi_g : \langle t \rangle \rightarrow G.$$

Now, Proposition 3.3 says the left vertical composition in Diagram (3.2.18) sends $[f] \in K_i(R)$ to

$$[f] \star \{\mu_{t^{-1}}\} \in K_{i+1}(R[t^{\pm 1}]).$$

So the commutativity of Diagram (3.2.18) says the right vertical composition sends $[f] \in K_i(R)$ in the summand $\bigoplus_{g \in G} K_i(R)$ indexed by $g \in G$ to the element

$$[f] \star \{\mu_{g^{-1}}\} \in K_{i+1}(R[G]).$$

□

Theorem 3.5 (An Explicit Formula for the Loday Assembly)

The filler in Diagram (3.2.6) is induced by the map

$$\begin{aligned}
G \times K_i(R) &\rightarrow K_{i+1}(R[G]) \\
(g, [f]) &\mapsto [f] \star \{\mu_{g^{-1}}\}.
\end{aligned}$$

Proof. Follows immediately from Proposition 3.4. □

Corollary 3.6 (The Loday Assembly on π_1 , [LR05, page 708])

For a regular ring R , the Loday assembly on π_1 is given by

$$K_1(i) \oplus \Phi_1 : K_1(R) \oplus [G_{ab} \otimes K_0(R)] \rightarrow K_1(R[G]), \tag{3.2.19}$$

for which $i : R \rightarrow R[G]$ is the inclusion, and Φ_1 is induced by the map

$$\begin{aligned} G \times K_0(R) &\rightarrow K_1(R[G]) \\ (g, [P]) &\mapsto \{\mathfrak{h}_g\}, \end{aligned} \tag{3.2.20}$$

for which $\mathfrak{h}_g : P \otimes R[G] \rightarrow R \otimes R[G]$ is the automorphism given by

$$\mathfrak{h}_g(x \otimes u) := x \otimes ug^{-1}.$$

Proof. If $g \in G$, and $[P] \in K_0(R)$, then Theorem 3.5 says the Loday assembly sends $(g, [P])$ to

$$\begin{aligned} [P] \star' \{\mu_g\} &= [P] \# \{\mu_{g^{-1}}\} \\ &= \{\mathfrak{h}_g\}. \end{aligned} \tag{Theorem 2.8}$$

□

3.3. Applications to the Second K-groups of Integral Group Rings. The following SES

$$0 \rightarrow K_2(\mathbb{Z}) \oplus [G_{ab} \otimes K_1(\mathbb{Z})] \rightarrow \pi_2(BG_+ \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}}) \rightarrow H_2(BG; K_0(\mathbb{Z})) \rightarrow 0 \tag{3.3.1}$$

is the extension problem of the AHSS for computing the second homotopy group $\pi_2(BG_+ \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}})$. See [Leh18, Theorem 12.2 on page 29]. Together with Theorem 3.5, we give some injectivity results for the Loday assembly on the second K-group. We begin by the following corollary:

Corollary 3.7 (The Loday Assembly on π_2 of an Integral Group Ring)

Let G be a group. The Loday assembly map

$$\alpha_{\text{Loday}} : BG_+ \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}} \rightarrow \mathbb{K}_{\mathbb{Z}[G]}^{\text{GW}} \tag{3.3.2}$$

for the integral group ring $\mathbb{Z}[G]$ on π_2 , when restricted onto the subgroup

$$K_2(\mathbb{Z}) \oplus [G_{ab} \oplus K_1(\mathbb{Z})],$$

is given by the formula:

$$K_2(i) \oplus \Phi_2 : K_2(\mathbb{Z}) \oplus [G_{ab} \oplus K_1(\mathbb{Z})] \rightarrow K_2(\mathbb{Z}[G]), \tag{3.3.3}$$

for which $i : \mathbb{Z} \rightarrow \mathbb{Z}[G]$ is the inclusion, and Φ_2 is induced by the map

$$\begin{aligned} G \times K_1(\mathbb{Z}) &\rightarrow K_2(\mathbb{Z}[G]) \\ (g, \pm 1) &\mapsto \{\pm 1, g\}_{\text{St}}. \end{aligned} \tag{3.3.4}$$

In particular, the Steinberg symbol $\{1, g\}_{\text{St}}$ is always the identity element in $K_2(\mathbb{Z}[G])$ for all $g \in G$.

Proof. Only the description of Φ_2 needs to be justified. Theorem 3.5 says Φ_2 is induced by the map

$$\begin{aligned} G \times K_1(\mathbb{Z}) &\rightarrow K_2(\mathbb{Z}[G]) \\ (g, \pm 1) &\mapsto \{\pm 1\} \star \{\mu_{g^{-1}}\}, \end{aligned} \tag{3.3.5}$$

and we have

$$\begin{aligned}
\{\pm 1\} \star \{\mu_{g^{-1}}\} &= -\{\pm 1, \mu_{g^{-1}}\}_{\text{St}} && \text{(Proposition 2.4)} \\
&= -\{\pm 1, g^{-1}\}_{\text{St}} && \text{(By definition of } \mu_{g^{-1}}) \\
&= \{\pm 1, (g^{-1})^{-1}\}_{\text{St}} && \text{(Bilinearity of the Steinberg symbol)} \\
&= \{\pm 1, g\}_{\text{St}}
\end{aligned}$$

as desired. $\square$

We now see that the Steinberg symbol $\{-1, g\}_{\text{St}}$ is closely related to the non-injectivity problem of α_{Loday} .

Example 3.8 (Injectivity on $K_2(\mathbb{Z}[C_2])$). Denote by C_2 the cyclic group of order two, with generator g . Dunwoody showed, in [Dun75, Theorem on page 482], that the second K-group of $\mathbb{Z}[C_2]$ is generated by two Steinberg symbols:

$$\begin{aligned}
K_2(\mathbb{Z}[C_2]) &= \langle \{-1, -1\}_{\text{St}}, \{-1, g\}_{\text{St}} \rangle \\
&\cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}.
\end{aligned} \tag{3.3.6}$$

Theorem 3.5 then says the Loday assembly

$$\alpha_{\text{Loday}} : (BC_2)_+ \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}} \rightarrow \mathbb{K}_{\mathbb{Z}[C_2]}^{\text{GW}} \tag{3.3.7}$$

for the cyclic group of order 2 is injective on π_2 . In fact, by the vanishing of the second Whitehead group $\text{Wh}_2(C_2)$ [Dun75, Theorem on page 482], we know it is, in fact, bijective on π_2 .

In fact, we have an expression for the Steinberg symbol $\{-1, g\}_{\text{St}}$ in terms of generators of the Steinberg group $\text{St}(\mathbb{Z}[G])$.

Theorem 3.9

Let G be a group. The Steinberg symbol $\{-1, g\}_{\text{St}} \in K_2(\mathbb{Z}[G])$ for $g \in G$ is given by

$$\{-1, g\}_{\text{St}} = w_{12}(-1) w_{12}(-1) w_{12}(g) w_{12}(g), \tag{3.3.8}$$

where

$$w_{ij}(u) := x_{ij}(u) x_{ji}(-u^{-1}) x_{ij}(u), \tag{3.3.9}$$

and the $x_{ij}(u)$ s are the generators of the Steinberg group $\text{St}(\mathbb{Z}[G])$.

Proof. The proof is about playing with the presentation of the Steinberg group.

We recall from [Ros95, Lemma 4.2.15 on page 195] the identities

$$w_{ij}(u)^{-1} = w_{ij}(-u), \tag{3.3.10}$$

$$w_{ij}(u) = w_{ji}(-u^{-1}), \tag{3.3.11}$$

$$w_{k\ell}(u) w_{ij}(v) w_{k\ell}(u)^{-1} = \begin{cases} w_{ij}(v) & \text{if } i, j, k, \ell \text{ are all distinct,} \\ w_{\ell j}(-u^{-1}v) & \text{if } k = i, \text{ and } i, j, \ell \text{ are all distinct,} \\ w_{i\ell}(-vu) & \text{if } k = j, \text{ and } i, j, k \text{ are all distinct,} \\ w_{ji}(-u^{-1}vu^{-1}) & \text{if } k = i, \text{ and } j = \ell. \end{cases} \quad (3.3.12)$$

In particular, Equation (3.3.12) gives

$$w_{13}(u) w_{12}(v) = w_{23}(v^{-1}u) w_{13}(u), \quad (3.3.13)$$

$$w_{13}(u) w_{23}(v) = w_{12}(-uv^{-1}) w_{13}(u). \quad (3.3.14)$$

Put

$$h_{ij}(u) := w_{ij}(u) w_{ij}(-1), \quad (3.3.15)$$

then a long and boring computation shows

$$\begin{aligned} \{-1, g\}_{\text{St}} &:= [h_{12}(-1), h_{13}(g)] \\ &= h_{12}(-1) h_{13}(g) h_{12}(-1)^{-1} h_{13}(g)^{-1} \\ &= w_{12}(-1) w_{12}(-1) w_{13}(g) w_{13}(-1) w_{12}(-1)^{-1} w_{12}(-1)^{-1} w_{13}(-1)^{-1} w_{13}(g)^{-1} \\ &= w_{12}(-1) w_{12}(-1) w_{13}(g) w_{13}(-1) w_{12}(1) w_{12}(1) w_{13}(1) w_{13}(-g) \\ &\quad (\text{Equation (3.3.10)}) \\ &= w_{12}(-1) w_{12}(-1) w_{13}(g) \underbrace{w_{13}(-1) w_{12}(1)}_{w_{23}(-1) w_{13}(-1)} w_{12}(1) w_{13}(1) w_{13}(-g) \\ &\quad (\text{Equation (3.3.13)}) \\ &= w_{12}(-1) w_{12}(-1) w_{13}(g) w_{23}(-1) \underbrace{w_{13}(-1) w_{12}(1)}_{w_{23}(-1) w_{13}(-1)} w_{13}(1) w_{13}(-g) \\ &\quad (\text{Equation (3.3.13)}) \\ &= w_{12}(-1) w_{12}(-1) w_{13}(g) w_{23}(-1) w_{23}(-1) \underbrace{w_{13}(-1) w_{13}(1)}_1 w_{13}(-g) \\ &\quad (\text{Equation (3.3.10)}) \\ &= w_{12}(-1) w_{12}(-1) \underbrace{w_{13}(g) w_{23}(-1)}_{w_{12}(g) w_{13}(g)} w_{23}(-1) w_{13}(-g) \\ &\quad (\text{Equation (3.3.14)}) \end{aligned}$$

$$\begin{aligned}
&= w_{12}(-1) w_{12}(-1) w_{12}(g) \underbrace{w_{13}(g) w_{23}(-1) w_{13}(-g)}_{w_{31}(-g^{-1}) w_{23}(-1) w_{31}(g^{-1})} \\
&\quad (\text{Equation (3.3.11)}) \\
&= w_{12}(-1) w_{12}(-1) w_{12}(g) \underbrace{w_{31}(-g^{-1}) w_{23}(-1) w_{31}(g^{-1})}_{w_{21}(-g^{-1})} \\
&\quad (\text{Equation (3.3.12)}) \\
&= w_{12}(-1) w_{12}(-1) w_{12}(g) \underbrace{w_{21}(-g^{-1})}_{w_{12}(g)} \\
&\quad (\text{Equation (3.3.11)}) \\
&= w_{12}(-1) w_{12}(-1) w_{12}(g) w_{12}(g)
\end{aligned}$$

as desired. $\square$

Theorem 3.9 motivates the following definition:

Definition 3.10 (The group $W_{12}(G)$)

Let G be a group. The subgroup $W_{12}(G)$ of $K_2(\mathbb{Z}[G])$ is generated by the Steinberg symbols:

$$W_{12}(G) := \langle \{-1, g\}_{\text{St}} \mid g \in G \rangle. \quad (3.3.16)$$

Notice the following:

- (1) The image of the map

$$\Phi_2 : G \times K_1(\mathbb{Z}) \rightarrow K_2(\mathbb{Z}[G])$$

in Equation (3.3.3) is precisely $W_{12}(G)$.

- (2) Readers who are familiar with the classical definition

$$\text{Wh}_2(G) := \frac{K_2(\mathbb{Z}[G])}{K_2(\mathbb{Z}[G]) \cap W(G)} \quad (3.3.17)$$

of the second Whitehead group proposed by Hatcher-Wagoner in [HW73, page 10] will see our group $W_{12}(G)$ follows their spirit. The group $W(G)$ is generated by the elements $w_{ij}(\pm g)$ for all $g \in G$, and for all i, j . Together with Theorem 3.9, we see that $W_{12}(G)$ is a subgroup of $W(G)$.

- (3) From the functoriality of the Steinberg group, we get a functor

$$W_{12}(-) : \text{Groups} \rightarrow \text{Abelian}. \quad (3.3.18)$$

Moreover, for every group G , there is a natural group homomorphism

$$\begin{aligned}\Psi_G : G &\rightarrow W_{12}(G) \\ g &\mapsto \{-1, g\}_{\text{St}},\end{aligned}\tag{3.3.19}$$

or equivalently the restriction

$$\Psi_G = \Phi_2|_{\{-1\} \times G}\tag{3.3.20}$$

for Φ_2 as in Equation (3.3.3). Non-injectivity of Ψ_G will imply the non-injectivity of the Loday assembly on π_2 .

Example 3.11 (Injectivity on $K_2(\mathbb{Z}[C_2^{\oplus n}])$). This example is due to Daniel Ramras.

Denote by C_2 the cyclic group of order 2. Fix $n \in \mathbb{N}$, and consider the n -fold direct sum. If $x \in C_2^{\oplus n}$ is non-zero, then there exists $1 \leq i \leq n$ such that the image $\text{proj}_i(x)$ of x under the i -th projection map

$$\text{proj}_i : C_2^{\oplus n} \rightarrow C_2$$

is non-zero. The functoriality of $W_{12}(-)$ then gives a commutative diagram

$$\begin{array}{ccc} C_2^{\oplus n} & \xrightarrow{\Psi_{C_2^{\oplus n}}} & W_{12}(C_2^{\oplus n}) \\ \text{proj}_i \downarrow & & \downarrow W_{12}(\text{proj}_i) \\ C_2 & \xrightarrow{\Psi_{C_2}} & W_{12}(C_2). \end{array}\tag{3.3.21}$$

We saw in Example 3.8 that Ψ_{C_2} is injective, so the commutativity of Diagram (3.3.21) says $\Psi_{C_2^{\oplus n}}(x) \in W_{12}(C_2^{\oplus n})$ is non-zero.

Let us recap what we did here. Fix a non-zero element $x \in C_2^{\oplus n}$. There exists $1 \leq i \leq n$ such that $\text{proj}_i(x) \in C_2$ is non-zero. Because Ψ_{C_2} is injective, we must have

$$\left(W_{12}(\text{proj}_i) \circ \Psi_{C_2^{\oplus n}} \right) (x) \neq 0,$$

and consequently $\Psi_{C_2^{\oplus n}}(x) \neq 0$, proving the group homomorphism

$$\Psi_{C_2^{\oplus n}} : C_2^{\oplus n} \rightarrow W_{12}(C_2^{\oplus n})$$

is injective for any $n \in \mathbb{N}$.

As a result, the Loday assembly map

$$\alpha_{\text{Loday}} : B(C_2^{\oplus n})_+ \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}} \rightarrow \mathbb{K}_{\mathbb{Z}[C_2^{\oplus n}]}^{\text{GW}}$$

for the group $C_2^{\oplus n}$ is injective on π_2 when restricted onto the subgroup $K_2(\mathbb{Z}) \oplus [C_2^{\oplus n} \otimes K_1(\mathbb{Z})]$.

Example 3.12 (Limitation). Our formula does not work well on studying the K-theory of the integral group ring $\mathbb{Z}[C_p]$ of cyclic group C_p of prime order $p > 2$. Since $C_p \otimes C_2 = 0$,

the map Φ_2 in Equation (3.3.4) is the zero map. Because $H_2(BC_p; \mathbb{Z}) = 0$ for all p , the SES in Equation (3.3.1) gives

$$\pi_2(BC_{p+} \wedge \mathbb{K}_{\mathbb{Z}}^{\text{GW}}) \cong K_2(\mathbb{Z}), \quad (3.3.22)$$

and therefore we learn nothing about structures of $K_2(\mathbb{Z}[C_p])$ from the assembly map approach, other than it contains $K_2(\mathbb{Z})$ as a subgroup.

INDEX

K_R^+ , [5](#)

$W_{12}(G)$, [21](#)

$\# : K_i(R) \otimes K_j(R) \rightarrow K_{i+j}(R \otimes R')$,
classical pairings in algebraic

K-theory, [7](#)

$i_G : \bigvee_{g \in G} \mathbb{S}^1 \hookrightarrow BG$, the inclusion from the
1-skeleton of BG into the BG , [14](#)

$i_t : \mathbb{S}^1 \rightarrow B \langle t \rangle$, the inclusion that sends $\mathbb{S}^1$
to the 1-cell of $B \langle t \rangle$ labelled by the
generator t , [14](#)

α_{Loday} , Loday assembly, [10](#)

$\Psi_G : G \rightarrow W_{12}(G)$, [22](#)

γ_{Loday} , the Loday pairing, [4](#)

γ'_{Loday} , modified Loday pairing, [6](#)

$\gamma_{m,n}$, [3](#)

$\text{AHSS}(X)_{p,q}^r$, the (p, q) -term on the
 E^r -page of the Atiyah-Hirzebruch
spectral sequence of the space X ,
represented by some spectrum, [10](#)

$\star$, Loday product, [4](#)

$\varphi_g : \langle t \rangle \rightarrow G$, [16](#)

Loday assembly, [10](#)

REFERENCES

- [BHM93] M. Bökstedt, W. C. Hsiang, and I. Madsen. “The cyclotomic trace and algebraic K -theory of spaces”. In: *Invent. Math.* 111.3 (1993), pages 465–539. ISSN: 0020-9910. DOI: [10.1007/BF01231296](https://doi.org/10.1007/BF01231296). URL: <https://doi.org/10.1007/BF01231296>.
- [Coh73] Marshall M. Cohen. *A Course in Simple-Homotopy Theory*. Graduate Texts in Mathematics 10. Springer, 1973. ISBN: 978-0387900551.
- [DL98] James F. Davis and Wolfgang Lück. “Spaces over a category and assembly maps in isomorphism conjectures in K - and L -theory”. In: *K-Theory* 15.3 (1998), pages 201–252. ISSN: 0920-3036. DOI: [10.1023/A:1007784106877](https://doi.org/10.1023/A:1007784106877). URL: <https://doi.org/10.1023/A:1007784106877>.
- [Dun75] M. J. Dunwoody. “ $K_2(\mathbf{Z}\Pi)$ for Π a group of order two or three”. In: *J. London Math. Soc. (2)* 11.4 (1975), pages 481–490. ISSN: 0024-6107. DOI: [10.1112/jlms/s2-11.4.481](https://doi.org/10.1112/jlms/s2-11.4.481). URL: <https://doi.org/10.1112/jlms/s2-11.4.481>.
- [ES52] Samuel Eilenberg and Norman E. Steenrod. *Foundations of Algebraic Topology*. 1st edition. Princeton University Press, 1952. ISBN: 978-0691079653.
- [Ger72] Stephen M. Gersten. “On the spectrum of algebraic K -theory”. In: *Bull. Amer. Math. Soc.* 78.2 (Mar. 1972), pages 216–219. URL: <https://projecteuclid.org:443/euclid.bams/1183533474>.
- [Hat01] Allen Hatcher. *Algebraic Topology*. Cambridge University Press, 2001. ISBN: 0521535379.
- [Hsi84] Wu Chung Hsiang. “Geometric applications of algebraic K -theory”. In: *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983)*. PWN, Warsaw, 1984, pages 99–118.
- [HW73] Allen Hatcher and John Wagoner. *Pseudo-isotopies of compact manifolds*. With English and French prefaces, Astérisque, No. 6. Société Mathématique de France, Paris, 1973, pages i+275.
- [Leh18] Georg Lehner. “The Atiyah-Hirzebruch Spectral Sequence and K -theory of Group Rings”. Master’s Thesis. Humboldt-Universität zu Berlin, 2018.
- [Lod76] Jean-Louis Loday. “ K -théorie algébrique et représentations de groupes”. French. In: *Annales scientifiques de l’École Normale Supérieure* 9.3 (1976), pages 309–377.
- [LR05] Wolfgang Lück and Holger Reich. “The Baum–Connes and the Farrell–Jones Conjectures in K - and L -Theory”. In: *Handbook of K-Theory*. Edited by Eric M. Friedlander and Daniel R. Grayson. Berlin, Heidelberg: Springer Berlin Heidelberg, 2005, pages 703–842. ISBN: 978-3-540-27855-9. DOI: [10.1007/978-3-540-27855-9_15](https://doi.org/10.1007/978-3-540-27855-9_15). URL: https://doi.org/10.1007/978-3-540-27855-9_15.
- [May80] J. Peter May. “Pairings of categories and spectra”. In: *Journal of Pure and Applied Algebra* 19 (1980), pages 299–346. ISSN: 0022-4049.
- [Mil72] John W. Milnor. *Introduction to Algebraic K-Theory*. Annals of Mathematics Studies 71. Princeton University Press, 1972. ISBN: 978-0691081014.
- [Ros95] Jonathan Rosenberg. *Algebraic K-Theory and Its Applications*. 2nd edition. Graduate Texts in Mathematics 147. Springer, 1995. ISBN: 0387942483.
- [Shi88] Kazuhisa Shimakawa. “Uniqueness of products in higher algebraic K -theory”. In: *Publ. Res. Inst. Math. Sci.* 24.1 (1988), pages 99–120. ISSN: 0034-5318. DOI: [10.2977/prims/1195175327](https://doi.org/10.2977/prims/1195175327). URL: <https://doi.org/10.2977/prims/1195175327>.
- [UW17] Mark Ullmann and Xiaolei Wu. “Note on the injectivity of the Loday assembly map”. In: *J. Algebra* 489 (2017), pages 460–462. ISSN: 0021-8693. DOI: [10.1016/](https://doi.org/10.1016/)

- j.jalgebra.2017.06.037. URL: <https://doi.org/10.1016/j.jalgebra.2017.06.037>.
- [Wag72] John B. Wagoner. “Delooping classifying spaces in algebraic K-theory”. In: *Topology* 11.4 (1972), pages 349–370. ISSN: 0040-9383. DOI: [https://doi.org/10.1016/0040-9383\(72\)90031-6](https://doi.org/10.1016/0040-9383(72)90031-6). URL: <http://www.sciencedirect.com/science/article/pii/0040938372900316>.
- [Wal78a] Friedhelm Waldhausen. “Algebraic K-Theory of Generalized Free Products, Part 1”. In: *Annals of Mathematics* 108.2 (1978), pages 135–204. ISSN: 0003486X. URL: <http://www.jstor.org/stable/1971165>.
- [Wal78b] Friedhelm Waldhausen. “Algebraic K-Theory of Generalized Free Products, Part 2”. In: *Annals of Mathematics* 108.2 (1978), pages 205–256. ISSN: 0003486X. URL: <http://www.jstor.org/stable/1971166>.
- [Wei13] Charles A. Weibel. *The K-Book: An Introduction to Algebraic K-Theory*. Graduate Studies in Mathematics 145. American Mathematical Society, 2013. ISBN: 0821891324.
- [Wei81] Charles A. Weibel. “A survey of products in algebraic K-theory”. In: *Algebraic K-theory, Evanston 1980 (Proc. Conf., Northwestern Univ., Evanston, Ill., 1980)*. Volume 854. Lecture Notes in Math. Springer, Berlin, 1981, pages 494–517.
- [Whi50] J. H. C. Whitehead. “Simple homotopy types”. In: *Amer. J. Math.* 72 (1950), pages 1–57. ISSN: 0002-9327. DOI: [10.2307/2372133](https://doi.org/10.2307/2372133). URL: <https://doi.org/10.2307/2372133>.
- [Wic15] Kirsten Wickelgren. *Atiyah-Hirzebruch spectral sequence*. 2015. URL: http://people.math.gatech.edu/~kwickelgren3/8803_Stable/L13_Atiyah_Hirzebruch_Spectral_sequence.pdf.
- [WW95] Michael Weiss and Bruce Williams. “Assembly”. In: *Novikov Conjectures, Index Theorems, and Rigidity*. Edited by Steven C. Ferry, Andrew Ranicki, and Jonathan M. Rosenberg. Volume 2. London Mathematical Society Lecture Note Series. Cambridge University Press, 1995, 332352. DOI: [10.1017/CB09780511629365.014](https://doi.org/10.1017/CB09780511629365.014).