ការផ្សាយពាណិជ្ជកម្ម
Your health questions, answered
Our experts share insights and advice in Health Essentials News
Sign Up

# ជំងឺស្ត្រេសស្រួចស្រាវ

ជំងឺស្ត្រេសស្រួចស្រាវ (ASD) អាចកើតឡើងក្នុងរយៈពេលមួយខែនៃការជួបប្រទះនូវព្រឹត្តិការណ៍ដ៏តក់ស្លុត ដូចជាគ្រោះធម្មជាតិ ឬការវាយប្រហារជាដើម។ វាទាក់ទងយ៉ាងជិតស្និទ្ធទៅនឹងជំងឺស្ត្រេសក្រោយការប៉ះទង្គិចផ្លូវចិត្ត (PTSD)។ ការព្យាបាលសំខាន់សម្រាប់ជំងឺស្ត្រេសស្រួចស្រាវគឺការព្យាបាលចិត្តសាស្ត្រ (ការព្យាបាលដោយការនិយាយ) ។

មាតិកា ↓ ទិដ្ឋភាពទូទៅ រោគសញ្ញានិងមូលហេតុ ការធ្វើរោគវិនិច្ឆ័យ និងការធ្វើតេស្ត ការគ្រប់គ្រង និងការព្យាបាល ការបង្ការ ទស្សនវិស័យ / ការព្យាករណ៍ រស់នៅជាមួយ

ការផ្សាយពាណិជ្ជកម្ម
healthessentials
Your health matters
Got a question? Our experts offer answers in Health Essentials News
Sign Up
Cleveland Clinic គឺជាមជ្ឈមណ្ឌលវេជ្ជសាស្ត្រសិក្សាមិនរកប្រាក់ចំណេញ។ ការផ្សាយពាណិជ្ជកម្មនៅលើគេហទំព័ររបស់យើងជួយគាំទ្រដល់បេសកកម្មរបស់យើង។ យើងមិនគាំទ្រផលិតផល ឬសេវាកម្មដែលមិនមែនជា Cleveland Clinic ទេ។ គោលនយោបាយ

ការថែទាំនៅ Cleveland Clinic

ការថែទាំសុខភាពផ្លូវចិត្ត

ធ្វើការណាត់ជួប

## ទិដ្ឋភាពទូទៅ

### តើជំងឺស្ត្រេសស្រួចស្រាវគឺជាអ្វី?

ជំងឺស្ត្រេសស្រួចស្រាវ (ASD) គឺជាស្ថានភាពសុខភាពផ្លូវចិត្តរយៈពេលខ្លី ដែលអាចកើតមានឡើងក្នុងខែដំបូង បន្ទាប់ពីជួបប្រទះនូវព្រឹត្តិការណ៍ប៉ះទង្គិចផ្លូវចិត្ត។ វាពាក់ព័ន្ធនឹងការឆ្លើយតប ស្ត្រេស រួមទាំង៖

- ការថប់បារម្ភ ។
- ការភ័យខ្លាចខ្លាំងឬភាពអស់សង្ឃឹម។
- ជួបប្រទះនឹងការត្រលប់មកវិញ ឬសុបិន្តអាក្រក់។
- មានអារម្មណ៍ស្ពឹក ឬដាច់ចេញពីខ្លួន។
- ជៀសវាងស្ថានភាព ទីកន្លែង ឬការរំលឹកផ្សេងទៀតដែលទាក់ទងនឹងព្រឹត្តិការណ៍ដ៏តក់ស្លុតនេះ។

ឧទាហរណ៍នៃព្រឹត្តិការណ៍របួសរួមមាន:

- គ្រោះធម្មជាតិ ដូចជាខ្យល់ព្យុះ ខ្យល់ព្យុះ ភ្លើង ឬទឹកជំនន់។
- ការរំលោភផ្លូវភេទ។
- ការរំលោភលើរាងកាយ។
- ការរំលោភបំពានពាក្យសំដី។
- ការធ្វើជាសាក្សីអំពីការខូចខាតរាងកាយឬការស្លាប់។
- គ្រោះថ្នាក់ធ្ងន់ធ្ងរ ដូចជាគ្រោះថ្នាក់ចរាចរណ៍។
- ជួបប្រទះការរងរបួសធ្ងន់ធ្ងរ ឬជំងឺភ្លាមៗ។
- សង្គ្រាម។

ជំងឺស្ត្រេសស្រួចស្រាវអាចប៉ះពាល់ដល់មនុស្សម្នាក់នៅគ្រប់ដំណាក់កាលនៃជីវិត - កុមារភាព វ័យជំទង់ និងពេញវ័យ។

### តើអ្វីជាភាពខុសគ្នារវាងជំងឺស្ត្រេសស្រួចស្រាវ និងជំងឺ PTSD?

ភាពខុសគ្នាសំខាន់រវាងជំងឺស្ត្រេសស្រួចស្រាវ និង ជំងឺស្ត្រេសក្រោយការប៉ះទង្គិចផ្លូវចិត្ត (PTSD) គឺជារយៈពេលនៃរោគសញ្ញា។

ជំងឺស្ត្រេសស្រួចស្រាវពាក់ព័ន្ធនឹងប្រតិកម្មស្ត្រេសដែលកើតឡើងរវាងបីថ្ងៃទៅបួនសប្តាហ៍បន្ទាប់ពីព្រឹត្តិការណ៍ប៉ះទង្គិចផ្លូវចិត្តមួយ។ ប្រតិកម្មស្ត្រេសដែលមានរយៈពេលយូរជាង 4 សប្តាហ៍អាចបំពេញតាមលក្ខណៈវិនិច្ឆ័យសម្រាប់ PTSD ។

សមាគមចិត្តសាស្រ្តអាមេរិកបានចាត់ថ្នាក់ជំងឺស្ត្រេសស្រួចស្រាវជាលើកដំបូងជាការធ្វើរោគវិនិច្ឆ័យសុខភាពផ្លូវចិត្តក្នុងឆ្នាំ 1994 នៅក្នុងការបោះពុម្ពលើកទី 4 នៃ *សៀវភៅណែនាំរោគវិនិច្ឆ័យ និងស្ថិតិនៃជំងឺផ្លូវចិត្ត (DSM-IV)* ។

### តើអ្វីជាភាពខុសគ្នារវាងជំងឺកែតម្រូវ និងជំងឺស្ត្រេសស្រួចស្រាវ?

វិបត្តិនៃការកែតម្រូវ គឺជាស្ថានភាពសុខភាពផ្លូវចិត្តដែលកើតឡើងក្នុងការឆ្លើយតបទៅនឹងព្រឹត្តិការណ៍ជីវិតដ៏តានតឹង។ រោគសញ្ញាផ្លូវចិត្ត និងអាកប្បកិរិយា ជាទូទៅត្រូវបានគេចាត់ទុកថា ធ្ងន់ធ្ងរជាងអ្វីដែលរំពឹងទុកសម្រាប់ប្រភេទនៃព្រឹត្តិការណ៍ដែលបានកើតឡើង។

The difference between acute stress disorder and adjustment disorder is that the stressful event in adjustment disorder (such as losing your job or the ending of a relationship) is typically less traumatic than an event that causes acute stress disorder (such as sexual assault or surviving a mass shooting).

### How common is acute stress disorder?

It's difficult for researchers to assess how common acute stress disorder is. This is partly because people may not seek professional help until their symptoms meet the criteria for PTSD.

According to various studies, the prevalence of acute stress disorder following a traumatic event may range from 6% to 33%. The rates also differ for different types of trauma. For example, survivors of accidents or natural disasters show lower rates of ASD. Survivors of violence, such as assaults and mass shootings, show higher rates of ASD.

## Symptoms and Causes

### What are the symptoms of acute stress disorder?

Symptoms of ASD are psychological and behavioral stress responses. They may include:

- Recurring, uncontrollable and distressing memories of the event.
- Recurring nightmares.
- Flashbacks of the traumatic event.
- Intense psychologic or physical distress when you're reminded of the event.
- Persistent difficulty feeling positive emotions, such as happiness, contentment or loving feelings.
- An altered sense of reality, like feeling you're in a daze or as if time is passing in slow motion.
- Memory loss regarding important aspects of the traumatic event.
- Efforts to avoid distressing memories, thoughts or feelings associated with the event.
- Efforts to avoid external reminders associated with the event (people, places or things).
- Disturbed sleep.
- Irritability or anger outbursts.
- Excessive attention to the possibility of danger (hypervigilance).
- Difficulty concentrating.
- An exaggerated response to loud noises, sudden movements or other stimuli (startle reflex).

ADVERTISEMENT

### What causes acute stress disorder?

It's unclear why people respond differently to traumatic events.

One theory involves the concept of "fear conditioning." This happens when your body exhibits a fear response to certain stimuli associated with a traumatic event. For example, if you were in a car accident and you had fast food in your vehicle and it was nighttime, future encounters with the smell of fast food at night may trigger your body to have the same fear response that you did during the traumatic event even though there's no threat to your safety.

Some people may adapt to fear conditioning via extinction learning, which involves a gradual reduction in response to the traumatic triggers. If this doesn't work, you could develop acute stress disorder and potentially PTSD.

#### What are the risk factors for acute stress disorder?

Acute stress disorder risk factors may include:

- History of prior mental health condition(s) (including prior trauma).
- Catastrophic worry.
- Avoidant coping style.
- Minimal support system.

### What are the complications of this condition?

The main complication of acute stress disorder is the potential development of post-traumatic stress disorder (PTSD). PTSD may develop if acute stress disorder isn't treated.

## Diagnosis and Tests

### How is acute stress disorder diagnosed?

There's no test to diagnose acute stress disorder. Instead, a healthcare provider makes the diagnosis after conducting a thorough psychosocial assessment. They'll ask about your current symptoms and about your medical and mental health history.

Providers use criteria explained in the *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition* (*DSM-5*) to diagnose acute stress disorder.

ADVERTISEMENT

## Management and Treatment

### What is the treatment for acute stress disorder?

Psychotherapy (talk therapy) is the main treatment for acute stress disorder. A form of cognitive behavioral therapy (CBT) called trauma-focused CBT is especially effective.

This therapy takes place with a trained, licensed mental health professional, such as a psychologist or psychiatrist. It provides support, education and guidance to you and, if necessary, your family/support system.

Trauma-focused CBT involves:

- Learning about how your body responds to trauma and stress.
- Symptom management skills.
- Identifying and reframing problematic thinking patterns.
- Exposure therapy.

Exposure therapy involves safely exposing you to your sources of fear and avoidance. People with acute stress disorder may avoid things or situations that they associate with the traumatic event. Because of this, they aren't able to learn that they can manage their fears when presented with these stimuli. Therapists use exposure therapy for people who have ASD to slowly encourage them to enter situations that cause them anxiety and to try to stay in that situation so that they can learn to cope.

#### Other treatments for acute stress disorder

While there's minimal evidence supporting the use of medications to treat acute stress disorder, some medications may alleviate the symptoms of PTSD, including:

- Selective serotonin reuptake inhibitors (SSRIs).
- Serotonin and norepinephrine reuptake inhibitors (SNRIs).

## Prevention

### Can acute stress disorder be prevented?

You can't necessarily prevent a traumatic event. But research shows that the intensity of acute stress disorder symptoms is manageable. The following can help:

- **Personal safety**: Experiencing a sense of safety is crucial following a traumatic event. The support of loved ones and professionals trained to provide assistance and guidance following a traumatic event can be greatly beneficial.
- **Physical health**: It's important to maintain a healthy daily routine following a traumatic event. Prioritizing a healthy diet, sleep schedule and involvement in physical activity and mindfulness exercises (like meditation) are important. You should avoid the use of alcohol or other substances as temporary coping tools.
- **Emotional support**: It's vital to seek emotional support from family and friends following a traumatic event. If this isn't possible, healthcare providers can serve as support.
- **Follow up with your healthcare team**: It's important to seek treatment with your healthcare provider and/or mental health provider following a traumatic event.

ADVERTISEMENT

## Outlook / Prognosis

### What is the prognosis of acute stress disorder?

The prognosis (outlook) for acute stress disorder varies. If symptoms are untreated, the potential for further difficulties increase, including:

- Alcohol use disorder and/or substance use disorder.
- Other mental health conditions, like PTSD, anxiety, depression or panic disorder.
- Thoughts about suicide/harming yourself or others.
- Problems at work and in your relationships.

**If you or a loved one are experiencing suicidal thoughts, dial 988 on your phone to reach the Suicide and Crisis Lifeline.**

## Living With

### How do I take care of myself if I have acute stress disorder?

In addition to seeking professional treatment for acute stress disorder, you may benefit from positive activities, including:

- Engaging in exercise to help reduce stress.
- Setting realistic goals and seeing them through to completion.
- Spending time with people you trust and educating them about your experience and ways in which they can be supportive.
- Identifying and seeking out comforting situations and places.
- Attending a trauma support group.
- Prioritizing patience and self-care. It's important to remember that symptoms will improve gradually, not immediately.

ADVERTISEMENT

### When should I see my healthcare provider?

It's also important to see your medical healthcare provider and/or mental health provider regularly following a traumatic event.

#### When should I go to the emergency department?

If you're experiencing thoughts of suicide or self-harm, it's crucial to contact 911, go to the nearest emergency department or call 988 to reach the Suicide and Crisis Lifeline.

### A note from Cleveland Clinic

It may be very difficult to seek professional help following a traumatic event. Know that treatment for acute stress disorder (ASD) is important, and with time, treatment can help you feel better. Talk to your healthcare provider about treatment options, and remember that they're available to help and support you.

### Care at Cleveland Clinic



Your mental well-being is just as important as your physical well-being. Cleveland Clinic's mental health experts can help you live life to the fullest.

Mental Health Care Make an Appointment

ពិនិត្យដោយវេជ្ជសាស្ត្រ

**ពិនិត្យចុងក្រោយនៅថ្ងៃទី 02/21/2023។**

ស្វែងយល់បន្ថែមអំពី បណ្ណាល័យសុខភាព និង ដំណើរការវិចារណកថា របស់យើង ។

ឯកសារយោង

ការណាត់ជួប
៨៦៦.៥៨៨.២២៦៤
ការណាត់ជួប និងទីតាំង
ស្នើសុំការណាត់ជួប

**សកម្មភាព**
ការណាត់ជួប និងការចូលប្រើប្រាស់
ការធានារ៉ាប់រងដែលទទួលយក
ប្រតិទិនព្រឹត្តិការណ៍
ជំនួយហិរញ្ញវត្ថុ
ផ្តល់ឱ្យទៅគ្លីនិក Cleveland
បង់វិក្កយបត្ររបស់អ្នកតាមអ៊ីនធឺណិត
យោងអ្នកជំងឺ
បញ្ជើឈ្មោះទូរសព្ទ
មតិទីពីរនិម្មិត
ការចូលមើលនិម្មិត

**ប្លុក ព័ត៌មាន និងកម្មវិធី**
ពិគ្រោះជាមួយ QD
សុខភាពសំខាន់
បន្ទប់ព័ត៌មាន
គ្លីនិក MyCleveland
គំនូសតាងរបស់ខ្ញុំ

**អំពីគ្លីនិក Cleveland**
100 ឆ្នាំនៃគ្លីនិក Cleveland
អំពីពួកយើង
ទីតាំង
គុណភាព និងសុវត្ថិភាពអ្នកជំងឺ
បទពិសោធន៍អ្នកជំងឺ
ការស្រាវជ្រាវ និងការច្នៃប្រឌិត
ការប្តេជ្ញាចិត្តសហគមន៍
អាជីព
សម្រាប់បុគ្គលិក
ធនធានសម្រាប់អ្នកជំនាញវេជ្ជសាស្ត្រ

**ព័ត៌មាន និងគោលការណ៍នៃគេហទំព័រ**
ផ្ញើមតិត្រឡប់មកយើង
ផែនទីគេហទំព័រ
អំពីគេហទំព័រនេះ។
រក្សាសិទ្ធិ បោះពុម្ពឡើងវិញ និងអាជ្ញាប័ណ្ណ
លក្ខខណ្ឌនៃការប្រើប្រាស់គេហទំព័រ
គោលការណ៍ឯកជនភាព
សេចក្តីជូនដំណឹងអំពីការអនុវត្តឯកជនភាព
សេចក្តីជូនដំណឹងអំពីការមិនរើសអើង

9500 Euclid Avenue, Cleveland, Ohio 44195 | 800.223.2273 | © 2025 គ្លីនិក Cleveland ។ រក្សាសិទ្ធិគ្រប់យ៉ាង។