

Model

The **inputs** are the real, continuous variable z_x , which standardises the increasing attempt numbers, x , of each individual and the dichotomous variable K_j which indicates whether the individual is a kid:

$$z_x = \frac{x - \bar{x}}{\hat{\sigma}_x}, \quad K_j = \begin{cases} 1 & \text{if } j \text{ is a kid} \\ 0 & \text{else} \end{cases} \quad (1)$$

where \bar{x} is the sample mean and $\hat{\sigma}_x$ the sample standard deviation of x .

The **outputs** are the reaction times, y_j , for each individual j , given the inputs. For our model, we use the $\log(y_j)$ and work with standardised values, $z_{\log(y_j)}$:

$$z_{\log(y_j)} = \frac{\log(y_j) - \overline{\log(y_j)}}{\hat{\sigma}_{\log(y_j)}} \quad (2)$$

The applied Bayesian **hierarchical model** expresses the probability of the output given the input and 9 parameters by redistributing the probability proportions as follows:

$$\begin{aligned} &P(z_x, K_j, \theta_{0_j}, \theta_{1_j}, \sigma, \mu_0, \varphi_0, \mu_1, \varphi_1, \tau_0, \tau_1 | z_{\log(y_j)}) \\ &\propto P(z_{\log(y_j)} | z_x, K_j, \theta_{0_j}, \theta_{1_j}, \sigma, \mu_0, \varphi_0, \mu_1, \varphi_1, \tau_0, \tau_1) \cdot P(\theta_{0_j}, \theta_{1_j}, \sigma, \mu_0, \varphi_0, \mu_1, \varphi_1, \tau_0, \tau_1) \\ &\propto P(z_{\log(y_j)} | z_x, \theta_{0_j}, \theta_{1_j}, \sigma) \cdot P(\theta_{0_j} | \mu_0, \varphi_0, \tau_0, K_j) \cdot P(\theta_{1_j} | \mu_1, \varphi_1, \tau_1, K_j) \cdot \\ &\quad P(\sigma) \cdot P(\mu_0) \cdot P(\mu_1) \cdot P(\varphi_0) \cdot P(\varphi_1) \cdot P(\tau_0) \cdot P(\tau_1) \end{aligned}$$

K_j and z_x are the outputs and have no priors. The joint posterior over all individuals can then be expressed as:

$$\prod_j^J P(z_x, K_j, \theta_{0_j}, \theta_{1_j}, \dots | z_{\log(y_j)}) \propto \prod_j^J P(z_{\log(y_j)} | z_x, \theta_{0_j}, \theta_{1_j}, \sigma) \cdot P(\theta_{0_j} | \mu_0, \varphi_0, \tau_0, K_j) \cdot P(\theta_{1_j} | \mu_1, \varphi_1, \tau_1, K_j) \cdot \dots$$

A graphical representation of the hierarchical model can be seen in Figure 2.

The **core** of the model is the distribution of the data (i.e. the likelihood function) which models a regression line for each individual with a certain noise around it:

$$z_{\log(y_j)} \sim \mathcal{N}(\theta_{0_j} + \theta_{1_j} \cdot z_x, \sigma^2)$$

The assumed shapes of the intermediate priors and pure (hyper-) priors are as follows:

$$\begin{aligned} \theta_{0_j} &\sim \mathcal{N}(\mu_0 + \varphi_0 \cdot K_j, \tau_0^2) \\ \theta_{1_j} &\sim \mathcal{N}(\mu_1 + \varphi_1 \cdot K_j, \tau_1^2) \\ \mu_0, \mu_1, \varphi_0, \varphi_1 &\sim \mathcal{U}(-\infty, +\infty) \\ \sigma, \tau_0, \tau_1 &\sim \mathcal{U}[0, +\infty) \end{aligned}$$

Note, that the priors in the first line above exclude 0.

Transformations

With the random variables $\Xi, \Theta, \Omega \sim \mathcal{N}(0, 1)$, we can rewrite the distributions as follows:

$$z_{\log(y_j)} = \theta_{0_j} + \theta_{1_j} \cdot z_x + \sigma \cdot \Xi$$

$$\theta_{0_j} = \mu_0 + \varphi_0 \cdot K_j + \tau_0 \cdot \Theta$$

$$\theta_{1_j} = \mu_1 + \varphi_1 \cdot K_j + \tau_1 \cdot \Omega$$

$$z_{\log(y_j)} = \mu_0 + \varphi_0 \cdot K_j + \tau_0 \cdot \Theta + (\mu_1 + \varphi_1 \cdot K_j + \tau_1 \cdot \Omega) z_x + \sigma \cdot \Xi$$

For the **individual** level, we keep the θ s and do not replace them with their priors:

$$z_{\log(y_j)} = \theta_{0_j} + \theta_{1_j} \cdot \frac{x - \bar{x}}{\hat{\sigma}_x} + \sigma \cdot \Xi$$

$$z_{\log(y_j)} = \theta_{0_j} - \theta_{1_j} \frac{\bar{x}}{\hat{\sigma}_x} + \frac{\theta_{1_j}}{\hat{\sigma}_x} x + \sigma \cdot \Xi$$

$$\log(y_j) = \underbrace{\hat{\sigma}_y \theta_{0_j} - \hat{\sigma}_y \theta_{1_j} \frac{\bar{x}}{\hat{\sigma}_x} + \overline{\log(y_j)}}_{\theta_{0_j, \text{unsc}}} + \underbrace{\hat{\sigma}_y \frac{\theta_{1_j}}{\hat{\sigma}_x} x}_{\theta_{1_j, \text{unsc}}} + \underbrace{\hat{\sigma}_y \sigma \cdot \Xi}_{\sigma_{\text{unsc}}}$$

$$\log(y_j) \sim \mathcal{N}(\theta_{0_j, \text{unsc}} + \theta_{1_j, \text{unsc}} \cdot x, \sigma_{\text{unsc}}^2)$$

$$E[y_j] = e^{\theta_{0_j, \text{unsc}} + \theta_{1_j, \text{unsc}} \cdot x + \frac{\sigma_{\text{unsc}}^2}{2}}$$

$$= \underbrace{e^{\theta_{1_j, \text{unsc}} \cdot x}}_{\text{effect of increasing attempts}} \cdot e^{\theta_{0_j, \text{unsc}} + \frac{\sigma_{\text{unsc}}^2}{2}}$$

For the **group** level, we can convolute the addition of the random variables Ξ, Θ and Ω assuming independence and merge them into a single one which we call again Ξ :

$$z_{\log(y_j)} = \mu_0 + \varphi_0 \cdot K_j + (\mu_1 + \varphi_1 \cdot K_j) z_x + \sqrt{\sigma^2 + \tau_0^2 + \tau_1^2 z_x^2} \cdot \Xi$$

De-standardising from Equations 1 and 2 yields:

$$z_{\log(y_j)} = \mu_0 - \mu_1 \frac{\bar{x}}{\hat{\sigma}_x} + \left(\varphi_0 - \varphi_1 \frac{\bar{x}}{\hat{\sigma}_x} \right) K_j + \left(\frac{\mu_1}{\hat{\sigma}_x} + \frac{\varphi_1}{\hat{\sigma}_x} \cdot K_j \right) x + \sqrt{\sigma^2 + \tau_0^2 + \tau_1^2 \frac{\bar{x}^2}{\hat{\sigma}_x^2} - 2 \frac{\tau_1^2 \bar{x}}{\hat{\sigma}_x^2} x + \frac{\tau_1^2}{\hat{\sigma}_x^2} x^2} \cdot \Xi$$

$$\log(y_j) = \underbrace{\hat{\sigma}_y \mu_0 - \hat{\sigma}_y \mu_1 \frac{\bar{x}}{\hat{\sigma}_x} + \overline{\log(y_j)}}_{\mu_{0, \text{unsc}}} + \underbrace{\hat{\sigma}_y \left(\varphi_0 - \varphi_1 \frac{\bar{x}}{\hat{\sigma}_x} \right) \cdot K_j}_{\varphi_{0, \text{unsc}}} + \left(\underbrace{\hat{\sigma}_y \frac{\mu_1}{\hat{\sigma}_x}}_{\mu_{1, \text{unsc}}} + \underbrace{\hat{\sigma}_y \frac{\varphi_1}{\hat{\sigma}_x} \cdot K_j}_{\varphi_{1, \text{unsc}}} \right) x + \sqrt{\underbrace{\hat{\sigma}_y^2 \sigma^2}_{\sigma_{\text{unsc}}^2} + \underbrace{\hat{\sigma}_y^2 \left(\tau_0^2 + \tau_1^2 \frac{\bar{x}^2}{\hat{\sigma}_x^2} \right)}_{\tau_{0, \text{unsc}}^2} + \underbrace{\left(-2 \hat{\sigma}_y^2 \frac{\tau_1^2 \bar{x}}{\hat{\sigma}_x^2} \right) x}_{\tilde{\tau}_{1, \text{unsc}}^2} + \underbrace{\hat{\sigma}_y^2 \frac{\tau_1^2}{\hat{\sigma}_x^2} x^2}_{\tau_{1, \text{unsc}}^2}} \cdot \Xi$$

$$\log(y_j) \sim \mathcal{N}(\mu_{0, \text{unsc}} + \varphi_{0, \text{unsc}} \cdot K_j + (\mu_{1, \text{unsc}} + \varphi_{1, \text{unsc}} \cdot K_j) x, \sigma_{\text{unsc}}^2 + \tau_{0, \text{unsc}}^2 + \tilde{\tau}_{1, \text{unsc}}^2 x + \tau_{1, \text{unsc}}^2 x^2)$$

$$E[y_j] = e^{\mu_{0, \text{unsc}} + \varphi_{0, \text{unsc}} \cdot K_j + (\mu_{1, \text{unsc}} + \varphi_{1, \text{unsc}} \cdot K_j) x + \frac{\sigma_{\text{unsc}}^2}{2} + \frac{\tau_{0, \text{unsc}}^2}{2} + \frac{\tilde{\tau}_{1, \text{unsc}}^2 x}{2} + \frac{\tau_{1, \text{unsc}}^2 x^2}{2}}$$

$$= e^{\mu_{0, \text{unsc}} + \varphi_{0, \text{unsc}} \cdot K_j + \frac{\sigma_{\text{unsc}}^2}{2} + \frac{\tau_{0, \text{unsc}}^2}{2}} \cdot \underbrace{e^{(\mu_{1, \text{unsc}} + \varphi_{1, \text{unsc}} \cdot K_j) x + \frac{\tilde{\tau}_{1, \text{unsc}}^2 x}{2} + \frac{\tau_{1, \text{unsc}}^2 x^2}{2}}}_{\text{effect of increasing attempts}}$$

$$= \underbrace{e^{\varphi_{0, \text{unsc}} \cdot K_j}}_{\text{child-effect on "intercept"}} e^{\mu_{0, \text{unsc}} + \frac{\sigma_{\text{unsc}}^2}{2} + \frac{\tau_{0, \text{unsc}}^2}{2}} \cdot \underbrace{e^{\varphi_{1, \text{unsc}} \cdot K_j x}}_{\text{child-effect on "slope"}} e^{\mu_{1, \text{unsc}} x + \frac{\tilde{\tau}_{1, \text{unsc}}^2 x}{2} + \frac{\tau_{1, \text{unsc}}^2 x^2}{2}}$$

The multiplicative effect of being a kid acts only on the group level, but the effect of increasing attempts acts on the individual and group level.