

EE23010 NCERT Exemplar

Vishal A - EE22BTECH11057

Question 23.2023

Let $X_1, X_2, X_3, \dots, X_n$ be a random sample of size $n (\geq 2)$ from a population having probability density function

$$f(x; \theta) = \begin{cases} \frac{2}{\theta x} (\log_e x) e^{-\frac{(\log_e x)^2}{\theta}} & , 0 < x < 1 \\ 0 & , otherwise \end{cases}$$

where $\theta > 0$ is an unknown parameter. The maximum likelihood estimator of θ is,

Solution:

$$L(\theta; X_1, X_2, \dots, X_n) = f(\theta; X_1, X_2, \dots, X_n) \quad (1)$$

Maximizing $L(\theta)$ is equivalent to maximizing the $\log L(\theta)$ as \log is a monotonically increasing function.

$$l(\theta) = \log L(\theta) \quad (2)$$

$$= \log \left(\prod_{i=1}^n f(X_i | \theta) \right) \quad (3)$$

$$= \sum_{i=1}^n \log f(X_i | \theta) \quad (4)$$

$$= -n \ln 2 - n \ln \theta + \sum_{i=1}^n \ln(-\ln x_i) - \sum_{i=1}^n (\ln X_i) - \sum_{i=1}^n \frac{(\log_e X_i)^2}{\theta} \quad (5)$$

Maximizing $l(\theta)$ with respect to θ gives the MLE estimation, therefore

$$\frac{\partial l(\theta)}{\partial \theta} = 0 \quad (6)$$

$$\frac{-n}{\theta} + \frac{1}{(\theta)^2} \sum_{i=1}^n (\log_e X_i)^2 = 0 \quad (7)$$

$$\theta = \frac{1}{n} \sum_{i=1}^n (\log_e X_i)^2 \quad (8)$$