

# Environmental Context Dependency in Species Interactions

*Owen Liu*

*19 July, 2018*

**Primary target journal:** Nature Ecology and Evolution

## **Journal Guidelines:**

- Maximum 3500 words main text (excluding introductory paragraph/abstract)
- Maximum 6 display items (figures/tables)
- The introductory paragraph is typically 150 words and is unreferenced; it contains a brief account of the background and rationale of the work, followed by a statement of the main conclusions introduced by the phrase “Here we show” or some equivalent. An introduction (without heading) of up to 500 words of referenced text expands on the background of the work (some overlap with the summary is acceptable), and is followed by a concise, focused account of the findings (headed ‘Results’), and one or two short paragraphs of discussion (headed ‘Discussion’).

## **Work confirmed/supported/debates:**

- Tegner et al 1997, Dayton et al 1999
- Bottom-up vs. top-down forcing? Disturbance or herbivory?
- Competition between urchin species?
- Physical drivers increase predictability of algae models!

## **Suggested future research**

- Predominance of mutualism- why the positive interactions of Pterygophora and Macro-cystis?
- Urchin grazing pressure as most important??
- Multiple stressors (e.g. herbivory, sst, wave height)

## **FIGURES**

Time series of study species Example attractor? (in supplementary) CCM results

---

## **Introductory paragraph (abstract)**

Ecological interactions are not uniform across time, and instead vary with environmental conditions. Interactions among species are often measured with short-term controlled experiments, but these experiments are subject to the particular environmental conditions

under which they are performed. As an alternative, we utilize empirical dynamic modeling applied to a 30-year time series to estimate species interaction strengths across a wide range of environmental conditions in a coastal marine ecosystem. By including large-scale climate indices, sea surface temperature, and a measure of physical disturbance in analyses, we show that environmental context influences the strength and direction of species interactions. In so doing, we are able to confirm and extend results from previous studies, as well as identify potentially important but understudied interactions. The significant context dependency in species interactions found in this study argues for a greater utilization of long-term data and empirical dynamic modeling in studies of ecosystem dynamics.

## 42 Introduction

Interactions between species drive patterns of diversity, stability, resilience, and productivity in nature<sup>1-4</sup>. In any ecosystem, the collection of species interactions determines community dynamics. Until recently, most studies viewed these dynamics—e.g., the bleaching and recovery of a coral reef, or the assembly and disassembly of terrestrial plant communities—as processes resulting from static, predictable species interactions. However, the observation that species interactions are not spatiotemporally uniform<sup>5-8</sup> calls into question assumptions of interaction stability.

Ecologists recognize now that important species interactions may vary over time, but this context dependency remains difficult to measure and describe. Experiments that measure interactions are generally performed over a limited spatiotemporal range, and are therefore subject to a specific environmental context that may not encompass the range of conditions experienced by that ecosystem over longer time scales<sup>9</sup>. This is worrying, since environmental context can profoundly influence the outcome of species interactions ranging anywhere from keystone predation<sup>7</sup> to competition<sup>5,10,11</sup>, to protective symbioses<sup>12-14</sup>.

Moreover, the focus of the search for context dependency has been on mean interaction strengths, at the expense of specific examinations of interaction variance<sup>8</sup>. This focus may be misguided, as it has been shown that interactions that are variable in magnitude and direction—and therefore “weak” when averaged—may actually be some of the most important in driving community dynamics<sup>4</sup>. If key species interactions are variable in this way across environmental gradients, then many studies may be attributing important ecological phenomena to observational noise.

A solution to these difficulties is to a) utilize ecological observations collected over a long time period, across a large range of environmental contexts, with b) an analytical method to directly estimate context-dependent species interactions from those observations. Such an approach could help to characterize environmental contingencies in species interactions and explicitly examine interaction variability. Here, we use empirical dynamic modelling (EDM,<sup>15</sup>) to estimate a varying species interaction network and establish environmental context dependency in interaction strength and direction. Empirical dynamic modelling uses information from single or multiple time series to empirically model relationships between variables through the reconstruction of dynamic attractors (<https://youtu.be/8DikuwwPWsY>). The general

modelling framework for all EDM methods is readily adaptable to many different sorts of time series variables, including environmental variables manifesting at different scales<sup>16–18</sup>. Because the methods are specifically designed for nonlinear dynamic systems, EDM—in theory—should be able to illuminate context-dependent patterns in species interactions.

Recently-developed EDM methods exist for uncovering dynamic species interactions from time series data<sup>15</sup>, but these methods have insofar been applied only to simulated and planktonic communities, and their utility to the study of other ecological systems remains untested. We focus here instead on giant kelp forests in southern California, a diverse and temporally dynamic ecosystem in which many important species interactions are well-documented<sup>19–21</sup>. The study of kelp forests has been foundational to ecological theory, especially regarding the relative influence of top-down and bottom-up structuring forces in ecosystems<sup>22–26</sup>. Recently, however, findings from long-term kelp forest research programs have begun to challenge many long-held beliefs about the drivers of kelp forest ecosystem dynamics<sup>27</sup>. In particular, a longer-term perspective has led to a recognition of the critical importance of environmental context—such as level of physical disturbance or the current state of El Niño conditions—for understanding kelp forest processes<sup>28–31</sup>. In this study we utilize monitoring data from one such effort at San Nicolas Island, a small, remote member of the California Channel Islands in the northeast Pacific<sup>32</sup>.

To characterize environmental context dependency in kelp forest interactions between species, we take three general steps (see Methods). First, we use empirical dynamic modeling causality tests called convergent cross-mapping<sup>33</sup> to construct a kelp forest species interaction network directly from time series data. In so doing, we test for all unidirectional causal signals between five common kelp forest species, as well as between five exogenous environmental variables and those species. Second, for each identified causal link between species, we reconstruct the actual species interactions over time, using another EDM tool called multivariate S-maps (sequential locally weighted global linear maps)<sup>15,34</sup>. Finally, we show how variability in key species interactions can be related back to the environmental context under which they took place.

## Results and discussion

Our analyses focus on the dynamics of five common southern California kelp forest species, whose interactions are thought to be important in structuring kelp forest ecosystems<sup>19,21,35</sup>. The giant kelp *Macrocystis pyrifera* is the eponymous foundation species<sup>36</sup>, the primary canopy- and habitat-forming kelp along most of the central and southern coast of California<sup>20</sup>. The monitoring data include young *Macrocystis* recruits (sporophytes identified as *Macrocystis* but less than 1 meter tall<sup>32</sup>). We explore *Macrocystis* dynamics and its interactions with two presumptive competitors and two abundant herbivores. The understory kelp species *Laminaria farlowii* and *Pterygophora californica* compete with *Macrocystis* for space, light, and nutrients<sup>37–39</sup>. The two herbivores—the purple sea urchin *Strongylocentrotus purpuratus* and the red sea urchin *Mesocentrotus franciscanus*—are thought in many places to control *Macrocystis* density and can sometimes wipe out entire giant kelp forests, leading to the alternative ecosystem state known as an urchin barren<sup>40,41</sup>.


114 In southern California, *Macrocystis* population dynamics can be driven by nutrient availability  
115 and physical disturbance<sup>26,30</sup>. The availability of nitrate is inversely related to seawater  
116 temperature<sup>42</sup> and, over longer time scales, is associated with oscillations in patterns of  
117 upwelling and oceanic currents. Accordingly, we include five environmental variables in our  
118 analyses to test their relationship to kelp forest species dynamics and interactions: sea surface  
119 temperature (SST), physical disturbance (measured by maximum seasonal wave height,  
120 SWH), and three indices of low-frequency climate modes: the Multivariate El Niño Index  
121 (MEI)<sup>43</sup>, the Pacific Decadal Oscillation (PDO)<sup>44</sup>, and the North Pacific Gyre Oscillation  
122 (NPGO)<sup>45</sup>.

123 Applying convergent cross mapping<sup>33</sup> (CCM) to the set of six biological and five physical  
124 variables, we find a relatively dense interaction network (Figure 1). Out of 90 possible  
125 unidirectional links between species and between species and the environmental variables,  
126 40 are significant. Adult *Macrocystis* density is driven by all five environmental variables,  
127 with SWH, SST, and the NPGO showing the strongest causal signals. This finding confirms  
128 recent work by others using different methods<sup>30</sup> that showed that these same three variables  
129 were the primary controls of giant kelp biomass dynamics across the California coast. More  
130 generally, although the included physical variables show significant links to many of the  
131 biological variables, the NPGO and SWH show the strongest links to almost all of the  
132 biological variables. Our analysis suggests that more attention should be focused on the  
133 effects of the NPGO.

134 Interestingly, in turn, adult *Macrocystis* shows strong causal links to every other biological  
135 variable. This is despite the fact that the study site at San Nicolas Island does not have  
136 a stable giant kelp forest (see raw time series, Figure 6); rather, the site has transitioned  
137 from an urchin barren<sup>41</sup> to a *Pterygophora* and *Laminaria*-dominated state, to a *Macrocystis*  
138 forest at various times throughout the 30-year time series. The implication is that, despite  
139 not maintaining dominance in the typical ecological sense of word (large abundance and  
140 biomass), *Macrocystis* is “dynamically dominant”, and remains the key foundation species in  
141 this ecosystem. Its dynamics are fundamentally important in driving the dynamics of all the  
142 other kelp forest species<sup>20</sup>.

143 CCM analysis confirms that the system studied represents a complex array of significant  
144 interactions between algal species and their herbivores. However, CCM alone does not  
145 elucidate the direction and magnitude of species interactions. To obtain estimates of the  
146 interactions themselves, we use multivariate S-maps<sup>15,34</sup> (see Methods). S-maps reconstructs  
147 dynamic “attractors” by casting the abundances of causally-related species into state space.  
148 For a set of causally-related species, a point in multivariate space can be plotted using each  
149 species’ abundance as an axis. The attractor is then created by tracing this multispecies  
150 trajectory forward in time (see example attractor in Figure 5). For each point along the  
151 attractor, S-maps computes a Jacobian matrix, the elements of which are the estimated  
152 partial derivatives between species. These interaction matrix elements are our measure of  
153 species interactions. Because Jacobians are computed sequentially for every point along  
154 reconstructed attractors, we obtain estimates of interaction strength that vary with ecosystem  
155 state.

156 There are two important aspects of S-maps that deserve mention. First, the S-map estimation






Figure 2: Distribution of estimated interaction by type

procedure, like all EDM methods, is specifically designed for nonlinear systems, and is therefore an appropriate tool for investigating ecosystems exhibiting nonlinear dynamics such as alternative stable states or hysteresis<sup>15</sup>. Secondly, because S-maps utilizes reconstructed multispecies attractors, each estimated interaction is fundamentally based on observations of similar past ecosystem states—in our case represented as multivariate collections of causally-linked species’ densities—rather than a phenomenological extrapolation of the most recent dynamics. In simple terms, instead of asking, “What is our prediction for the strength of herbivory based on last year’s observed dynamics,” S-maps is concerned with, “What is our best estimate for the strength of herbivory, based on our knowledge of times in the past when the ecosystem was most similar to today?”

In the San Nicolas kelp forest, we find a striking prevalence of neutral and positive species interactions (Figure 2). After grouping species interactions by type, only herbivory (the effect of urchins on algae species) is predominantly negative. Conversely, the effect of the algae species on the urchins has the flattest distribution, with occasional strong negative and strong positive interactions. Likewise, contrary to our expectations, interactions between the algae species and between the urchin species are not always antagonistic. These results suggest that facilitation is common in kelp forests when context is considered. Since there is evidence here of strong herbivory, for example, then the indirect facilitative effects between algae species




Figure 3: Mean and standard deviation of each estimated unidirectional species interaction

may sometimes outweigh their direct competition. Additionally, *Macrocystis* forests can mediate current strength and as a result, can also help retain reproductive propagules of other species<sup>20</sup>. These types of indirect facilitation in kelp forests have received little attention[46; Benes2015] relative to a strong focus on exploitative competition between these species for light and nutrients<sup>37,39</sup>, but similar effects have been documented in other systems<sup>???</sup>. Our analysis does not contradict the importance of competition in kelp forest ecosystems, but suggests that facilitation, especially indirect facilitation, may be an important structuring factor in kelp forest ecosystems.

- CCM results and variable interaction network- centrality of *Macrocystis*
- Mean vs. variance in certain algal interactions?
- Bidirectional mean interactions (across two dimensions, A on B and B on A)?
- Frequency of positive vs. negative interactions? Across environmental gradients?




Figure 4: Macrocystis effect on Pterygophora






Figure 5: Example reconstructed dynamic attractor




Figure 6: Raw data for species and physical drivers, normalized to zero mean and unit variance

## 187 Methods

## 188 References

- 189 1. Connell, J. H. Diversity in Tropical Rain Forests and Coral Reefs. *Science* **199**, 1302–1310  
190 (1978).
- 191 2. Suttle, K. B., Thomsen, M. A. & Power, M. E. Species Interactions Reverse Grassland  
192 Responses to Changing Climate. *Science* **315**, 640–642 (2007).
- 193 3. Tilman, D. The ecological consequences of changes in biodiversity: a search for general  
194 principles. *Ecology* **80**, 1455–1474 (1999).
- 195 4. Berlow, E. L. Strong effects of weak interactions in ecological communities. *Nature* **398**,  
196 330–334 (1999).
- 197 5. Odadi, W. O., Karachi, M. K., Abdulrazak, S. A. & Young, T. P. African wild ungulates  
198 compete with or facilitate cattle depending on season. *Science* **333**, 1753–1755 (2011).
- 199 6. Pennings, S. C. & Silliman, B. R. Linking Biogeography and Community Ecology :  
200 Latitudinal Variation in Plant-Herbivore Interaction Strength. *Ecology* **86**, 2310–2319 (2005).
- 201 7. Menge, B. A., Berlow, E. L., Blanchette, C. A., Navarrete, S. A. & Yamada, B. The  
202 Keystone Species Concept : Variation in Interaction Strength in a Rocky Intertidal Habitat.  
203 *Ecology* **64**, 249–286 (1994).
- 204 8. Chamberlain, S. A., Bronstein, J. L. & Rudgers, J. A. How context dependent are species  
205 interactions? *Ecology Letters* **17**, 881–890 (2014).
- 206 9. Vaughn, K. J. & Young, T. P. Contingent Conclusions: Year of Initiation Influences  
207 Ecological Field Experiments, but Temporal Replication is Rare. *Restoration Ecology* **18**,  
208 59–64 (2010).
- 209 10. Lima, M., Ernest, S. K. M., Brown, J. H., Belgrano, A. & Stenseth, N. C. Chihuahuan  
210 Desert Kangaroo Rats: Nonlinear Effects of Population Dynamics, Competition, and Rainfall.  
211 *Ecology* **89**, 2594–2603 (2008).
- 212 11. Holzapfel, C. & Mahall, B. E. Bidirectional Facilitation and Interference between Shrubs  
213 and Annuals in the Mojave Desert. *Ecology* **80**, 1747–1761 (1999).
- 214 12. Daskin, J. H. & Alford, R. A. Context-dependent symbioses and their potential roles  
215 in wildlife diseases. *Proceedings of the Royal Society B: Biological Sciences* **279**, 1457–1465  
216 (2012).
- 217 13. Glynn, P. W. Coral reef bleaching: Facts, hypotheses and implications. *Global Change*  
218 *Biology* **2**, 495–509 (1996).
- 219 14. White, J. D., Sarnelle, O. & Hamilton, S. K. Unexpected population response to increasing  
220 temperature in the context of a strong species interaction. *Ecological Applications* (2017).

doi:10.1002/eap.1558

15. Deyle, E. R., May, R. M., Munch, S. B. & Sugihara, G. Tracking and forecasting ecosystem interactions in real time. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20152258 (2016).

16. Deyle, E. R. *et al.* Predicting climate effects on Pacific sardine. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 6430–6435 (2013).

17. Chang, C. W., Ushio, M. & Hsieh, C. hao. Empirical dynamic modeling for beginners. *Ecological Research* 1–12 (2017). doi:10.1007/s11284-017-1469-9

18. Ye, H., Deyle, E. R., Gilarranz, L. J. & Sugihara, G. Distinguishing time-delayed causal interactions using convergent cross mapping. *Scientific reports* **5**, 14750 (2015).

19. Foster, M. S. & Schiel, D. R. Ecology of giant kelp forests in California: a community profile. *United States Fish and Wildlife Service Biological Report* **85**, 1–152 (1985).

20. Graham, M. H., Vásquez, J. A. & Buschmann, A. H. Global Ecology of the Giant Kelp *Macrocystis* : From Ecotypes To Ecosystems. *Oceanography and Marine Biology* **45**, 39–88 (2007).

21. Teagle, H., Hawkins, S. J., Moore, P. J. & Smale, D. A. The role of kelp species as biogenic habitat formers in coastal marine ecosystems. *Journal of Experimental Marine Biology and Ecology* **492**, 81–98 (2017).

22. Estes, J. A. & Palmisano, J. F. Sea otters: their role in structuring nearshore communities. *Science* **185**, 1058–1060 (1974).

23. Graham, M. *et al.* Population dynamics of giant kelp *Macrocystis pyrifera* along a wave exposure gradient. *Marine Ecology Progress Series* **148**, 269–279 (1997).

24. Kvitek, R. G., Iampietro, P. J. & Bowlby, C. E. Sea Otters and Benthic Prey Communities : a Direct Test of the Sea Otter As Keystone Predator in Washington State. *Marine Mammal Science* **14**, 895–902 (1998).

25. Foster, M. S. & Schiel, D. R. Loss of predators and the collapse of southern California kelp forests (?): Alternatives, explanations and generalizations. *Journal of Experimental Marine Biology and Ecology* **393**, 59–70 (2010).

26. Reed, D. C. *et al.* Wave disturbance overwhelms top-down and bottom-up control of primary production in California kelp forests. *Ecology* **92**, 2108–2116 (2011).

27. Reed, D. *et al.* Extreme warming challenges sentinel status of kelp forests as indicators of climate change. *Nature Communications* **7**, 13757 (2016).

28. Parnell, P. E. *et al.* The response of giant kelp (*Macrocystis pyrifera*) in southern California to low-frequency climate forcing. *Limnology and Oceanography* **55**, 2686–2702 (2010).

29. Byrnes, J. E. *et al.* Climate-driven increases in storm frequency simplify kelp forest food

- 257 webs. *Global Change Biology* **17**, 2513–2524 (2011).
- 258 30. Bell, T. W., Cavanaugh, K. C., Reed, D. C. & Siegel, D. A. Geographical variability in  
259 the controls of giant kelp biomass dynamics. *Journal of Biogeography* **42**, 2010–2021 (2015).
- 260 31. Young, M. A. *et al.* Environmental controls on spatial patterns in the long-term  
261 persistence of giant kelp in central California. *Ecology* **86**, 45–60 (2015).
- 262 32. Kenner, M. C. *et al.* A multi-decade time series of kelp forest community structure at  
263 San Nicolas Island, California (USA). *Ecology* **94**, 2654–2654 (2013).
- 264 33. Sugihara, G. *et al.* Detecting causality in complex ecosystems. *Science* **338**, 496–500  
265 (2012).
- 266 34. Sugihara, G. Nonlinear forecasting for the classification of natural time series. *Philo-*  
267 *sophical Transactions of the Royal Society of London Series A-Mathematical Physical and*  
268 *Engineering Sciences* **348**, 477–495 (1994).
- 269 35. Foster, M. S. Organization of macroalgal assemblages in the Northeast Pacific: the  
270 assumption of homogeneity and the illusion of generality. *Hydrobiologia* **192**, 21–33 (1990).
- 271 36. Dayton, P. Toward an understanding of resilience and the potential effects of enrichments  
272 to the benthos at McMurdo Sound, Antarctica. 96 (1972).
- 273 37. Dayton, P. K. *et al.* Patch dynamics and stability of some California USA kelp  
274 communities. *Ecological Monographs* **54**, 253–290 (1984).
- 275 38. Dayton, P. K., Tegner, M. J., Edwards, P. B. & Riser, K. L. Temporal and spatial scales  
276 of kelp demography: the role of oceanographic climate. *Ecological Monographs* **69**, 219–250  
277 (1999).
- 278 39. Reed, D. C. & Foster, M. S. The effects of canopy shadings on algal recruitment and  
279 growth in a giant kelp forest. *Ecology* **65**, 937–948 (1984).
- 280 40. Ling, S. *et al.* Global regime shift dynamics of catastrophic sea urchin overgrazing. *Phil.*  
281 *Trans. R. Soc. B* **370**, 20130269 (2014).
- 282 41. Harrold, C. & Reed, D. C. Food Availability , Sea Urchin Grazing , and Kelp Forest  
283 Community Structure. *Ecology* **66**, 1160–1169 (1985).
- 284 42. Zimmerman, R. C. & Kremer, J. N. Episodic nutrient supply to a kelp forest ecosystem  
285 in Southern California. *Journal of Marine Research* **42**, 591–604 (1984).
- 286 43. Wolter, K. & Timlin, M. S. El Niño/Southern Oscillation behaviour since 1871 as  
287 diagnosed in an extended multivariate ENSO index (MEI.ext). *International Journal of*  
288 *Climatology* **31**, 1074–1087 (2011).
- 289 44. Mantua, N. J. & Hare, S. R. The Pacific Decadal Oscillation. **58**, 35–44 (2002).
- 290 45. Di Lorenzo, E. *et al.* North Pacific Gyre Oscillation links ocean climate and ecosystem  
291 change. *Geophysical Research Letters* **35**, 2–7 (2008).
- 292 46. Arkema, K. K., Reed, D. C. & Schroeter, S. C. Direct and indirect effects of giant kelp

**293** determine benthic community structure and dynamics. *Ecology* **90**, 3126–3137 (2009).