

Ammended Claims

Dated : 27.04.2011

EP 07 011 162.0

1

Claims

1. A composition for prophylaxis or treatment of urethritis or cystitis, comprising a matrix metalloproteinase inhibitor, X
- ~~2.~~ ³ The composition of claim 1, wherein the matrix metalloproteinase inhibitor is selected from the group consisting of ~~Zizyphus jujuba, Malva sylvestris,~~ Opuntia, Streptacantha, ~~Cinnamomi Cortex,~~ Hippophae rhamnoides, Panax ginseng, Eriobotrya japonica, Citrus depressa Hayata, ~~Nelumbo nucifera,~~ Mosla chinensis, Fructus schisandrae, Dalbergia odorifera, Smilax glabra, kale and their extracts. X
2. ~~3.~~ The composition of claim 1, wherein the matrix metalloproteinase inhibitor is selected from the group consisting of inhibitors of MMP-2 and MMP-9. X
3. ~~4.~~ The composition of claim 1, which further comprises at least one component selected from the group consisting of a calcineurin enzyme activator, cranberry, and a urine pH stabilizer. X
4. ~~5.~~ ³ The composition of claim ~~4~~, wherein the calcineurin enzyme activator is selected from the group consisting of Lonicera japonica Thunb., Cordeceps sinensis, Sophorae Radix, Hippophae rhamnoides, Flos Lonicerae, Pulsatillae Radix, extract of the dried leaf of Camellia sinensis (L.) C. Kuntze, Ganoderma lucidum, Gynostemma pentaphyllum, Cynara Scolymus L, Thyme, Fructus Schisandrae, Prunellae spica, Lophatheri spica, Coptis Chinensis Franchet, Dandelion (Taraxacum officinale F. Weber ex Wiggers), ~~Portulacae Herba,~~ Gardeniae Fructus, Sophora subprostrata Radix, Houlttuynia cordata Thum, Forsythia suspense (Thum) Vahl, Smilax glabra and Coptis groenlandica Salisb. X

- 5 ~~8~~. The composition of claim ~~4~~³, wherein the urine pH stabilizer is selected from the group consisting of glycolic acid, gallic acid, ellagic acid, vanillic acid, citric acid, malic acid, tartaric acid, lactic acid, and vitamin C. X
- 6 ~~7~~. The composition of claim ~~4~~³, which comprises a calcineurin enzyme activator. X
- 7 ~~8~~. The composition of claim ~~7~~⁶, wherein ratio of weight percent of the matrix metalloproteinase inhibitor to the calcineurin enzyme activator is between 10:1 and 1:5. X
- 8 ~~9~~. The composition of claim 1, which contains pharmaceutically acceptable excipients, carriers or diluents. X
- 9 ~~10~~. Use of a composition for the preparation of a medicament for the prophylaxis or treatment of urethritis or cystitis, wherein the composition comprises a matrix metalloproteinase inhibitor, X
- ~~11. The use of claim 10, wherein the matrix metalloproteinase inhibitor is selected from the group consisting of Zizyphus-jujuba, Malva-sylvestris, Opuntia Streptacantha, Cinnamomi-Cortex, Hippophae rhamnoides, Panax ginseng, Eriobotrya japonica, Citrus depressa Hayata, Nelumbo-nucifera, Mosla chinensis, Fructus schisandrae, Dalbergia odorifera, Smilax glabra, kale and their extracts.~~ X
X
X
X
- 10 ~~12~~. The use of claim ~~10~~³, wherein the matrix metalloproteinase inhibitor is selected from the group consisting of inhibitors of MMP-2 and MMP-9. X

- 11 ⁹ 13. The use of claim ~~10~~, wherein the composition further comprises at least one component selected from the group consisting of a calcineurin enzyme activator, cranberry, and a urine pH stabilizer. X
- 12 ¹¹ 14. The use of claim ~~13~~, wherein the calcineurin enzyme activator is selected from the group consisting of *Lonicera japonica* Thunb., *Cordeceps sinensis*, *Sophorae Radix*, *Hippophae rhamnoides*, *Flos Lonicerae*, *Pulsatillae Radix*, extract of the dried leaf of *Camellia sinensis* (L.) C. Kuntze, *Ganoderma lucidum*, *Gynostemma pentaphyllum*, *Cynara Scolymus* L, Thyme, *Fructus Schisandrae*, *Prunellae spica*, *Lophatheri spica*, *Coptis Chinensis* Franchet, Dandelion (*Taraxacum officinale* F. Weber ex Wiggers), ~~Portulacae—Herba,~~ *Gardeniae Fructus*, *Sophora subprostrata Radix*, *Houttuynia cordata* Thum, *Forsythia suspense* (Thum) Vahl, *Smilax glabra* and *Coptis groenlandica* Salisb. X
- 13 ¹¹ 15. The use of claim ~~13~~, wherein the urine pH stabilizer is selected from the group consisting of glycolic acid, gallic acid, ellagic acid, vanillic acid, citric acid, malic acid, tartaric acid, lactic acid, and vitamin C. X
- 14 ⁹ 16. The use of claim ~~10~~, wherein the composition contains pharmaceutically acceptable excipients, carriers or diluents. X