

11-8-2011

CLAIMS

1. — ~~An eriocitrin-containing material which is obtained by separating eriocitrin from an eriocitrin-containing citrus extract prepared from a citrus fruit using an extractant, the eriocitrin-containing material being characterized in that the eriocitrin separation from the citrus extract comprises: bringing the citrus extract into contact with a porous synthetic adsorption resin such that eriocitrin in the citrus extract is adsorbed on the porous synthetic adsorption resin, the porous synthetic adsorption resin comprising a phenol-formaldehyde resin as a main framework and having amino and phenolic hydroxyl groups; and then eluting the eriocitrin adsorbed on the porous synthetic adsorption resin using an elution solvent.~~
2. — ~~The eriocitrin-containing material according to claim 1, wherein the elution solvent is hydrous ethanol having an ethanol concentration of 20 to 90% by volume.~~
3. — ~~The eriocitrin-containing material according to claim 1 or 2, wherein the porous synthetic adsorption resin and the elution solvent are a first porous synthetic adsorption resin and a first elution solvent, respectively, and~~
- ~~wherein the eriocitrin separation from the citrus extract further comprises: bringing an eluate obtained by eluting the eriocitrin adsorbed on the first porous synthetic adsorption resin using the first elution solvent into contact with a second porous synthetic adsorption resin such that the eriocitrin in the eluate is adsorbed on the second porous synthetic adsorption resin, the second porous synthetic adsorption resin comprising at least one selected from styrenic and acrylic adsorption resins; and then eluting the eriocitrin adsorbed on the second porous synthetic~~

~~adsorption resin using a second elution solvent.~~

~~4. The eriocitrin-containing material according to claim 1 or 2, wherein the porous synthetic adsorption resin and the elution solvent are a first porous synthetic adsorption resin and a first elution solvent, respectively,~~

~~wherein the eriocitrin separation from the citrus extract further comprises: bringing the citrus extract into contact with a second porous synthetic adsorption resin such that eriocitrin in the citrus extract is adsorbed on the second porous synthetic adsorption resin, the second porous synthetic adsorption resin comprising at least one selected from styrenic and acrylic adsorption resins; and then eluting the eriocitrin adsorbed on the second porous synthetic adsorption resin using a second elution solvent, and~~

~~wherein the contact of the citrus extract with the first porous synthetic adsorption resin is carried out by bringing an eluate obtained by eluting the eriocitrin adsorbed on the second porous synthetic adsorption resin using the second elution solvent into contact with the first porous synthetic adsorption resin.~~

~~5. The eriocitrin-containing material according to claim 3 or 4, wherein the second elution solvent is hydrous ethanol having an ethanol concentration of 20 to 50% by volume.~~

~~6. A food or drink product characterized by comprising an eriocitrin-containing material according to any one of claims 1 to 5 formulated therein.~~

~~7. A pharmaceutical characterized by comprising an eriocitrin-containing material according to any one of claims 1 to 5 formulated therein.~~

~~8. A cosmetic characterized by comprising an eriocitrin-containing material according to any one of~~

~~claims 1 to 5 formulated therein.~~

1 9. A process for producing an eriocitrin-containing material, characterized by comprising the steps of:

5 preparing an eriocitrin-containing citrus extract from a citrus fruit using an extractant; and

separating eriocitrin from the citrus extract, wherein the step of separating eriocitrin from the citrus extract comprises the steps of:

10 bringing the citrus extract into contact with a porous synthetic adsorption resin such that eriocitrin in the citrus extract is adsorbed on the porous synthetic adsorption resin, the porous synthetic adsorption resin comprising a phenol-formaldehyde resin
15 as a main framework and having amino and phenolic hydroxyl groups; and

eluting the eriocitrin adsorbed on the porous synthetic adsorption resin using an elution solvent.

2 ~~10~~. The process for producing an eriocitrin-containing material according to claim 1 9, wherein the
20 elution solvent is hydrous ethanol having an ethanol concentration of 20 to 90% by volume.

3 ~~11~~. The process for producing an eriocitrin-containing material according to claim 1 9 or 2 ~~10~~,
25 wherein in the step of separating eriocitrin from the citrus extract, a step is repeated twice or more that involves brining the citrus extract into contact with the porous synthetic adsorption resin and subsequently eluting the eriocitrin adsorbed on the porous synthetic
30 adsorption resin.

4 ~~12~~. The process for producing an eriocitrin-containing material according to any one of claims 1 9 to 3 ~~11~~, wherein the porous synthetic adsorption resin and the elution solvent are a first porous synthetic
35 adsorption resin and a first elution solvent,

respectively, and wherein the step of separating eriocitrin from the citrus extract further comprises the steps of:

bringing an eluate obtained by eluting the
5 eriocitrin adsorbed on the first porous synthetic
adsorption resin using the first elution solvent into
contact with a second porous synthetic adsorption resin
such that the eriocitrin in the eluate is adsorbed on
the second porous synthetic adsorption resin, the
10 second porous synthetic adsorption resin comprising at
least one selected from styrenic and acrylic adsorption
resins; and

eluting the eriocitrin adsorbed on the second
porous synthetic adsorption resin using a second
15 elution solvent.

5 ~~13~~. The process for producing an eriocitrin-
containing material according to any one of claims 1 ~~9~~
to 3 ~~11~~, wherein the porous synthetic adsorption resin
and the elution solvent are a first porous synthetic
20 adsorption resin and a first elution solvent,
respectively, and wherein the step of separating
eriocitrin from the citrus extract further comprises
the steps of:

bringing the citrus extract into contact with a
25 second porous synthetic adsorption resin such that
eriocitrin in the citrus extract is adsorbed on the
second porous synthetic adsorption resin, the second
porous synthetic adsorption resin comprising at least
one selected from styrenic and acrylic adsorption
30 resins; and

eluting the eriocitrin adsorbed on the second
porous synthetic adsorption resin using a second
elution solvent,

wherein the step of bringing the citrus extract
35 into contact with the first porous synthetic adsorption

resin is carried out by bringing an eluate obtained by eluting the eriocitrin adsorbed on the second porous synthetic adsorption resin using the second elution solvent into contact with the first porous synthetic
5 adsorption resin.

6 14. The process for producing an eriocitrin-containing material according to claim 4 12 or 5 13, wherein the second elution solvent is hydrous ethanol having an ethanol concentration of 20 to 50% by volume.
10